

# PELCZYNSKI'S PROPERTY (V) ON SPACES OF VECTOR-VALUED FUNCTIONS

NARCISSE RANDRIANANTOANINA

Preliminary Version

**Abstract.** Let  $E$  be a separable Banach space and  $X$  be a compact Hausdorff space. It is shown that the space  $C(X; E)$  has property (V) if and only if  $E$  does. Similar result is also given for Bochner spaces  $L^p(X; E)$  if  $1 < p < \infty$  and  $\mu$  is a finite Borel measure on  $X$ .

## 1. Introduction

Let  $E$  and  $F$  be Banach spaces and suppose  $T : E \rightarrow F$  is a bounded linear operator. The operator  $T$  is said to be unconditionally converging if  $T$  does not  $\times$  any copy of  $c_0$ . A Banach space  $E$  is said to have Pelczynski's property (V) if every unconditionally converging operator with domain  $E$  is weakly compact. In a fundamental paper [14], Pelczynski showed that if  $X$  is a compact Hausdorff space then the space  $C(X)$ , of all continuous scalar valued functions on  $X$ , has property (V); and he asked ([14] Remark 1, p. 645; see also [9] p. 183) if for a Banach space  $E$  the abstract continuous function space  $C(X; E)$  has property (V). This question has been considered by several authors. Perhaps the sharpest result so far is in the paper of Cembranos, Kalton, E. Saab and P. Saab [5] where they proved that if  $E$  has property (U) and contains no copy of  $\ell_1$  then  $C(X; E)$  has property (V). There are however many known examples of Banach spaces that have property (V) but fail to satisfy the above conditions. For instance, Kisliakov in [12] (see also Dealban [7] independently) showed that the disk algebra has property (V); Bourgain did the same for ball algebras and polydisk algebras in [4] and  $H^1$  in [3]. For more information and examples of spaces with property (V), we refer to [11] and [17].

In this note, we obtain a positive answer to the above question for the separable case; namely we prove that if  $E$  is a separable Banach space then  $C(\cdot; E)$  has property (V) if and only if  $E$  does. We present also some applications of the main theorem to Banach spaces of compact operators as well as for Bochner function spaces.

Our notation is standard Banach space terminology as may be found in the books [8] and [9].

## 2. Definitions and some preliminary results

**Definition 1.** Let  $E$  be a Banach space. A series  $\sum_{n=1}^{\infty} x_n$  in  $E$  is said to be weakly unconditionally Cauchy (WUC) if for every  $x$  in  $E$ , the series  $\sum_{n=1}^{\infty} |x(x_n)|$  is convergent.

There are many criteria for a series to be a WUC series (see for instance [8]).

The following proposition was proved by Pelczynski in [14].

**Proposition 1.** For a Banach space  $E$ , the following assertions are equivalent:

- (i) A subset  $H \subset E$  is relatively weakly compact whenever  $\lim_{n \rightarrow \infty} \sup_{x \in H} \sum_{k=1}^n |x(x_k)| = 0$  for every WUC series  $\sum_{n=1}^{\infty} x_n$  in  $E$ ;
- (ii) For any Banach space  $F$ , every bounded operator  $T: E \rightarrow F$  that is unconditionally converging is weakly compact.

**Definition 2.** A subset  $H \subset E$  is called a (V)-subset if  $\lim_{n \rightarrow \infty} \sup_{x \in H} \sum_{k=1}^n |x(x_k)| = 0$  for every WUC series  $\sum_{n=1}^{\infty} x_n$  in  $E$ .

So a Banach space  $E$  has property (V) if (and only if) every (V)-subset of  $E$  is relatively weakly compact. This leads us to study (V)-subsets of the dual of  $C(\cdot; E)$  for a given Banach space  $E$  and a compact Hausdorff space  $\cdot$ .

Recall that the space  $C(\cdot; E)$  is isometrically isomorphic to the Banach space  $M(\cdot; E)$  of all weak\*-regular  $E$ -valued measures of bounded variation defined on the  $\sigma$ -field of Borel subsets of  $\cdot$  and equipped with norm  $\|m\| = \sum |j(m)|$ , where  $\sum |j(m)|$  is the variation of  $m$ . In this section we study different structures of subsets of  $M(\cdot; E)$ .

Let us begin by recalling some classical facts: Fix a probability measure on and let  $m \in M(\mathbb{R}; E)$  with  $\|m\|_1 = 1$  and  $\nu$  be a lifting of  $L^1(\mathbb{R})$  (see [10] and [20]). For  $x \in E$ , the scalar measure  $xm$  has density  $dxm = d \in L^1(\mathbb{R})$ . We define  $(m)(\cdot)(x) = (dxm = d)(\cdot)$ . It is well known that

$$x(m(A)) = \int_A h(m)(\cdot); xid(\cdot)$$

and

$$\|m\|_1(A) = \int_A \|h(m)(\cdot)\|_1 d(\cdot)$$

for every measurable subset  $A$  of  $\mathbb{R}$ . Note also that  $\|h(m)(\cdot)\|_1(\cdot) \in E$  is weak\*-scalarly measurable.

The following proposition can be deduced from [2] but we will present a direct proof for sake of completeness.

**Proposition 2.** Let  $H$  be a bounded subset of  $M(\mathbb{R}; E)$ . If  $H$  is a (V)-subset then  $V(H) = \{ \sum_{n=1}^{\infty} m_n \in H \}$  is relatively weakly compact in  $M(\mathbb{R}; E)$ .

**Proof.** Assume that  $V(H)$  is not relatively weakly compact. Since the space  $C(\mathbb{R})$  has property (V), there exists a WUC series  $\sum_{n=1}^{\infty} e_n$  in  $C(\mathbb{R})$ , sequence  $(m_n)_n$  in  $H$  and  $\epsilon > 0$  so that  $\|h_{m_n}; j_n\|_1 \geq \epsilon$  for each  $n \in \mathbb{N}$ . Let  $\nu = \sum_{n=1}^{\infty} \frac{1}{2^n} j_n$ . Since  $\|j_n\|_1 \leq 2^n$ , for a lifting  $\nu$  of  $L^1(\mathbb{R})$ , there exists a weak\*-scalarly measurable map  $g_n : \mathbb{R} \rightarrow E$  so that:

- (a)  $\|g_n(\cdot)\|_1 \leq 1$  and  $\|m_n\|_1(A) = \int_A \|g_n(\cdot)\|_1 d(\cdot)$  for all  $A \in \mathcal{B}$ ;
- (b)  $\int_A h_{m_n}(\cdot) d\nu = \int_A h_{g_n}(\cdot) d\nu$  for  $e \in E$  and  $A \in \mathcal{B}$ ;
- (c)  $\|g_n\|_1 = 1$ .

Now since  $C(\mathbb{R}; E)$  is norming for  $M(\mathbb{R}; E)$ , there exists  $\epsilon_n \in C(\mathbb{R}; E)$ , with  $\|e_n\|_1 = 1$  and such that  $\|h_{m_n}; e_n\|_1 \geq \epsilon_n$ , i.e.,

$$\int_{\mathbb{R}} \|h_{m_n}(\cdot); e_n(\cdot)\|_1 d\nu \geq \int_{\mathbb{R}} \|g_n(\cdot)\|_1 d\nu \geq \frac{\epsilon_n}{2}$$

or

$$\int_{\mathbb{R}} \|h_n(\cdot); g_n(\cdot)\|_1 d\nu \geq \int_{\mathbb{R}} \|g_n(\cdot)\|_1 d\nu \geq \frac{\epsilon_n}{2} :$$

Notice also that since  $\sum_{n \in \mathbb{N}} \|h_n\|_{\infty} < \infty$ ,  $\sum_{n \in \mathbb{N}} \|g_n\|_{\infty} < \infty$  and we get that

$$\begin{aligned} & \sum_{n \in \mathbb{N}} \|e_n\|_{\infty} \|h_n\|_{\infty} \|g_n\|_{\infty} < \infty \\ & = \sum_{n \in \mathbb{N}} \|e_n\|_{\infty} (\|h_n\|_{\infty} \|g_n\|_{\infty}) < \infty \\ & \sum_{n \in \mathbb{N}} \|e_n\|_{\infty} \|h_n\|_{\infty} \|g_n\|_{\infty} < \infty \end{aligned}$$

So for each  $n \in \mathbb{N}$ ,

$$\sum_{n \in \mathbb{N}} \|e_n\|_{\infty} \|h_n\|_{\infty} \|g_n\|_{\infty} < \infty$$

Fix  $f_n = e_n \otimes h_n$ ; the function  $f_n$  belongs to  $C(X; E)$  and we claim that  $\sum_{n=1}^{\infty} f_n$  is a WUC series in  $C(X; E)$ . For that it is enough to notice that for any finite subset of  $\mathbb{N}$ , we get

$$\sum_{n \in \mathbb{N}} \|e_n\|_{\infty} \|h_n\|_{\infty} < \infty$$

Now  $\sum_{n \in \mathbb{N}} \|h_n\|_{\infty} < \infty$ ,  $\sum_{n \in \mathbb{N}} \|g_n\|_{\infty} < \infty$ . Contradiction with the assumption that  $H$  is a  $(V)$ -subset.  $\square$

For the next proposition, we will use the following notation: for a given measure  $\mu \in M(X; E)$  and  $A \in \mathcal{A}$ ,  $\mu_A$  denotes the measure  $\mu_A(B) = \mu(A \cap B)$  for any  $B \in \mathcal{A}$ .

**Proposition 3.** Let  $H$  be a  $(V)$ -subset of  $M(X; E)$  and  $(A_n)_{n \in \mathbb{N}}$  a collection of measurable subsets of  $X$ . Then the subset  $\{\mu_{A_n} : \mu \in H\}$  is a  $(V)$ -subset of  $M(X; E)$ .

**Proof.** Assume that  $H$  is a  $(V)$ -subset of  $M(X; E)$ . By Proposition 3,  $V(H)$  is relatively weakly compact in  $M(X)$ . Let  $\nu$  be a control measure for  $V(H)$ . Fix a sequence  $(\mu_n)_{n \in \mathbb{N}}$  in  $\{\mu_{A_n} : \mu \in H\}$ . We need to show that the countable subset  $\{\mu_n : n \in \mathbb{N}\}$  is a  $(V)$ -subset. Let  $\sum_{n=1}^{\infty} f_n$  be a WUC series in  $C(X; E)$  with  $\sum_{n=1}^{\infty} \|f_n\|_{\infty} < \infty$ . For  $\varepsilon > 0$  (fixed), there exists  $\delta > 0$  such that if  $A \in \mathcal{A}$ ,  $\mu(A) < \delta$  then  $\sum_{n=1}^{\infty} \mu(f_n(A)) < \varepsilon$ ,  $\mu \in H$ ; for each  $n \in \mathbb{N}$ , choose a compact set  $C_n$  and an open set  $O_n$  such that  $C_n \subset A_n \subset O_n$  and  $\nu(O_n \setminus C_n) < \delta$ . Fix a continuous

$$\begin{aligned} \int_{\mathbb{M}_n(A_m)} f_n \, d\mu_n &= \int_{\mathbb{M}_n(A_m)} f_n \, d\mu_n \\ &= \int_{\mathbb{M}_n(A_m)} f_n \, d\mu_n \\ &\quad + \int_{\mathbb{M}_n(C_n)} f_n \, d\mu_n + \int_{\mathbb{M}_n(A_m \cap C_n)} f_n \, d\mu_n \\ &\quad + \int_{\mathbb{M}_n(O_n \cap C_n)} f_n \, d\mu_n + \int_{\mathbb{M}_n(A_m \cap C_n)} f_n \, d\mu_n \\ \int_n; m_n \, d\mu_n &+ \int_n; j(O_n \cap C_n) + \int_n; j(A_m \cap C_n) \\ \int_n; m_n \, d\mu_n &+ 2 \int_n; j(O_n \cap C_n) \\ \int_n; m_n \, d\mu_n &+ \dots \end{aligned}$$

If we denote by  $M^1(\cdot; E)$  the set of  $M(\cdot; E)$  in  $J$  then we obtain the following corollary.

**Proof.** Let  $g_m : \mathbb{R}_+$  be the density of  $m$  with respect to  $\lambda$ .

$$\lim_{N \rightarrow \infty} \int_{\mathbb{R}^d} g_N(x) dx = 0 \text{ uniformly on } H :$$

$$\sum_{f \in \mathcal{F}_m} g_m(f) d(f) < \infty$$

and let  $A_m = f!; g_m(!)$  N g. It is clear that  $H_* = \text{fin } A_m; m \in \mathbb{N}$  is a subset of  $N M^{-1} ( ; E )$  and is a  $(V)$ -subset by Proposition 4. Also each measure  $m$  in  $H$  satisfies  $m = m_{A_m} + m_{A_m^c}$  with  $m_{A_m^c} k < \infty$ .  $\square$

Our next proposition can be viewed as a generalization of Theorem 1 of [15] for sequences of weak\*-scalarly measurable maps. We denote by  $(e_n)$  the unit vector basis of  $C_0$ ,  $(\Omega, \mathcal{F}, \mu)$  a probability space and for any Banach space  $F$ ,  $F_1$  stands for the closed unit ball of  $F$ .

**Proposition 4.** Let  $Z$  be a separable subspace of a real Banach space  $E$  and  $(f_n)_n$  be a sequence of maps from  $\Omega$  to  $E$  that are weak\*-scalarly measurable with  $\sup_n \|f_n\|_{k_1} < \infty$ . Let  $a, b$  be real numbers with  $a < b$  then:

There exist a sequence  $g_n \in \text{conv}\{f_n, f_{n+1}, \dots\}$ , measurable subsets  $C$  and  $L$  of  $\Omega$  with  $\mu(C \cup L) = 1$  such that

(i)  $\Omega \subset C$  and  $\Omega \subset L$  ( $C_0; Z$ ) $_1$  then either

$$\limsup_{n \rightarrow \infty} \int_{\Omega} g_n(\omega) d\mu(\omega) = b \text{ or } \\ \liminf_{n \rightarrow \infty} \int_{\Omega} g_n(\omega) d\mu(\omega) = a;$$

(ii)  $\Omega \subset L$ , there exists  $k \in \mathbb{N}$  so that for each infinite sequence  $(\epsilon_n)$  of zeros and ones, there exists  $T \in L$  ( $C_0; Z$ ) $_1$  such that for  $n \geq k$ ,

$$\epsilon_n = 1 \Rightarrow \int_{\Omega} g_n(\omega) d\mu(\omega) = b \\ \epsilon_n = 0 \Rightarrow \int_{\Omega} g_n(\omega) d\mu(\omega) = a;$$

The proof is a further extension of the techniques used in [19] and [15]. We will begin by introducing some notations, some of which were already used in [19] and [15].

Let  $f_n : \Omega \rightarrow E$  be a sequence as in the statement of the proposition. We write  $u_n = f_n$  (or  $(u_n) = (f_n)$ ) if there exists  $k \in \mathbb{N}$  and  $p_1 < q_1 < p_2 < q_2 < \dots < p_n < q_n < \dots$  so that for  $n \geq k$ ,

$$u_n = \sum_{i=p_n}^{q_n} f_i; \text{ with } f_i \in [0, 1] \text{ and } \sum_{i=p_n}^{q_n} f_i = 1;$$

Consider  $L(C_0; Z)_1$  the closed unit ball of  $L(C_0; Z)$  with the strong operator topology. It is not difficult to see (using the fact that  $Z$  is separable) that  $L(C_0; Z)_1$  is a Polish space; in particular it has a countable basis  $(O_n)_n$ . Since  $L(C_0; Z)_1$  is a metric space, we can assume that the  $O_n$ 's are open balls.

The letter  $K$  will stand for the set of all (strongly) closed subsets of  $L(C_0; Z)_1$ .

We will say that  $\mathbb{T} \cap K(\cdot)(\cdot) \cap K$  is measurable if the set  $f : K(\cdot) \setminus O_n \in \mathcal{G}$  is a measurable subset of  $\mathbb{T}$  for every  $n \in \mathbb{N}$ .

Let  $h_n = \sum_{i=p_n}^{p_{n+1}} f_i$  with  $\sum_{i=p_n}^{p_{n+1}} 1 = 1$ ,  $f_i \geq 0$  and  $p_1 < p_2 < p_3 < \dots$ ;  $V$  be an open subset of  $L(C_0; \mathbb{Z})_1$  and  $\mathbb{T} \cap K(\cdot)$  a fixed measurable map, we set

$$(1) \quad \overline{h_n}(\cdot) = \sup_{k \in \mathcal{G}_n} \sup_{f \in h_n(\cdot)} f; T \in_k i; T \in V \setminus K(\cdot) \mathcal{G}$$

$$(2) \quad (h)(\cdot) = \lim_{n \rightarrow \infty} \sup \overline{h_n}(\cdot)$$

Notice that the definition of  $\overline{h_n}$  depends on the representation of  $h_n$  as a block convex combination of  $f_n$ 's. Similarly we set

$$(3) \quad \mathfrak{h}_n(\cdot) = \inf_{k \in \mathcal{G}_n} \inf_{f \in h_n(\cdot)} f; T \in_k i; T \in V \setminus K(\cdot) \mathcal{G}$$

$$(4) \quad (h)(\cdot) = \lim_{n \rightarrow \infty} \inf \mathfrak{h}_n(\cdot) :$$

The proof of the following lemma is just a notational adjustment of the proof of Lemma 2 of [15].

Lemma 1. There exists  $(g_n) \in (f_n)$  such that if  $(h_n) \in (g_n)$  then

$$\lim_{n \rightarrow \infty} k(g) \overline{h_n} k_1 = 0 \text{ and } \lim_{n \rightarrow \infty} k'(g) \mathfrak{h}_n k_1 = 0.$$

Main construction:

Fix  $a < b$  and let  $\omega$  be the first uncountable ordinal. Set  $h_n^0 = f_n$ , we construct as in [15] for  $\alpha < \omega$ , sequences  $h = (h_n)_n$ , measurable maps  $K : \mathbb{T} \cap K$  with the following properties:

$$(5) \quad \text{for } \alpha < \omega; \quad h = h^0;$$

For  $\alpha < \omega$  and  $h = f$  with  $h = \sum_{j=p_n}^{p_{n+1}} f_j$  we define

$$(6) \quad \overline{h_{n,\alpha}}(\cdot) = \sup_{k \in \mathcal{G}_n} \sup_{f \in h_n(\cdot)} f; T \in_k i; T \in O \setminus K(\cdot) \mathcal{G}$$

$$; (h)(\cdot) = \lim_{n \rightarrow \infty} \sup \overline{h_{n,\alpha}}(\cdot)$$

$$\mathfrak{h}_{n,\alpha}(\cdot) = \inf_{k \in \mathcal{G}_n} \inf_{f \in h_n(\cdot)} f; T \in_k i; T \in O \setminus K(\cdot) \mathcal{G}$$

$$; (h)(\cdot) = \lim_{n \rightarrow \infty} \inf \mathfrak{h}_{n,\alpha}(\cdot) :$$

Then for each  $\alpha$  of the form  $\beta + 1$  and each  $h \in h_\alpha$ , we have

$$\lim_{n \rightarrow 1} k_n(\alpha, h) = \bar{h}_\alpha; k_1 = 0$$

and

$$\lim_{n \rightarrow 1} k'_n(\alpha, h) = \bar{h}_\alpha; k_1 = 0 :$$

If  $\alpha$  is limit, we set

$$(7) \quad K(\alpha) = \bigcup_{\beta < \alpha} K(\beta) :$$

If  $\alpha = \beta + 1$

$$(8) \quad K(\alpha) = fT \cap K(\beta); T \cap O(\alpha) \cap h_\alpha(\alpha) \cap b; \alpha, h_\alpha(\alpha) \cap a_g :$$

The construction is done in the same manner as in [15] and is a direct application of Lemma 1.

As in [15], one can fix an ordinal  $\alpha < \omega_1$  such that for a.e.  $\alpha \in \omega_1$ ,  $K(\alpha) = K_{\alpha+1}(\alpha) :$

Let  $h = h_{\alpha+1}$ ,

$$C = f! : K(\alpha) = ; g \text{ and}$$

$$M = f! : K(\alpha) = K_{\alpha+1}(\alpha) \notin ; g :$$

Clearly  $C$  and  $M$  are measurable and  $(C \cap L) = 1$ .

The next lemma is the analogue of Lemma 4 of [15].

Lemma 2. Let  $\alpha \in \omega_1$  and  $T \in L(C_0; Z)_1$ . If  $u \in h$  then either

$$\limsup_{n \rightarrow 1} u_n(\alpha); T e_n = b \text{ or}$$

$$\liminf_{n \rightarrow 1} u_n(\alpha); T e_n = a$$

Proof. Let  $\alpha \in \omega_1$ ,  $T \in L(C_0; Z)_1$  and  $x \in u \in h$  (say  $u = \sum_{j=1}^{\infty} f_j$ ); let  $S : C_0 \rightarrow C_0$  be an operator defined as follows  $Se_{b_n} = e_n$  and  $Se_j = 0$  if  $j \notin b_n$ ,  $n \in \mathbb{N}$ . The operator  $S$  is obviously bounded linear with  $\|S\| = 1$ . So  $T \in L(C_0; Z)_1 = K_0(\alpha)$ . Since  $T \in S \notin K(\alpha)$ , there exists a least ordinal  $\beta$  for which  $T \in S \notin K(\beta)$ . The ordinal  $\beta$  cannot be a limit so  $\beta = \gamma + 1$  and  $T \in S \notin K(\gamma)$ . By



the definition of  $K(\cdot)$ , there exists  $\delta \in \mathbb{N}$  with  $T \in S_2$ . But either  $\delta_j(h)(\cdot) \leq b$  or  $\delta_j(h)(\cdot) \geq a$ . Now since  $u \leq h$ , we get that either

$$\limsup_{n \rightarrow \infty} \sup_{i \in I} (h_n(\cdot); T e_n i) = \limsup_{n \rightarrow \infty} \sup_{i \in I} (h_n(\cdot); T e_n i) \leq S_{\mathbb{Q}} i$$

$$\delta_j(u)(\cdot) \leq \delta_j(h)(\cdot) \leq b$$

or

$$\liminf_{n \rightarrow \infty} \inf_{i \in I} (h_n(\cdot); T e_n i) \geq \delta_j(u)(\cdot) \geq \delta_j(h)(\cdot) \geq a :$$

The lemma is proved.  $\square$

The following property of the measurable subset  $M$  is somewhat stronger than that obtain in Lemma 5 of [15] and is the main adjustment of the entire proof.

**Lemma 3.** There exists a subsequence  $(n(i))$  of integers such that for a.e.  $\delta \in M$ , if  $\delta$  is an infinite sequence of zeros and ones then there exists an operator  $T \in L(C_0; Z)_1$  (which may depend on  $\delta$  and  $\delta$ ) such that:

$$\delta_i = 1 \Rightarrow \|h_{n(i)}(\cdot); T e_i\| \leq b$$

$$\delta_i = 0 \Rightarrow \|h_{n(i)}(\cdot); T e_i\| \geq a :$$

**Proof.** Let us denote by  $F$  the set of finite sequences of zeros and ones and  $F^\infty$  the set of infinite sequences of zeros and ones. For  $s \in F$ ,  $|s|$  will denote the length of  $s$ . Let  $s = (s_1; \dots; s_n)$  and  $r = (r_1; \dots; r_m)$  with  $n \leq m$ . We say that  $s < r$  if  $s_i = r_i$  for  $i \leq m$ . Let us fix a representation of  $(h_n)$  as block convex combination of  $(f_n)$ :

$$h_n = \sum_{i=p_n}^{q_n} \lambda_i f_i :$$

We will construct sequences of integers  $n(i)$  and  $m(i)$ ; measurable sets  $B_i \subset M$  and measurable maps  $Q(s; \cdot) : M \rightarrow \mathbb{N}$  (for  $s \in F$ ) such that:

- (9)  $q_{n(1)} < m(1) < q_{n(2)} < m(2) < \dots < q_{n(i)} < m(i) < \dots$  ;
- (10)  $\delta \in F^\infty$  ;  $\sup \{Q(s; \delta) ; \delta \in M\} < 1$  ;
- (11)  $(M \cap B_i) \subset \delta^i$  ;

$$(12) \quad \text{For all } s \in F; \text{diam}(O_{Q(s;!)}) \leq \frac{1}{j};$$

$$(13) \quad \text{For } s, r \in F; s < r \text{ and} \\ \bigcup_{j \leq i \leq j} B_i; \text{ one has } O_{Q(r;!)} \subset O_{Q(s;!)};$$

$$(14) \quad \exists! 2^M; s \in F; K \setminus O_{Q(s;!)} \neq \emptyset;$$

$$(15) \quad \exists s \in F; \exists i \in \mathbb{N} \text{ p. } j; \exists! 2^{\bigcup_{i \leq j \leq i} B_j}; \\ s_i = 1 \Rightarrow \exists T \in O_{Q(s;!)}; \sup_{q_n(i)} h_{n(i)}(!); T e_k i = b \\ s_i = 0 \Rightarrow \exists T \in O_{Q(s;!)}; \inf_{q_n(i)} h_{n(i)}(!); T e_k i = a:$$

The construction is done in a similar fashion as in [15]; the only difference is on the selection of the measurable map  $Q(s; !): \mathbb{N} \rightarrow \mathbb{N}$  so that (12) is satisfied. For that we consider instead of  $N$ , the subset  $M \subset N$  defined by

$$M = \{k \in N; \text{diam}(O_k) \leq \frac{1}{j}\}$$

and since  $N$  and  $M$  are equipped with the discrete topology, we can replace  $N$  by  $M$  and use the same argument to get  $Q(s; !): \mathbb{N} \rightarrow M$  f0g.

To complete the proof, let  $L = \bigcup_{k \in \mathbb{N}} \bigcap_{i \in \mathbb{N}} B_i$ . It is clear that  $(M \cap L) = \emptyset$ . Fix  $\bigcup_{i \in \mathbb{N}} B_i$  and  $\bigcup_{i \in \mathbb{N}} F^1$ , denote  $\mathbf{m} = (m_i)_{i \in \mathbb{N}}$ . Let

$$\mathbf{m} = (m_1; \dots; m_m) \in F; \exists m \in \mathbb{N};$$

By (15), we get that for  $m \in \mathbb{N}$  and  $\exists i \in \mathbb{N}; \exists! 2^{\bigcup_{i \leq j \leq m} B_j}$ ,

$$s_i = 1 \Rightarrow \exists T \in O_{Q(\mathbf{m};!)}; \sup_{q_n(i)} h_{n(i)}(!); T e_k i = b \\ s_i = 0 \Rightarrow \exists T \in O_{Q(\mathbf{m};!)}; \inf_{q_n(i)} h_{n(i)}(!); T e_k i = a:$$

It is easy to check that the same conclusion holds for  $T \in \overline{O_{Q(\mathbf{m};!)}}$  (the closure of  $O_{Q(\mathbf{m};!)}$  for the strong operator topology). So if we let  $A = \bigcap_{m \in \mathbb{N}} \overline{O_{Q(\mathbf{m};!)}}$ ,  $A \neq \emptyset$ ; in fact  $(\overline{O_{Q(\mathbf{m};!)}})_{m \in \mathbb{N}}$  is a nested sequence of nonempty closed sets (by (13)) of a complete metric space and such that  $\text{diam}(\overline{O_{Q(\mathbf{m};!)}}) \rightarrow 0$  ( $m \rightarrow \infty$ ) (by (12)) so  $A \neq \emptyset$ ; (see for instance [13] p.270).

It is now clear that if  $\sum_{i=1}^{\infty} \|B_i\| < \infty$  and  $T \in A$ , for  $i \leq k$ ,

$$\begin{aligned} i = 1 & \Rightarrow \sup_{q_n(i) \leq k \leq m(i)} \|h_n(i)(!); T e_k\| = \|b\| \\ i = 0 & \Rightarrow \sup_{q_n(i) \leq k \leq m(i)} \|h_n(i)(!); T e_k\| = \|a\| : \end{aligned}$$

We complete the proof as in [15]: choose  $k(i) \in [q_n(i), m(i)]$  such that

$$\sup_{q_n(i) \leq k \leq m(i)} \|h_n(i)(!); T e_k\| = \|h_n(i)(!); T e_{k(i)}\|$$

for  $i = 1$  and

$$\inf_{q_n(i) \leq k \leq m(i)} \|h_n(i)(!); T e_k\| = \|h_n(i)(!); T e_{k(i)}\|$$

for  $i = 0$ . The sequence  $(k(i))$  is an increasing sequence by (9) so one can construct an operator  $S : c_0 \rightarrow c_0$  with  $S e_i = e_{k(i)}$   $\forall i \in \mathbb{N}$  and it is now clear that if

$$\begin{aligned} i = 1 & \Rightarrow \|h_n(i)(!); T S e_i\| = \|b\| \\ i = 0 & \Rightarrow \|h_n(i)(!); T S e_i\| = \|a\| : \end{aligned}$$

The proof of the lemma is complete. For the proposition, we take  $g_i = h_n(i)$  for  $i \in \mathbb{N}$ .  $\square$

### 3. Main Theorem

**Theorem 1.** Let  $E$  be a separable Banach space and  $M$  be a compact Hausdorff space. Then the space  $C(M; E)$  has property (V) if and only if  $E$  has property (V).

**Proof.** If  $C(M; E)$  has property (V), then the space  $E$  has property (V) since  $E$  is isomorphic to a complemented subspace of  $C(M; E)$ . Conversely, assume that  $E$  has property (V). Let  $H$  be a  $(V)$ -subset of  $M(M; E)$ . Our goal is to show that  $H$  is relatively weakly compact. Using Corollary 1, one can assume without loss of generality that there exists a probability measure  $\mu$  on  $M$  such that  $\int_M f d\mu = 0$  for each  $f \in H$ . Observe that if  $E$  has property (V), then  $E$  is weakly sequentially complete and thus  $M(M; E)$  is weakly sequentially complete as shown in [19] (Theorem 17). If  $H$  is not relatively weakly compact, then it contains a sequence  $(m_n)_n$  that is equivalent to the  $\ell^1$ -basis. By Theorem 14 of [19], there exists

$m_n^0 \in \text{conv} \{m_n; m_{n+1}; \dots; g\}$  and  $\epsilon_n^0 > 0$  so that for  $n \geq n_0$ , there exists  $N \in \mathbb{N}$  such that  $(m_n^0(\cdot))_{n \geq N}$  is equivalent to the  $\mathcal{H}$ -basis in  $E$ . Let

$$f_n = (m_n^0(\cdot))_{n \geq N}; \quad n \in \mathbb{N} :$$

$(f_n)_n$  is a sequence of weak\*-scalarly measurable maps and  $\sup_n \|f_n\|_{k_1} < 1$ .

**Proposition 5.** There exist a sequence  $g_n \in \text{conv} \{f_n; f_{n+1}; \dots; g\}$ , a positive number and a strongly measurable map  $T : \mathbb{R} \rightarrow L(C_0; E)_1$  such that

$$\lim_{n \rightarrow \infty} \inf_{i \in \mathbb{N}} \langle g_n(\cdot); T(\cdot)e_n \rangle = 0 :$$

For the proof of the proposition, let  $(a(k); b(k))_{k \in \mathbb{N}}$  be an enumeration of all pairs of rationals with  $a < b$ . By induction, we construct sequences  $(g^k)$ , measurable sets  $C_k, L_k$  of  $\mathcal{H}$  satisfying the following:

- (i)  $g^{k+1} = g^k$  for each  $k \in \mathbb{N}$ ;
- (ii)  $C_{k+1} \subset C_k, L_k \subset L_{k+1}, (C_k \cap L_k) = \emptyset$
- (iii)  $\exists n \in \mathbb{N}, n \geq k$ , and  $T \in L(C_0; E)_1$ , then either

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{i \in \mathbb{N}} \langle g_n^j(\cdot); T(\cdot)e_n \rangle &= b(k) \\ \lim_{n \rightarrow \infty} \inf_{i \in \mathbb{N}} \langle g_n^j(\cdot); T(\cdot)e_n \rangle &= a(k); \end{aligned}$$

- (iv)  $\exists n \in \mathbb{N}, n \geq k$ , there exists  $j \in \mathbb{N}$  such that for each infinite sequence of zeros and ones, there exists  $T \in L(C_0; E)_1$  such that if  $n_j = 1$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{i \in \mathbb{N}} \langle g_n^k(\cdot); T(\cdot)e_n \rangle &= b(k) \text{ or} \\ \lim_{n \rightarrow \infty} \inf_{i \in \mathbb{N}} \langle g_n^k(\cdot); T(\cdot)e_n \rangle &= a(k); \end{aligned}$$

This is just an application of Proposition 4 inductively starting from  $g^0 = f$ .

Let  $P = \{k \in \mathbb{N}; b(k) > 0\}$  and  $N = \{k \in \mathbb{N}; a(k) < 0\}$ . It is clear that  $\mathbb{N} = P \cup N$ . Consider  $C = \bigcup_{k \in P} C_k$  and  $L = \bigcup_{k \in N} L_k$ ; we have  $(C \cap L) = \emptyset$ .

**Case 1:**  $(L) > 0$

Since  $L = \bigcup_{k \in N} L_k$ , there exists  $k \in \mathbb{N}$  such that  $(L_k) > 0$ . Let  $(g_n) = (g_n^k)$ . If  $k \in P$  (i.e.,  $b(k) > 0$ )  $x = (1; 1; 1; \dots)$ . For each  $n \in \mathbb{N}$ , there exists  $T \in L(C_0; E)_1$  such that  $\langle g_n^k(\cdot); T(\cdot)e_n \rangle = b(k) - \epsilon_n$ . We can choose the above operator measurably using the following lemma:

**Lemma 4.** There exists a strongly measurable map  $T : \mathbb{R} \rightarrow L(C_0; E)_1$  such that:

$$(\cdot) T(!) = 0 \text{ if } ! \notin L_k;$$

$$(\cdot) ! \in L_k, \text{ there exists } j \in \mathbb{N} \text{ such that if } n \geq j, h_{g_n}(!); T(!) e_n \leq b(k).$$

To see the lemma, consider  $L(c_0; E)_1$  with the strong operator topology and  $E_1$  with the weak\*-topology. The space  $E_1$  is a compact metric space and hence is a Polish space. The space  $E_1^{\mathbb{N}} = L(g; E)_1$  equipped with the product topology is a Polish space. Let  $A$  be the following subset of  $E_1^{\mathbb{N}} = L(g; E)_1$ :

$$f(x_n); Tg \in A, \text{ if } j \in \mathbb{N}; h_{x_n}; T e_n \leq b(k) \text{ for } n \geq j;$$

The set  $A$  is clearly a Borel subset of  $E_1^{\mathbb{N}} = L(g; E)_1$  and if  $\pi: E_1^{\mathbb{N}} = L(g; E)_1 \rightarrow E_1$  is the first projection,  $\pi(A)$  is an analytic subset of  $E_1$ . By Theorem 8.5.3 of [6], there exists a universally measurable map  $\phi: \pi(A) \rightarrow L(c_0; E)_1$  such that the graph of  $\phi$  is a subset of  $A$ . Notice that if  $! \in L_k$ ,  $(g_n(!))_{n \geq 1} \in \pi(A)$ . We define

$$T(!) = \begin{cases} (g_n(!))_{n \geq 1} & \text{if } ! \in L_k \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to check that  $T$  satisfies all the requirements of the lemma. The lemma is proved.

Back to the proof of the proposition, we have  $h_{g_n}(!); T(!) e_n \leq b(k)$  for  $! \in L_k$ , and  $n \geq j$ ; so  $\liminf_{n \rightarrow \infty} h_{g_n}(!); T(!) e_n \leq b(k)$  for  $! \in L_k$ , and by Fatou's lemma,

$$\liminf_{n \rightarrow \infty} h_{g_n}(!); T(!) e_n \leq b(k) \quad (I_k)$$

so if  $b(k) \cdot (I_k) > 0$ , the proof is complete for  $k \in \mathbb{P}$ .

Now if  $k \in \mathbb{N}$ , we consider  $\phi = (0; 0; \dots)$  and choose a strongly measurable map  $\psi: \pi(T(!))$  (using similar argument as in the above lemma) with  $T(!) = 0$  for  $! \notin L_k$  and for  $! \in L_k$ , there exists  $j \in \mathbb{N}$  such that  $h_{g_n}(!); T(!) e_n \leq a(k) < 0$  for  $n \geq j$ . So we get that

$$\limsup_{n \rightarrow \infty} h_{g_n}(!); T(!) e_n \leq a(k)$$

for each  $! \in L_k$  and hence

$$\limsup_{n \rightarrow \infty} h_{g_n}(!); T(!) e_n \leq a(k) \quad (I_k)$$

which implies that

$$\liminf_{n \rightarrow \infty} h_{g_n}(!); T(!) e_n \leq -a(k) \cdot (I_k)$$

so the proof is complete if  $\|L\| > 0$ .

Case 2:  $\|L\| = 0$

Since  $\|C\| \|L\| = 1$ , we have that  $\|C_n\| = 0$ . Choose a sequence  $(g_n)$  so that  $(g_n) \subset (g_n^k)$  for every  $k \in \mathbb{N}$ . By the definition of the  $C_k$ 's and (iii) we have either  $\limsup_{n \rightarrow \infty} \|g_n\|_{E_n} > 0$  or  $\liminf_{n \rightarrow \infty} \|g_n\|_{E_n} = 0$  for  $k \in \mathbb{N}$ , and therefore for each  $\varepsilon > 0$ ,  $\lim_{n \rightarrow \infty} \|g_n\|_{E_n} > \varepsilon$  exists for every  $T \in L(C_0; E)_1$  (\*).

But for  $\varepsilon > 0$ , the sequence  $(f_n(\varepsilon))_n$  is equivalent to the  $\varepsilon$ -basis in  $E$  and since  $(g_n) \subset (f_n)$ ,  $(g_n(\varepsilon))_n$  is also equivalent to the  $\varepsilon$ -basis in  $E$  and therefore  $\{g_n(\varepsilon); n \in \mathbb{N}\}$  cannot be a  $(V)$ -subset of  $E$ , i.e., there exists an operator  $T \in L(C_0; E)_1$  such that  $\limsup_{n \rightarrow \infty} \|g_n(\varepsilon)\|_{E_n} > 0$ ; but condition ( ) insures that the limit exists so for each  $\varepsilon > 0$ , there exists  $T \in L(C_0; E)$  such that  $\lim_{n \rightarrow \infty} \|g_n(\varepsilon)\|_{E_n} > 0$ . We now choose the operator  $T$  measurably using the same argument as in the above lemma: i.e., there exists  $T : \mathbb{T} \rightarrow L(C_0; E)_1$  strongly measurable such that  $T(\varepsilon) = 0$  for  $\varepsilon \neq 0$  and  $\lim_{n \rightarrow \infty} \|g_n(\varepsilon)\|_{E_n} > 0$  for  $\varepsilon > 0$ . Let  $\phi(\varepsilon) = \lim_{n \rightarrow \infty} \|g_n(\varepsilon)\|_{E_n}$  for  $\varepsilon > 0$  and 0 otherwise.

The map  $\varepsilon \mapsto \phi(\varepsilon)$  is measurable and we obtain

$$\lim_{n \rightarrow \infty} \int_{\mathbb{T}} \|g_n(\varepsilon)\|_{E_n} d\mu(\varepsilon) = \int_{\mathbb{T}} \phi(\varepsilon) d\mu(\varepsilon) = \int_{\mathbb{T}} \phi(\varepsilon) d\mu(\varepsilon) > 0 :$$

The proof of the proposition is complete.

To complete the proof of the theorem, let  $(g_n) \subset (f_n)$ ,  $T : \mathbb{T} \rightarrow L(C_0; E)_1$  strongly measurable and  $\phi > 0$  as in Proposition 5. For each  $n \in \mathbb{N}$ , let  $G_n : \mathbb{T} \rightarrow E$  be the measure in  $M(\mathbb{T}; E)$  defined by:

$$G_n(A) = \int_A \langle g_n, T(\varepsilon) \rangle d\mu(\varepsilon) :$$

It is clear that  $G_n \in \text{conv} \{m_n, m_{n+1}, \dots\}$  and we will show that  $\{G_n; n \in \mathbb{N}\}$  is not a  $(V)$ -subset of  $M(\mathbb{T}; E)$  to get a contradiction by virtue of Proposition 3: Since  $\varepsilon \mapsto T(\varepsilon)e_n$  is norm measurable for each  $n \in \mathbb{N}$ , one can choose (using Lusin's Theorem) a compact subset  $K_n$  with  $\mu(K_n^c) < \frac{1}{3}$  and such that the map  $\varepsilon \mapsto T(\varepsilon)e_n$  ( $\varepsilon \in K_n$ ) is continuous for each  $n \in \mathbb{N}$ .

Let  $\psi : C(\mathbb{T}; E) \rightarrow C(\mathbb{T}; E)$  be an extension operator (the existence of such operator is given by Theorem 21.1.4 of [18]) and consider  $t_n = \psi(T(\cdot)e_n)$ . The series  $\sum_{n=1}^{\infty} t_n$  is a WUC series in  $C(\mathbb{T}; E)$ . In fact the operator  $S : C_0 \rightarrow C(\mathbb{T}; E)$

given by  $S e_j = T(e_j)$  is easily checked to be linear and bounded and  $t_n = S(e_n)$  so  $\sum_{n=1}^{\infty} t_n$  is a WUC series.

The following estimate concludes the proof.

$$\begin{aligned} \| \sum_{n=1}^N t_n \|_{G_N} &= \| \sum_{n=1}^N h_{g_n}(e_n); t_n(e_n) \text{id}(e_n) \| \\ &= \left\| \sum_{n=1}^N h_{g_n}(e_n); T(e_n) e_n \text{id}(e_n) \right\| + \left\| \sum_{n=1}^N h_{g_n}(e_n); t_n(e_n) \text{id}(e_n) \right\| \end{aligned}$$

so

$$\begin{aligned} \| \sum_{n=1}^N t_n \|_{G_N} &\leq \left\| \sum_{n=1}^N h_{g_n}(e_n); T(e_n) e_n \text{id}(e_n) \right\| + \left\| \sum_{n=1}^N h_{g_n}(e_n); t_n(e_n) \text{id}(e_n) \right\| \\ &= \left\| \sum_{n=1}^N h_{g_n}(e_n); t_n(e_n) \text{id}(e_n) \right\| + \left\| \sum_{n=1}^N h_{g_n}(e_n); T(e_n) e_n \text{id}(e_n) \right\| \\ &\leq \| \sum_{n=1}^N h_{g_n}(e_n); T(e_n) e_n \text{id}(e_n) \| + \frac{2}{3} \end{aligned}$$

which implies that

$$\left\| \sum_{n=1}^N h_{g_n}(e_n); T(e_n) e_n \text{id}(e_n) \right\| \leq \frac{2}{3} + \| \sum_{n=1}^N t_n \|_{G_N}.$$

Hence

$$\liminf_{n \rightarrow \infty} \| \sum_{n=1}^n t_n \|_{G_n} \leq \frac{2}{3}.$$

This of course shows that  $\{ \sum_{n=1}^n t_n \}_{n \in \mathbb{N}}$  is not a (V)-set. The theorem is proved.  $\square$

Theorem 1. above has the following consequences relative to Banach spaces of compact operators. For what follows if  $X$  and  $Y$  are Banach spaces,  $K_w(X; Y)$  denotes the Banach space of weak\* to weakly continuous compact operators from  $X$  to  $Y$  equipped with the operator norm. We have the following corollaries.

Corollary 2. Let  $X$  and  $Y$  be Banach spaces. If  $X$  is injective and  $Y$  is separable and has property (V) then  $K_w(X; Y)$  has property (V).

Proof. The space  $K_w(X; Y)$  is isometrically isomorphic to  $K_w(Y; X)$  which is a complemented subspace of  $K_w(Y; C(B_X)) = C(B_X; Y)$  has property (V) by Theorem 1.  $\square$

Corollary 3. Let  $X$  be a  $L_1$ -space and  $Y$  a separable Banach space with property (V). The space  $K(X; Y)$  has property (V).

Proof. The space  $K(X; Y)$  is isomorphic to  $K_w(X; Y)$  (see [16]) and it is well known that  $X$  is injective and so  $K_w(X; Y)$  has property (V) by Corollary 2.  $\square$

We now turn our attention to Bochner spaces. In [1], Bombal observed that if  $E$  is a closed subspace of an order continuous Banach lattice, then  $L^p(\cdot; E)$  has property (V) if  $1 < p < \infty$  and  $E$  has property (V). Our next result shows that for the separable case, property (V) can be lifted to the Bochner space  $L^p(\cdot; E)$ .

Theorem 2. Let  $E$  be a separable Banach space and  $(\cdot; \cdot)$  be a finite measure space. If  $1 < p < \infty$ , then the space  $L^p(\cdot; E)$  has property (V) if and only if  $E$  does.

Proof. Without loss of generality, we will assume that  $\cdot$  is a compact Hausdorff space,  $\cdot$  is a Borel measure and  $\cdot$  is the completion of the field of Borel measurable subsets of  $\cdot$ . For  $1 < p < \infty$ , let  $q$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ . It is a well known fact that the dual of  $L^p(\cdot; E)$  is isometrically isomorphic to the space  $M^q(\cdot; E')$  of all vector measures  $F : \Sigma \rightarrow E'$  with

$$\|F\|_{M^q} = \sup_{A \in \Sigma} \left( \sum_{i=1}^{\infty} \|F(A_i)\|_{E'}^q \right)^{1/q} < \infty$$

(see for instance, [9], p. 115).

Let  $H$  be a (V)-subset of  $M^q(\cdot; E')$  and assume that  $H \not\subset M^1(\cdot; E')$ . Since  $C(\cdot; E) \subset L^p(\cdot; E)$  and  $C(\cdot; E)$  has property (V) by Theorem 1,  $H$  is relatively weakly compact in  $M^q(\cdot; E')$ . Let  $(f_n)_n \subset H$  and  $\mu$  a lifting of  $\mu$ . There exists  $G_n \in \text{conv}\{f_n, f_{n+1}, \dots\}$  and  $G \in M^1(\cdot; E')$  such that  $\|G_n(\cdot)\|_{E'} - \|G(\cdot)\|_{E'}$  converges to zero for  $\mu$ -a.e.  $\cdot$ . By the Lebesgue dominated convergence

$$\sum_{n=1}^{\infty} \|G_n(\cdot)\|_{E'} - \|G(\cdot)\|_{E'} \leq \sum_{n=1}^{\infty} \|G_n(\cdot)\|_{E'} < \infty$$

But this is equivalent to say that  $\|G_n - G\|_{M^1} \rightarrow 0$  which proves that  $H$  is relatively weakly compact in  $M^q(\cdot; E')$  (see for instance [21]). The theorem is proved.  $\square$

Remark 1. As it was observed in [17], the property (V) cannot be lifted from  $E$  to  $L^1(\cdot; E)$ . In fact the space  $E = (c_0^n)_1$  has property (v) but  $L^1(\cdot; E)$  contains a complemented copy of  $c_0$  hence failing property (v).



## References

1. F. Bombal. On  $\ell_1$  subspaces of vector-valued function spaces. *Math. Proc. Camb. Phil. Soc.*, 101:107{112, (1987).
2. F. Bombal. On (V) sets and Pelczynski's property (V). *Glasgow Math. J.*, 32:109{120, (1990).
3. J. Bourgain.  $H^1$  is a Grothendieck space. *Studia Math.*, 75:193{216, (1983).
4. J. Bourgain. On weak compactness of the dual of spaces of analytic and smooth functions. *Bull. Soc. Math. Belg.*, B 35:111{118, (1983).
5. P. Cembranos, N. J. Kalton, E. Saab, and P. Saab. Pelczynski's Property (V) on  $C(\mathbb{R})$  spaces. *Math. Ann.*, 271:91{97, (1985).
6. D. L. Cohn. *Measure Theory*. Birkhauser, Basel, Stuttgart, (1980).
7. F. Dearing. Weakly compact operators on the disk algebra. *J. of Algebra*, 45:284{294, (1977).
8. J. Diestel. *Sequences and Series in Banach Spaces*, volume 92 of Graduate Text in Mathematics. Springer Verlag, New York, first edition, (1984).
9. J. Diestel and Jr. J. J. Uhl. *Vector Measures*, volume 15 of Math Surveys. AMS, Providence, RI, (1977).
10. N. Dinculeanu. *Vector Measures*. Pergamon Press, New York, (1967).
11. G. Godefroy and P. Saab. Weakly unconditionally convergent series in  $M$ -ideals. *Math. Scand.*, 64:307{318, (1990).
12. S. V. Kisliakov. Uncomplemented uniform algebras. *Mat. Zametki*, 18:91{96, (1975).
13. J. Munkres. *Topology, a first course*. Prentice Hall, Englewood Cliffs, New Jersey, (1975).
14. A. Pelczynski. On Banach spaces on which every unconditionally converging operator is weakly compact. *Bull. Acad. Polon. Sci.*, 10:641{648, (1962).
15. N. Randrianantoanina. Complemented copies of  $\ell_1$  and Pelczynski's property (V) in Bochner spaces. Submitted for publication.
16. W. Ruess. Duality and geometry of compact operators. *Math. Studies* 90, North Holland, (1984).
17. E. Saab and P. Saab. Stability problems in Banach spaces. *Lecture Notes in Pure and Appl. Math.*, 136:367{394, (1992).
18. Z. Semadeni. *Banach spaces of continuous functions*. PWN, Warszawa, (1971).
19. M. Talagrand. Weak Cauchy sequences in  $L^1(E)$ . *Amer. J. Math.*, 106:703{724, (1984).
20. A. Tulecea and C. Ionescu. *Topics in the theory of lifting*, volume 48 of *Ergebnisse der Mathematik und Ihrer Grenzgebiete*. Springer-Verlag, Berlin and New York, first edition, (1969).
21. A. Unger. Weak compactness in  $L^1(\mathbb{R}; X)$ . *Proc. Amer. Math. Soc.*, 113:143{149, (1991).

Department of Mathematics, The University of Texas at Austin, Austin,  
TX 78712-1082

E-mail address: nrandri@math.utexas.edu