

Eigenfunctions on the Finite Poincare Plane
by Jinghua Kuang

1. Introduction. Let F_q be a finite field with q elements (q odd.) Fix a non-square element $\delta \in F_q$.

$$H_q = F_q \left(\frac{p}{-} \right) \quad F_q = \{fx + y \frac{p}{-} jx; y \in 0, 2, F_q\}$$

is called the finite Poincare plane. From [1], [2], [4], [7], we have the following facts:

1. $G = GL_2(F_q)$ acts on H_q by linear transformation:

$$gz = \frac{az + b}{cz + d}; \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G; \quad g \neq \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in H_q;$$

Let $K = \begin{pmatrix} a & b \\ b & a \end{pmatrix}$. One may identify $G=K$ with H_q and with $P = \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \quad y \in 0; x \in F_q$.

2. There is a G -invariant pseudo-distance on H_q :

$$(z_1; z_2) = \frac{N(z_1 - z_2)}{\operatorname{Im} z_1 \operatorname{Im} z_2} \quad 8z_1, z_2 \in H_q;$$

where N is the norm from $F_q \left(\frac{p}{-} \right)$ to F_q and $\operatorname{Im} z = y$ if $z = x + y \frac{p}{-}$.

3. For a $z \in F_q$ and $a \in 4$, a graph structure $X_q(\cdot; a)$ is defined on H_q : $(z_1; z_2)$ is an edge if $(z_1; z_2) = a$. Katz [3] and Li [5] proved that $X_q(\cdot; a)$ are Ramanujan graphs using different methods.

However, distributions of eigenvalues of the adjacency matrices have not been well understood. We will calculate the first and second moments of the asymptotic distribution of the eigenvalues of the adjacency matrices and provide evidence for Terras' conjecture in [7].

Let V be the space of all complex functions on H_q . Using character sums, Evans [2] constructed an orthogonal basis of V that diagonalizes all the adjacency matrices. Kuang [4], using group representation theory, also constructed such an orthogonal basis. This paper will also compare these two bases.

2. Hecke Algebra. Since K is the analogue of $O(2)$ in the classical Poincare upper half plane, it is natural to consider the Hecke algebra $H(G; K)$, which is defined as the algebra of all bi- K -invariant complex functions on G under convolution. Since K is the isotropic subgroup of $\frac{p}{-}$, $(g; 1) = (g^{-1}; 1)$ implies that $KgK = Kg^{-1}K$. Therefore, $H(G; K)$ is commutative. Obviously, $H(G; K)$ has dimension q . Hence there exist q idempotents in $H(G; K)$.

3. Idempotents. For an irreducible representation ρ of G , let $\rho^0(g) = \frac{1}{|K|} \sum_{k \in K} \operatorname{tr}(\rho(kg))$. ρ is spherical (i.e. $\operatorname{Res}_K^G(\rho) = \rho_K$) if $\rho^0(g) \neq 0$. From [4], we have

$$\rho_1^0 \rho_2^0 = \begin{cases} 0; & \text{if } \rho_1 \neq \rho_2; \\ \frac{\dim(\rho)}{\dim(G)} \rho; & \text{if } \rho_1 = \rho_2 = \rho. \end{cases}$$

Let χ be a multiplicative character of $F_q \left(\frac{p}{-} \right)$ of order $q-1$, ψ be a multiplicative character of order $q-1$, and $s = \chi(q-1)=2$. There are q irreducible spherical representations of G (see [6]):

$$= 1_G;$$

$$\text{or } = (s; s) = q \text{ dimensional irreducible component of } \operatorname{Ind}_B^G(s; s);$$

$$\text{or } = (\psi; \psi) = \operatorname{Ind}_B^G(\psi; \psi); \quad \psi = 1; \dots; \frac{q-3}{2};$$

$$\text{or } = \psi_j = \text{cuspidal representation corresponding to } \psi_j = \chi^j(q-1); \quad \psi_j = 1; \dots; \frac{q-1}{2};$$

(B is the Borel subgroup of G .) Let $\chi_j = \frac{\dim(\chi_j)}{\dim(G)}$. Let $\rho_0 = 1_G$, $\rho_j = (\psi; \psi); \psi_j = 1; \dots; (q-3)=2$, $\frac{q-1}{2} = (s; s)$ and $\frac{q-1}{2} + j = \psi_j; \psi_j = 1; \dots; (q-1)=2$: Then

Research at MSRI supported in part by NSF grant # DMS 9022140

Theorem 1 (Kuang [4]). $e_0, e_1, \dots, e_{q-1}$ are the q idempotents in $H(G; K)$.

4. Representation of $H(G; K)$. $H(G; K)$ has a natural representation on $L^2(H_q) = L^2(P)$, the space of all complex functions on $H_q = P$:

$$T_{\mathbf{h}}(f)(p) = \frac{1}{|K|} \sum_j f(j^{-1}p), \quad T_{\mathbf{h}}^{-1}(p) = \frac{1}{|K|} \sum_{h \in G} f(h) \overline{h^{-1}p} = \sum_{p_1 \in P} f(p_1) \overline{p_1^{-1}p};$$

for $\mathbf{h} \in H(G; K)$, $f \in L^2(P)$ and $p \in P$.

For the usual inner product $\langle f, g \rangle$ on $L^2(P)$, we have $\langle T_{\mathbf{h}}(f), T_{\mathbf{h}'}(g) \rangle = \langle f, g \rangle$ where $\mathbf{h}' = \overline{\mathbf{h}^{-1}}$. That is, $H(G; K)$ acts on $L^2(P)$ self-adjointly. Hence, there exists a basis of simultaneous eigenfunctions of $H(G; K)$.

5. Construction of the Bases. Let χ be a fixed non-trivial additive character of F_q . Denote $\chi_a(x) = \chi(ax)$. Define

$$e_i(g) = \chi_i(y) \text{ if } gK = \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} K, i = 1, \dots, q-1.$$

$$\chi_a(g) = \chi(y) \chi_a(x) \text{ if } gK = \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} K, a \in F_q,$$

where $\chi(y) = 1$ if $y = 1$; $= 0$ otherwise. Then

Theorem 2 (Kuang [4]). $e_i, i = 1, \dots, q-1$ and $\chi_a, i = 1, \dots, q-1; a \in F_q$ are simultaneous eigenfunctions of $H(G; K)$ and make up an orthogonal basis of $L^2(P)$. Moreover, $T_j(e_i) = e_i$ if $j = (i, i)$; $= 0$ otherwise. $T_j(\chi_a) = \chi_a$ if $j = (j, 0)$; $= 0$ otherwise.

Terras [8] constructed the following functions:

$$K_{i, a} \left(\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \right) = \chi_a(x) \chi_i(y) \sum_{u \in F_q} \chi_a(u) \chi_i(u^2 - y^2);$$

Evans [2], based on Velasquez's work according to [9], constructed the following functions:

$$H_{t, i, a} \left(\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \right) = \frac{\chi_a(x)}{q+1} \sum_{u \in F_q} \chi_a(u) \sum_{s \in U_1} \chi_i(s) + \frac{1}{t} + \frac{y}{t} + \frac{t}{y} - \frac{u^2}{ty};$$

where $U_1 = \{f \in F_q \mid fN(\cdot) = 1\}$ and the asterisk indicates that when $y = t$, the $u = 0$ term is multiplied by $q+1$.

Theorem 3 (Terras, Velasquez, Evans [2]). $K_{i, a}, i = 1, \dots, q-1, a \in F_q$ and $H_{t, i, a}, t \in F_q, i = 1, \dots, q-1, a \in F_q$ are simultaneous eigenfunctions of $H(G; K)$. There exist a_0 for any fixed i and a such that $K_{i, 0}, i = 1, \dots, q-1, K_{i, a}, i = 1, \dots, (q-1)=2, a \in F_q$ and $H_{t_0, i, a}, i = 1, \dots, (q-1)=2, a \in F_q$ make up an orthogonal basis of $L^2(P)$.

6. Comparison of the Bases. We now prove that the two bases constructed above are essentially same. Let us denote by e_i the spherical representation that give rise to e_i .

Theorem 4. The following identities hold.

$$\begin{aligned} \overline{C_i(0)} e_i &= K_{i, 1}, i = 1, \dots, q-1; \\ e_a e_i &= \frac{\dim(e_i)}{|G|} C_i(a) K_{i, a}, i = 1, \dots, (q-1)=2; \\ e_{a+i+(q-1)=2} &= \frac{\dim(e_{i+(q-1)=2})}{|G|} H_{t, i, a}, i = 1, \dots, (q-1)=2; \end{aligned}$$

where $C_i(a) = \sum_{x \in F_q} a(x)^{-i} (x^2)^i$.

Proof. $K_{i,1} = \overline{C_i(0)}$ is obvious. Since for any pair of $a, b \in F_q$, $K_{i,b}$ and $\overline{a - i}$ are eigenfunctions of $H(G; K)$ of the same eigenvalues ($i = 1; \dots; (q-1)/2$), $\overline{a - i}$ is a complex linear combination of $K_{i,b}$ ($b \in F_q$). Now,

$$\begin{aligned} hK_{i,b}; \overline{a - i} &= \sum_{x \in F_q} K_{i,b}(x) \overline{a - i}(x) \\ &= \sum_{x \in F_q} \sum_{u \in F_q} b(u - x)^{-i} (y)^{-i} (u^2 - y^2)^{-i} \sum_{v \in F_q} (a + b)v \\ &= 0 \quad \text{if } a + b \neq 0 \end{aligned}$$

Hence $\overline{a - i}$ is a multiple of $K_{i,a}$. Using the equation (2.17) in [2], we calculate

$$\begin{aligned} \overline{a - i}(1) &= \sum_{u \in F_q} a(u)^{-i} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \\ &= \frac{\dim(\overline{a - i})}{\#j(q+1)} C_i(a) \overline{C_i(a)}: \end{aligned}$$

And $K_{i,a}(1) = \overline{C_i(a)}$. Therefore, $\overline{a - i} = \frac{\dim(\overline{a - i})}{\#j(q+1)} C_i(a) K_{i,a}$.

For the third identity, we use the equation (2.16) in [2]. Let $\#(z)$ be the cardinality of the K -orbit of z in H_q . Let $\hat{i} = i + (q-1)/2$. Then,

$$\begin{aligned} \overline{a - \hat{i}}(p) &= \sum_{u \in F_q} a(u)^{-\hat{i}} \begin{pmatrix} y & x & u \\ 0 & 1 & \end{pmatrix} \\ &= \sum_{u \in F_q} a(x - u)^{-\hat{i}} \begin{pmatrix} y & u \\ 0 & 1 \end{pmatrix} \\ &= \sum_{u \in F_q} a(x) \sum_{u \in F_q} a(u) \frac{\dim(\overline{a - \hat{i}})}{\#j} \frac{1}{\#(u + y)} \sum_{j \in 2U_1} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}^{\hat{i}} (s) + \frac{1}{2} + \frac{(u + y)^{p-p-1}}{j} \\ &= \sum_{u \in F_q} a(x) \sum_{u \in F_q} a(u) \frac{\dim(\overline{a - \hat{i}})}{\#j} \frac{1}{\#(u + y)} \sum_{j \in 2U_1} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}^{\hat{i}} (s) + \frac{1}{2} + \frac{u^2}{y} - \frac{1}{y} \\ &= \frac{\dim(\overline{a - \hat{i}})}{\#j} H_{i, \overline{a - i}}(p) \end{aligned}$$

7. Comments and Questions. For any function f on G , define the Fourier coefficient

$$F(g; a; f) = \sum_{x \in F_q} f \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g(a)(x):$$

From [4], we see $h_{a - i}; \overline{a - i} = \#j(q-1) F(1; a; \overline{a - i})$. So $F(1; a; \overline{a - i}) \neq 0$. But can one evaluate $F(1; a; \overline{a - i})$? We have $F(1; a; \overline{a - i}) = \overline{a - i}(1)$. Hence, for $i = 1; \dots; (q-1)/2$,

$$\begin{aligned} F(1; a; \overline{a - i}) &= \frac{\dim(\overline{a - i})}{\#j(q+1)} \overline{C_i(a)}^2 = \frac{\dim(\overline{a - i})}{\#j(q+1)} \overline{C_i(a)}^2 \\ F(1; a; \overline{a - \hat{i}}) &= \frac{\dim(\overline{a - \hat{i}})}{\#j} \sum_{u \in F_q} a(u) \frac{1}{\#(u + y)} \sum_{j \in 2U_1} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}^{\hat{i}} (s) + \frac{1}{2} + \frac{u^2}{y} - 2 \end{aligned}$$

However, we don't know much about $C_i(a)$ and about the last set of sums. We note that $C_i(0)$ is essentially a Jacobisum, in fact, $C_i(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} J(\hat{i}; s)$.

8. Connection to $X_q(\cdot; a)$. There are q double cosets: $D_a = KgK$, where $(g; 1) = a \in \mathbb{F}_q$. Let χ_a be the characteristic function of the set D_a . Then $A_a = \frac{1}{K} \sum_j T_{a,j}$ acting on $L^2(H_q)$ is the adjacency matrix of $X_q(\cdot; a)$. Therefore each basis constructed in Section 5 diagonalizes all the adjacency matrices. Let $S_a = D_a \setminus P$. Then

$$\lambda_i(a) = \frac{\sum_j \chi_a(j)}{\dim(S_a)} \chi_a(j_i) \chi_a(j_i); i = 0, 1, \dots, q-1$$

are all the eigenvalues of A_a . Katz [3], Li [5] proved that $|\lambda_i(a)| \leq 2\sqrt{q}$ for $i = 1, \dots, q-1$, which confirms that $X_q(\cdot; a)$ is Ramanujan.

Fix $a \in \mathbb{F}_q$, Terras [7] conjectured that $\sum_{j=1}^{q-1} \lambda_j(a) = \sum_{j=1}^{q-1} \sqrt{q} \chi_j(a)$ asymptotically has Sato-Tate distribution, i.e. for $E \subset [0, 2]$,

$$\lim_{q \rightarrow \infty} \frac{1}{q-1} \sum_{j=1}^{q-1} \lambda_j(a) \chi_j(a) = \int_E \sqrt{q} \chi_j(a) d\mu = \frac{1}{2} \int_E \sqrt{4-x^2} dx$$

9. Moments. Define a $q \times q$ matrix

$$M = \left(\frac{\sum_j \chi_a(j)}{\dim(S_a)} \chi_a(j_i) \chi_a(j_i) \right)_{i=0,1,\dots,q-1}$$

The idempotent property of χ_i ($i = 0, \dots, q-1$) implies that $MM^0 = \frac{1}{K} \sum_j \chi_j \chi_j = I_q$. (M^0 is the transpose of M .) Hence, $M^0 M = \frac{1}{K} \sum_j \chi_j \chi_j = I_q$, that gives

$$\begin{aligned} \frac{1}{q-1} \sum_{i=0}^{q-1} \lambda_i(a) \chi_i(a) &= 0 \\ \frac{1}{q-1} \sum_{i=0}^{q-1} \lambda_i(a)^2 &= \frac{\sum_j \chi_j \chi_j}{K \sum_j (q-1)} \end{aligned}$$

That, in turn, implies

$$\begin{aligned} \lim_{q \rightarrow \infty} \frac{1}{q-1} \sum_{i=0}^{q-1} \lambda_i(a) \chi_i(a) &= 0 \\ \lim_{q \rightarrow \infty} \frac{1}{q-1} \sum_{i=0}^{q-1} \lambda_i(a)^2 &= 1 \end{aligned}$$

Therefore, the first and second moments of $\sum_{j=1}^{q-1} \lambda_j(a) \chi_j(a)$ asymptotically match with those of the Sato-Tate distribution.

Acknowledgement. This is a written-up of a talk that the author delivered at the Workshop on Spectral Theory of Automorphic Forms and Number Theory, in MSRI, Berkeley, California, October 24-28, 1994. Some calculation was completed while the author was visiting MSRI. He thanks the MSRI for its hospitality.

References.

- [1]. J. Angel, N. Celniker, S. Poulos, A. Terras, C. Trimble, E. Velasquez Special Functions on Finite Upper Half Planes, in AMS Contemporary Math. Vol. 138, 1992.
- [2]. R. Evans Character sums as orthogonal eigenfunctions of adjacency operators for Cayley graphs, in Proc. of a conference on finite fields, AMS contemporary Math. Vol. 168, 1994.
- [3]. N. Katz Estimates for Sato-Andrade Sums, J. für die Reine und Angew. Math. 438 (1993), 143-161.
- [4]. J. Kuan A Natural Orthogonal Basis of Eigenfunctions of the Hecke Algebra Acting on Cayley Graphs, to appear in the Proceedings of American Mathematical Society.
- [5]. W. Li A survey of Ramanujan graphs, Proc. of a Conference on Arithmetic Geometry and Coding Theory at Luminy, 1993 (to appear).

- [6]. I. Piatetski-Shapiro Complex Representation of $GL_2(\mathbb{F}_q)$, *AMS Contemporary Math.* Vol. 16, 1983.
- [7]. A. Terras Are Finite Upper Half Plane Graphs Ramanujan? in *DIMACS Series in Discrete Mathematics and Theoretic Computer Science*, Vol. 10 (1993), 125-142
- [8]. A. Terras Fourier Analysis on Finite Groups and Applications, *U.C.S.D. Lecture Notes*, 1991-2.
- [9]. E. Velasquez, Private Conversation, October 27, 1994.

Department of Mathematics
 Pennsylvania State University, at Fayette
 Uniontown, PA 15401
 E-mail address: kuang@math.psu.edu