Environm ent-induced dynam ical chaos

Bidhan Chandra Bag and Deb Shankar Ray Indian Association for the Cultivation of Science, Jadavpur, Calcutta 700 032, INDIA. (M arch 30, 2022)

We exam ine the interplay of nonlinearity of a dynam ical system and therm al uctuation of its environment in the \physical lim it" of sm all damping and slow di usion in a sem iclassical context and show that the trajectories of c-num ber variables exhibit dynam ical chaos due to the therm al uctuations of the bath.

PACS number(s): 05.45 M t

The interplay of nonlinearity of a dynam ical system and therm al uctuations of its environment has been one of the major areas of investigation in the recent past. These studies have enriched our understanding of nonequilibrium processes in several contexts, such as, symm etry between the growth and decay of classical uctuations in equilibrium [1], interesting topological features of patterns of paths of large uctuations in nonlinear system s [1], existence of generalized nonequilibrium potential [2], in uence of nonlinearity on dissipation in multiphoton processes [3] and higher order di usion in a nonlinear system [4], etc. W hile the developm ent in these areas is largely con ned to classical dom ain we exam ine a related issue in the sem iclassical context. Since quantization is likely to add a new dimension to the interplay of nonlinearity and stochasticity in a weakly dissipative system, it is worthwhile to consider the physical lim it of sm all dam ping and slow di usion due to therm al uctuations of the environm ent and look for the thermal uctuations-induced features of nonlinearity in the dynam ics. In this communication we speci cally explore som e interesting aspects of dynam ical chaos in a driven bistable system whose origin lies at the uctuations of the environm ental degrees of freedom .

To describe the dissipative quantum dynamics of a system we rst consider the traditional system -reservoir model developed over the last few decades [3,5,6]. The H am iltonian of the bare system is coupled to an environ-ment modeled by a reservoir of harm onic oscillatorm odes characterized by a frequency set f_jg . The quantum dynamics is generated by the overall H am iltonian operator H for the system, environment and their coupling as follows;

$$H = H_{0} + h_{j=1} x^{i} b_{j}^{y} b_{j} + h_{j=1} g(y_{j}) b_{j} + g^{2}(y_{j}) b_{j}^{y} x;$$
(1)

where

$$H_0 = \frac{p^2}{2} + V(x)$$
; (2)

de nes the usual kinetic and potential energy term s corresponding to the system , x and p being the position and m om entum operators, respectively. The second and the third terms in (1) specify the reservoir modes and their linear coupling to the system . g() denotes the system - reservoir coupling constant.

System atic elimination of the reservoir modes in the usualway, using Born and Markov approximations leads one to the following standard reduced density matrix equation of the system only [5].

$$\frac{d}{dt} = \frac{i}{h} [H_0;] + \frac{1}{2} (1 + n) (2a a^{y} a^{y} a a^{y} a) + \frac{1}{2} n (2a^{y} a aa^{y} aa^{y}) ; \qquad (3)$$

Here the system operator co-ordinate x is related to the creation and the annihilation operators a^{y} ; a respectively as $x = (\frac{p-1}{2!})(a + a^{y})$. ! is the linearised frequency of the system. Also the spectral density function of the reservoir is replaced by a continuous density D (!). > 0 is the limit of 2 $\dot{g}(!)^{2} \frac{D(!)}{!}$ as ! ! 0_{+} and is assumed to be nite. is the relaxation or dissipation rate and n is the di usion coe cient D. n (= [exp $\frac{h!}{kT}$ 1] ¹) is the average therm all photon number of the reservoir. The term s analogous to Stark and Lam b shifts have been neglected.

The rst term in Eq.(3) corresponding to the dynamical motion of the system refers to Liouville ow. The term s containing arise due to the interaction of the system with the surroundings. W hile $\frac{n}{2}$ term s denote the di usion of uctuations of the reservoir modes into the system mode, $\frac{1}{2}$ term s refer to the loss of energy from the system into reservoir. In the lim it T ! 0; i.e.; n ! 0 the system is in uenced by pure quantum noise or vacuum uctuations [N ote that 1 in (n + 1) in Eq.(3) corresponds to the vacuum].

O ur next task is to go over from a full quantum operator problem described by the Eq.(3) to an equivalent c-num ber problem. To this end we consider the quasiclassical distribution function W (x;p,t) of W igner. x;p are now c-num ber variables. Rewriting Eq.(3) in a quasiclassical language we obtain

$$\frac{@W}{@t} = p\frac{@W}{@x} + \frac{@V}{@x}\frac{@W}{@p} + \frac{2}{2} \frac{@xW}{@x} + \frac{@pW}{@p}$$

$$+ \frac{h}{2!} (n+1) \frac{\theta^2 W}{\theta x^2} + \frac{h!}{2} (n+1) \frac{\theta^2 W}{\theta p^2} + \frac{X}{2^{2n}} \frac{h^{2n} (1)^n}{(2n+1)!} \frac{\theta^{2n+1} V}{\theta x^{2n+1}} \frac{\theta^{2n+1} W}{\theta p^{2n+1}} :$$
(4)

in Eq.7 is a parameter used to identify the environmentinduced e ect on the dynamics described by Eq.(4) (kept for bookkeeping in the calculation and put = 1 at the end).

In the sem iclassical lim it h! << kT, we have n + 1 - nand D kT so that Eq.(7) reduces to

$$\frac{@W}{@t} = p\frac{@W}{@x} + \frac{@V}{@x}\frac{@W}{@p} + \frac{1}{2} \frac{@xW}{@x} + \frac{@pW}{@p} + p \frac{1}{2}\frac{@^2W}{@x^2} + \frac{1}{2}\frac{@^2W}{@p^2} + \frac{X}{12}\frac{h^{2n}(1)^n}{2^{2n}(2n+1)!}\frac{@^{2n+1}V}{@x^{2n+1}}\frac{@^{2n+1}W}{@p^{2n+1}} :$$
(5)

The overall dynam ics described above is a superposition of two contributions, i. e. , the Liouville-W igner dynam ics and the system reservoir dissipative dynam – ics. That the two contributions act independently is an assumption. The master equation (3) [or its W igner function version (4)] is the most popular one in quantum optics. It has been extensively used [7], for the strongly nonlinear processes like three-wave, four-wave mixing and strong coherent light-matter interaction phenom ena. The equation has also been applied in the context of chaos, e. g., in the dissipative standard map [8], dissipative logistic map [9], sem iclassical theory of quantum noise in open chaotic systems [10,11] and in the studies of decoherence in relation to chaos for analysis of quantum -classical correspondence [12,13].

In the sem iclassical (h ! 0) lim it the dissipative quantum dynam ics can be conveniently described by $\ KB -$ like" ansatz (we refer to $\ W \ KB -$ like" since we are considering m ore that one dimension. Traditional W KB refer to one dimension only) [1,14] of Eq.(5) for W igner function of the form

W
$$(x;p;t) = Z(t) \exp(\frac{s}{h})$$
: (6)

Here Z (t) is a prefactor and s(x;p;t) is a classical action which is a function of c-number variables x and p, satisfying the following H am ilton-Jacobi equation

$$\frac{\vartheta s}{\vartheta t} + p \frac{\vartheta s}{\vartheta x} \quad \frac{\vartheta s}{\vartheta x} \frac{\vartheta s}{\vartheta p} \quad \frac{\omega s}{2} \left(x \frac{\vartheta s}{\vartheta x} + p \frac{\vartheta s}{\vartheta p} \right) + \left(\frac{\vartheta s}{\vartheta x} \right)^{2}$$

$$+ \frac{1^{2} (\vartheta s)^{2}}{(\vartheta p)^{2}} + \frac{X}{n} \frac{x^{2n} (1)^{3n+1}}{2^{2n} (2n)!} \frac{\vartheta^{2n+1} V}{\vartheta x^{2n+1}} \frac{\vartheta s}{\vartheta p} = 0 : (7)$$

The derivation of Eq.(7) is based on the consideration of the \physical lim it" of weak dissipation and slow uctuations in the sense $\frac{D_1}{h^2} = \frac{1}{h}$ where $D_1 = \frac{D}{2!^2}$ (note that

 $\rm D_{1}$ and h have same dimension). The above equation can be solved by integrating the H am iltonian equations of motion,

$$\underline{\mathbf{x}} = \mathbf{p} \quad \frac{1}{2} \quad \mathbf{x} + 2 \quad \mathbf{P}$$

$$\underline{\mathbf{x}} = \mathbf{P} \quad \frac{1}{2} \quad \mathbf{X}$$

$$\underline{\mathbf{p}} = \mathbf{V}^{0} + \frac{1}{2} \quad \mathbf{p} \quad 2!^{2} \quad \mathbf{X} \quad \frac{\mathbf{X}}{\frac{(1)^{3n+1}}{2^{2n}(2n)!}} \frac{(2^{2n+1}\mathbf{V})^{2n}}{(2^{2n+1})!} \mathbf{X}^{2n}$$

$$\underline{\mathbf{P}} = \mathbf{V}^{00}\mathbf{X} + \frac{1}{2} \quad \mathbf{P} \quad \frac{\mathbf{X}}{\frac{(1)^{3n+1}}{2^{2n}(2n+1)!}} \frac{(2^{2n+2}\mathbf{V})^{2n+1}}{(2^{2n+2}\mathbf{V})!} \mathbf{X}^{2n+1} \quad (8)$$

which are derived from the following H am iltonian H $_{\rm eff}$

$$H_{eff} = pP \quad V^{0}X \quad \frac{1}{2} (xP + pX) + (P^{2} + !^{2}X^{2}) \\ + \frac{X}{n} \frac{(1)^{3n+1}}{2^{2n} (2n+1)!} \frac{e^{2n+1}V}{e^{x^{2n+1}}} X^{2n+1} :$$
(9)

Here we have put $\frac{@s}{@x} = P$ and $\frac{@s}{@p} = X$. The introduction of additional degree-of-freedom by incorporating the auxiliary m om entum (P) the and co-ordinate (X) m akes the system e ectively a two-degree-of-freedom system. The origin of these two variables is the therm al uctuations of the reservoir. It is easy to identify the environm entin Eqs.(7-9). The auxiliary related terms containing Ham iltonian (9) is therefore not to be confused with the m icroscopic H am iltonian (1) which describes a system in contact with a reservoir with in nite degrees of freedom . A Ithough the phase space trajectories concern uctuations of c-num ber variables in the form al sense, because of the equations of motion (8) described by a Ham iltonian (9), the motion is strictly deterministic. The experiments on the corresponding classical version of the problem by Luchinsky and M cC lintock [1] have dem onstrated that a trajectory of uctuation is indeed a part of physical reality. W e em phasize, how ever, here a num ber of distinguishing features in this context. W hile the studies by Luchinsky and M cC lintock [1] and G raham and Tel [2] concern overdam ped lim it, we consider here a weakly dissipative system . Furtherm ore because of the quantum correction, the phase space trajectories of uctuations are signi cantly modied by sem iclassical features. The introduction of these quantum features at a sem iclassical level through a c-num ber H am iltonian description of a dissipative evolution in the physical lim it of weak dam ping and slow di usion due to therm alnoise is the essential content of the present work. Since the Eq.(9) describes determ inistic evolution under nonlinear potential, the pattern of trajectories of uctuations may display chaotic behaviour. In what follows we investigate this dynamical aspect of the dissipative system .

The equations derived in the weak therm alnoise limit for the weakly dissipative sem iclassical systems are fairly general. For illustration we now consider a simple model system H am iltonian H $_0$ (see Eq2)

$$H_{0} = \frac{p^{2}}{2} + ax^{4} bx^{2} + gx\cos_{0}t$$
 (10)

which describes a bistable potential driven by a timeperiodic eld. a and b are the constants of the potential V (x). The fourth term in (10) includes the e ect of coupling of the system as well as the strength of the eld of frequency $_0$. For the H am iltonian (10) the Eqs.(8) read as

$$\underline{x} = p - \frac{1}{2} x + 2 P$$

$$X_{-} = P - \frac{1}{2} X$$

$$\underline{p} = 4ax^{3} 2bx + g\cos t + (\frac{1}{2}p - 2!^{2}X) 3axX^{2}$$

$$P_{-} = (12ax^{2} - 2b)X + \frac{1}{2} P - aX^{3}$$
(11)

which are derivable from the auxiliary Ham iltonian

$$H_{eff} = Pp \quad (4ax^{3} \quad 2bx + g\cos!t)X \quad \frac{1}{2} \quad (xP + pX) + (P^{2} + !^{2}X^{2}) + axX^{3}: \quad (12)$$

The system H am iltonian (10) has served as a standard paradigm for a number of theoretical and experimental investigations [1,10,15] over m any years. For the present purpose we choose the following parameter values; $a = \frac{1}{4}$, $b = \frac{1}{2}$, $_0 = 6.07$ and g = 10. Since we are considering the physical limit of weak damping and small di usion we take the initial conditions for the auxiliary variables X and P (which originate from the uctuations due to environment) as P = 0 and X ! 0 (we have used X = 1.5 10⁶). This ensures a vanishing H am iltonian H eff for the entire num erical investigation that follows below. We rst consider a specie c trajectory with the initial

condition p = 0 and x = 2.512 for sm all values of (typical 0.01). Under this condition (= 1) the system is vanishingly coupled to the surroundings and consequently the dynam ical behaviour is e ectively due to the weak dissipation only. We illustrate this situation in F ig. 1 in terms of a Poincare m ap for the phase space which exhibits strong global chaos. On the other hand when the parameter is switched o (= 0) the system displays typical weak chaos (F ig.2). The similar behaviour has been observed for other sets of initial conditions for x and p (W e have not reproduced them here for the sake of brevity).

The elect of weak dissipation and slow dilusion due to thermal uctuations from the surroundings can be seen in the case of other sets of initial conditions also. For the initial condition p = 0 and x = 2:49 one observes for = 0.01 dissipative strong chaos(= 1). This is illustrated in Fig.3. It is interesting to note that for = 0 the

sam e trajectory gets localized in the left well as a regular

one as shown in Fig.4. The weakly chaotic and regular trajectories in Figs. 2 and 4, respectively, are purely sem iclassical in nature (in the absence of any coupling to the surroundings). The strong global chaotic behaviour as shown in Figs. 1 and 3 has therefore its origin in the therm all uctuations of the reservoir. In other words the chaotic behaviour or its enhancement is exclusively due to therm all uctuations from the surroundings, which becomes appreciable even in the physical limit of weak damping and slow dilusion of the therm all noise within the sem iclassical description. We have checked this assertion for some other values of the initial conditions for the system oscillator.

The reduction of the system -reservoir H am iltonian description [H in Eq.1] for a dissipative quantum system to an auxiliary H am iltonian description [H $_{eff}$ in Eq.9] e ectively reduces the in nite-degree-of-freedom system to a two-degree-of-freedom system where the auxiliary degree of freedom characterized by X and P owe their origin in the uctuations of the reservoir. Since X and P appear as the multiplicative factors in the auxiliary H am iltonian H $_{eff}$, the weak therm alnoise lim it makes H $_{eff}$ a vanishing H am iltonian. The observed sem iclassical chaos m ay therefore be regarded as a dynam icalm anifestation of the interplay of nonlinearity and therm all uctuations.

In this paper we have exam ined the weak therm alnoise lim it of a sem iclassical dissipative nonlinear system . We have shown that the vanishing H am iltonian method can be suitably extended to follow the phase space trajectories of uctuations of c-num ber variables which exhibit dynam ical chaos. In view of the accessibility of the model to analogue electronic circuits [1] we believe that the results discussed bear further experim ental relevance in the sem iclassical context.

B.C.Bag is indebted to the CSIR for partial nancial support and to J.Ray Chaudhuri for helpful discussions.

- [1] D.G. Luchinsky and P.V.E.M cC lintock, Nature 389 403 (1997).
- [2] R.Graham and T.Tel, Phys. Rev. Letts 52 9 (1984).
- [3] G.G angopadhyay and D.S.Ray, J.Chem. Phys. 46 4693 (1992).
- [4] M .Battezzati, J.Chem .Phys.111, 9932 (1999).
- [5] W. H. Louisell, Quantum Statistical Properties of Radiation (W iley, New York, 1973); A. J. Leggett and A.O. Caldeira, Physica A 121 587 (1983); G.W. Ford, J.T. Lew is and R. J. O'Connell, Ann. Phys. (N.Y.) 185 270 (1988).
- [6] See, for example, the review by G.G angopadhay and D. S.Ray, in Advances in Multiphoton Processes and Spectroscopy Vol8, edited by S.H.Lin, A.A. Villayes and F. Fujimura, (W orld Scientic, Singapore, 1993).

- [7] P. M eystre and M. Sargent III, E lements of Quantum Optics (Springer-Verlag, Berlin, 1989); M. D. Reid and D. F. W alls, Phys. Rev. A 33 4465 (1986); 34 4929 (1986).
- [8] T.D ittrich and R.Graham, Ann.Phys.200 363 (1990);
 Phys.Scr.40 409 (1989).
- [9] M.E.Goggin, B.Sundaram and P.W. Milonni, Phys. Rev.A 41, 5705 (1990).
- [10] B.C.Bag, S.Chaudhuri, J.Ray Chaudhuri and D.S. Ray, Physica D 125 47 (1999).
- [11] B.C.Bag and D.S.Ray, J.Stat. Phys. 96 271 (1999).
- [12] W . H. Zurek and J. P. Paz, Phys. Rev. Lett. 72 2508 (1994); Physica D 83, 300 (1995).
- [13] A.K.Pattanayak, Phys.Rev.Lett.83, 4526 (1999).
- [14] C.M. Bender and S.A.Orszag, Advanced M athem atical M ethods for Scientists and Engineers (M cG raw H ill, New York, 1978).
- [15] W .A.Lin and L.E.Ballentine, Phys. Rev. Lett. 65 2927 (1990).

FIG.1. Plot of x vs p on the Poincare surface of section (X = 0) for Eq.(11) with initial condition x = 2.512, p = 0, X ! 0, P = 0, = 0:01 = 1. (Units are arbitrary).

FIG.2. Same as in Fig.1 but for = 0.

FIG.3. Same as in Fig.1 but for x = 2:49.

FIG.4. Same as in Fig.3 but for = 0.



Fig. 1

B. C. Bag and D. S. Ray



Fig. 2

B. C. Bag and D. S. Ray



Fig. 3



Fig. 4