Universala-matrix as integral operator.

S D erkachov^{a,c1}, D . K arakhanyan^{b2} & R . K irschner^{c3}

- ^a Department of Mathematics, St Petersburg Technology Institute, Sankt Petersburg, Russia
- b Yerevan Physics Institute,
 BrAlikhanian st2, 375036, Yerevan, Armenia.
- ^c Naturw issenschafflich-Theoretisches Zentrum und Institut für Theoretische Physik, Universitat Leipzig, Augustusplatz 10, D-04109 Leipzig, Germany

Abstract.

We derive the integral operator form for the general rational solution of the Yang-Baxter equation with s'(2jl) symmetry. Considering the dening relations for the kernel of the Roperator as a system of second order dierential equations we observe remarkable reduction to a system of simple rst order equations. The obtained kernel of Roperator has a very simple structure. To illustrate all this in the simplest situation we treat also the s'(2) case.

¹e-m ail: Sergey D erkachov@ itp uni-leipzig de

²e-m ail: karakhan@ lx2.yerphi.am

³e-mail:Roland:Kirschner@itp.uni-leipzig.de

C ontents

1	Introduction	2
2	s'(2)-invariant R-m atrix 2.1 A lgebra s'(2) and lowest weight modules	5
3	The operator R $_{^1}$ $_{^2}$ (u) as integral operator 3.1 Dierential equations	
4	The supersym metric extension: $s'(2j)$ -invariant R -matrix 4.1 The algebra $s'(2j)$ and lowest weights modules	11
5	The operator $R_{\gamma_1\gamma_2}$ (u) as integral operator 5.1 Dierential equations	17
6	C on clusions	20
7	A cknow ledgm ents	20

1 Introduction

Solutions of the Yang-Baxter equation, in particular ones with s'(2) or s'(2jl) symmetry, are used in constructing a number of statistical models [1, 2, 3, 4, 5, 6, 8, 26] and are applied in analyzing the high-energy Regge [16, 17, 18, 22] and B jorken asymptotics [19, 20, 21] of four-dimensional gauge theories.

In the previous paper [9] we have studied the s'(2jl) sym m etric rational solutions of the Yang-Baxter equation. We have obtained the general R-operator acting on the tensor product of two arbitrary representation spaces of lowest weight in terms of the matrix elements in a basis of lowest weight vectors appearing in the tensor product.

The lowest weight representation spaces have been chosen in terms of polynomials in one even variable z and two odd variables; Then the tensor product representation consists of polynomial two-point functions, i.e. functions of two sets $(z_i; i; i)$. This formulation is the appropriate one for treating arbitrary in nite dimensional lowest weight representations. It is the natural formulation from the viewpoint of applications to the B jorken limit of four-dimensional eld theories like QCD. In the B jorken asym totics the interaction reduces in dimension electively to a line on the light cone, being the range of the even variable z. The G rassmann variables; appear in the B jorken limit of supersymmetric (N = 1) Yang-Mills theories allowing to unify the gluon and quark elds in terms of super elds. The elective interaction can be formulated in terms of integral operators, the kernels of which are (super)conformal four-point functions.

This physical context provides a motivation to look for a corresponding formulation of the R-operators in terms of integral kernels. Instead of constructing these kernels from the matrix elements derived earlier we present here a direct calculation. We consider the dening

relations for the general R-operator now as dierential equations on the kernel. The essential observation is that this set of dierential equation reduces to simple rst order dierential equations, one for each variable involved. The resulting kernel is expressed in powers of combinations of the variables having simple s'(2jl) transform ation properties.

The method proposed here seems to be general enough to become useful in solving the Yang-Baxter equation with other symmetries. In particular the observed reduction of the dening relation to simple instance of the restorder differential equations is expected to work as a powerful tool in other cases. This reduction relies on the structure of the Lax operator, i.e. the Roperator acting in the tensor product of the fundamental and an arbitrary representation. Its matrix representation decomposes into matrices times derivatives, where these matrices are tensor products of particular vectors. In the case of s'(2) this structure appeared earlier in calculations by L N Lipatov [16] and has been used in [17, 18, 22]. The kernel of Roperator is nally obtained as the similarity transformation transforming these instance retorder differential operators into the simple derivatives with respect to variables z_i ; i; i.

A im ing to give a clear presentation of the method we discuss the simplest case of s '(2) rst. Note that the main idea to represent the universal R-matrix as an integral operator originates from the paper [10]. Moreover, the nalanswer for the kernel of R-operator has been obtained by a straightforward solution of the system of dierential equations of the second order in the paper [11]. See this paper for the applications of the integral representation of R-matrix.

The presentation is organized as follows. In Section 2 we collect the standard facts about the algebra s'(2) and its representations. We represent the lowest weight modules by polynomials in one variable (z) and the s'(2)-generators as instandard i erential operators. We derive the dening relation for the general R-matrix, i.e. the solution of the Yang-Baxter equation acting on tensor products of two arbitrary representations, the elements of which are polynomial functions of z_1 and z_2 . We solve this dening relation in the space of lowest weights. The result is well known [28, 24, 25, 27] but we represent all calculations in the form which is appropriate for the supersymmetric generalisation. In Section 3 we give the simple method of derivation of the integral operator representation for the R-matrix. The main observation is that the dening system of differential equations of the second order for the kernel of R-matrix is equivalent to the system of the rest-order differential equations.

These sections have introductory character and starting from the Section 4 we follow the same strategy for the supersymmetric algebras '(2]1). We represent the s'(2]1) lowest weight modules by polynomials in one even (z) and two odd variables (;) and the s'(2]1)-generators as rst order dierential operators and derive the dening relation for the general R-matrix, i.e. the solution of the Yang-Baxter equation acting on tensor products of two arbitrary representations, the elements of which are polynomial functions of $(z_1; 1; 1)$ and $(z_2; 2; 2)$. We solve this dening relation in the space of lowest weights. All this is the short summary of our previous paper [9] where the reader cannot all details.

In Section 5 we reduce the de ning system of di erential equations of the second order for the kernel of R-m atrix to the equivalent system of the rst-order di erential equations. Next we demonstrate by direct calculation that the obtained integral operator coincides with ones obtained in our previous paper.

Finally, in Section 6 we sum marize. Appendix A contains some technical details about the construction of superconformal N-point functions. In Appendix B we give the list of superintegrals which are needed for the calculations of matrix elements.

2 s'(2)-invariant R-m atrix

2.1 A lgebra s'(2) and lowest weight modules

The algebra s'(2) has three generators $S = (S^1; S^2; S^3)$ with standard commutation relations:

$$[S^{j};S^{k}] = i^{jkp}S^{p}$$

We shall use the generators S;S

$$S = S^1 ext{ iS}^2 ; S = S^3 ; [S;S] = S ; [S^+;S] = 2S:$$

which have the following form in the fundam ental representation

$$S^{i} = \frac{1}{2}^{i};$$
 $_{1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix};$ $_{2} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix};$ $_{3} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix};$ $S^{+} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix};$ $S = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix};$ $S = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix};$

We represent the generators as sixt order dierential operators, acting on the space of polynomials (z):

$$S = 0; S^{+} = z^{2}0 + 2z; S = z0 + z$$
 (2.1.1)

Below we shall use the global form of s'(2)-transform ations

$$e^{S}$$
 (z) = (z); $e^{S^{+}}$ (z) = $\frac{1}{(1-z)^{2}}$ $\frac{z}{1-z}$ (2.1.2)

The lowest weight s'(2)-module V, is built on the lowest weight vector obeying:

$$S = 0; S = `$$

The center of the enveloping algebra of the algebra s'(2) is generated by the C asim ir operator C_2 and the module is characterised uniquely by the action of the C asim ir operators on its elements:

$$C_2 = S^2 S + S^+ S ; C_2 v = '(' 1)v:$$
 (2.1.3)

It is a vector space spanned by the following basis $V_k = (S^+)^k$; $k \ 2 \ Z_+ : W$ e shall use the above realization of the s '(2)-generators as the di-erential operators of instance on the in nite-dimensional (for generic ') space V_* of polynomials (z) of variable z. The coherent state e^{S^+} for the lowest weight = 1 is the generating function, the power expansion in of which produces the basis:

$$e^{S^{+}} 1 = (1 z)^{2}; V_{k} = (2)_{k} z^{k}$$
 (2.1.4)

There exist some special values of ': '= n; n 2 $\frac{1}{2}Z_+$, for which the module V_* becomes a nite-dimensional vector space. Indeed it is evident from (2.1.4) that all basis vectors are equal zero for k n+1. Let us introduce the special notation for the fundamental s '(2)-module: $V_{\frac{1}{2}}$ V_f .

The tensor product of two s'(2)-m odules has the well known direct sum decomposition:

$$V_{1} V_{2} = \sum_{n=0}^{x^{2}} V_{1+2+n}$$
 (2.1.5)

For the proof of (2.1.5) in the generic situation one has to determ ine all possible lowest weight vectors appearing in the tensor product $V_1 - V_2$. In the realization on functions of z the space $V_1 - V_2$ is isomorphic to the space of polynomials in two variables $(z_1;z_2)$. The s'(2)-generators acting on the $V_1 - V_2$ are just the sum s of corresponding generators acting in V_1 . The lowest weight vectors of the irreducible representations in the decomposition of $V_1 - V_2$ are defined as the solutions of the equation:

$$S = 0$$
) $(z_1; z_2) = (z_1 z_2)$ (2.1.6)

Now, the lowest weight vectors in the decomposition of the tensor product are constructed from functions (2.1.6) being eigenfunctions of generator S. The eigenfunctions of the operator S are homogeneous polynomials of degree n; n 2 $\rm Z_+$ and nally we obtain that all lowest weights in the space $\rm V_1$ $\rm V_2$ are:

$$_{n}$$
 $(z_{1} z_{2})^{n}$ (2.1.7)

2.2 Yang-Baxter equation and general operator R₁, (u)

Let V_{i_1} ; i=1;2;3 be three lowest weight s`(2)-m odules. Let us consider the three operators $R_{i_1i_2}(u)$ which are acting in $V_{i_1} - V_{i_2}$ and obey the Yang-Baxter equation in the space $V_{i_3} - V_{i_1} - V_{i_2}$:

$$R_{i_3i_1}(u)R_{i_3i_2}(v)R_{i_1i_2}(v-u) = R_{i_1i_2}(v-u)R_{i_3i_2}(v)R_{i_3i_1}(u)$$
 (2.2.1)

To obtain the de ning relation for the general R-operator we consider the special case $'_1 = f$ in (2.2.1)

$$R_{f_1}(u)R_{f_2}(v)R_{f_1}(v) = R_{f_2}(v)R_{f_1}(u)$$
: (2.2.2)

W e can choose the m atrix realization in V_f and use the operator (S are generators acting in V_{ν})

$$R_{f}(u) = u + \frac{1}{2} + S = u + \frac{1}{2} + S S S + u + \frac{1}{2} S$$
; (2.2.3)

which is the well known solution of the Yang-Baxter equation:

$$R_{f_1f_2}(u \quad v)R_{f_1i_3}(u)R_{f_2i_3}(v) = R_{f_2i_3}(v)R_{f_1i_3}(u)R_{f_1f_2}(u \quad v)$$
; $R_{f_1f_2}(u) = 1 + u \quad P_{2i_3}(u)$

where P_{12} is the permutation. The operators $R_{f'_1}$, $R_{f'_2}$ are linear functions of the spectral parameter u

$$R_{f_{i}}(u) = u + \frac{1}{2} + R_{i}; R_{i} = \begin{cases} S_{i} & S_{i} \\ S_{i}^{+} & S_{i} \end{cases}$$
; i= 2;3:

Now the general R-matrix R $_{1,2}$ (u) acting in the tensor product $V_{1} - V_{2}$ of arbitrary modules, is xed by the condition (222). After separation of v+u and v u dependence we obtain two equations (v u!u):

$$(R_1 + R_2)R_{i_1i_2}(u) = R_{i_1i_2}(u)(R_1 + R_2)$$
 (2.2.4)

$$\frac{u}{2} (R_1 - R_2) + R_1 R_2 - R_{1'2} (u) = R_{1'2} (u) - \frac{u}{2} (R_1 - R_2) + R_2 R_1$$
 (2.2.5)

The rst equation (22.4) expresses the fact that R $_{1}$ $_{2}$ (u) has to be invariant with respect to the action of s (2)-algebra

h i
$$S;R_{i_1i_2}(u) = 0; S = S_1 + S_2$$

and the second equation is the wanted de ning relation for the operator $R_{i_1 i_2}$ (u). In the matrix form the de ning relation reads as follows:

$$K_{AB}R_{112}(u) = R_{112}(u)K_{AB}$$
; A;B = 1;2 (2.6)

where:

$$K = \begin{pmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{pmatrix} \begin{pmatrix} u \\ 2 \end{pmatrix} (R_1 & R_2) + R_1 R_2 ; K \begin{pmatrix} u \\ 2 \end{pmatrix} (R_1 & R_2) + R_2 R_1$$
:

Due to s'(2)-invariance there is only one independent equation so that the system of equations (22.4), (22.5) for R-m atrix is equivalent to the system

Indeed, the operators $R_{f_1}(u)R_{f_2}(v)$ and $R_{f_2}(v)R_{f_1}(u)$ are s'(2)—invariant by construction

$$S + \frac{1}{2} \sim R_{f_1}(u) R_{f_2}(v) = 0 ; S + \frac{1}{2} \sim R_{f_2}(v) R_{f_1}(u) = 0$$

so that we obtain the commutation relations between s'(2)-generators and K-operators

$$S_{K}^{i} = \frac{1}{2} [K_{K}^{i}] ; S_{K}^{i} = \frac{1}{2} \sim K_{K} :$$

For example

$$S^{+}$$
; $K_{12} = K_{11}$ K_{22} ; S^{+} ; $K_{11} = K_{21}$; S^{+} ; $K_{22} = K_{21}$; S^{+} ; $K_{21} = 0$

and it is possible to obtain the full system of equations (2.2.6) starting from the K $_{12}$ -equation in (2.2.7).

2.3 Eigenvalues of operator R_{1,2} (u)

The s'(2)-invariance of the operator R $_{^1}$ $_{^2}$ (u) allows to simplify the eigenvalue problem. Due to s'(2)-invariance any eigenspace of the operator R $_{^1}$ $_{^2}$ is a lowest weight s'(2)-module generated by some lowest weight eigenvector. Therefore without loss of generality we can solve the dening relation (2.2.5) in the space of lowest weights. Let us consider in more details the structure of eigenspace of the s'(2)-invariant operator acting on the tensor product V_{1} V_{2} . As we have seen from the direct sum decomposition (2.1.5) for any xed nother space of lowest weight vectors is one-dimensional and therefore the operator R $_{1}$ $_{2}$ is diagonal on lowest weight vectors $_{1}$:

$$R_{1} \cdot (u) = R_{n} \cdot (2.3.1)$$

The relation (2.2.7) leads to a sim ple recurrence relation for the eigenvalues R $_{\rm n}$. The operator K $_{12}$ com m utes w ith the low ering generator S $_{12}$ and therefore K $_{12}$ m aps low est w eight vectors to low est w eight vector and decrease its power by one:

$$K_{12} = n = n (u) = n (u + '1 + '2 + n = 1);$$

where the concrete form of the function $_{n}$ (u) is derived by the explicit calculation using

$$K_{12} = (z_1 z_2)@_1@_2 \frac{u}{2} z_2 Q_1 + \frac{u}{2} z_2 z_2 z_2)^n$$
:

Under the permutation:

the operators K_{AB} and lowest weights transform as follows:

so that the action of operator K $_{12}$ on lowest weight vectors can be obtained from formulae for K $_{12}$ by the form alsubstitution $`_1;z_1$ \$ $`_2;z_2$ and u \$ u:

$$K_{12} = n = n (u)_{n=1}$$
:

W e project the operator equation K $_{12}R = RK_{12}$ onto the lowest weight vectors $_n$:

$$K_{12}R_{n} = RK_{12}_{n} =)$$
 $_{n}(u)R_{n}_{n-1} = R_{n-1}_{n}(u)_{n-1}$

which results in the recurrent relation:

$$_{n}(u)R_{n} = _{n}(u)R_{n-1}$$

with the general solution [28, 24, 25, 27]:

$$R_{n}(u) = (1)^{n} \frac{(u + i_{1} + i_{2})}{(u + i_{1} + i_{2})} \frac{(u + i_{1} + i_{2} + n)}{(u + i_{1} + i_{2} + n)}$$
(2.3.2)

where we xed the overall normalisation of R-operator in such a way that R:17 1.

3 The operator R 1/2 (u) as integral operator

It is useful to derive the explicit representation of the R-m atrix as integral operator acting on polynom ials $(z_1; z_2)$ 2 V_{i_1} V_{i_2}

$$\mathbb{R}_{1,2}(u) \] (z_1; z_2) = dz_3 dz_4 \mathbb{R} (z_1; z_2; z_3; z_4) (z_3; z_4)$$
(3.0.1)

We suppose that it is possible to integrate by parts so that the equations (22.4), (22.5) result in the set of dierential equations for the kernel R $(z) = R(z_1; z_2; z_3; z_4)$

$$(R_1 + R_2 + R_3 + R_4)R$$
 (z) = 0 (3.0.2)

$$\frac{u}{2}(R_1 - R_2) + R_1R_2 - R_2(z) = \frac{u}{2}(R_4 - R_3) + R_4R_3 - R_2(z)$$
 (3.0.3)

where the matrices R_i ; i = 3;4 arise after integration by parts

$$R_{i} = \begin{pmatrix} S_{i} & S_{i} \\ S_{i}^{+} & S_{i} \end{pmatrix} = \begin{pmatrix} z_{i} \theta_{i} + \mathbf{i}_{i} & \theta_{i} \\ z_{i}^{2} \theta_{i} + 2 \mathbf{i}_{i} z_{i} & z_{i} \theta_{i} \end{pmatrix}; \mathbf{i} \quad \mathbf{i}$$

First we shall solve this system of dierential equations and then specify selfconsistently the contours of integration. The main system of equations can be simplied drastically.

Proposition 1 The system of dierential equations (3.0.2), (3.0.3) of the second order is equivalent to the system of dierential equations of the rst order

$$D_1R = 0$$
; $D_2R = 0$; $D_3R = 0$; $D_4R = 0$: (3.0.4)

where operators D i are de ned in a following way

$$D_{1} = \theta_{1} + \frac{u + \frac{1}{1} + \frac{1}{4}}{z_{14}} = \frac{u + \frac{1}{4} + \frac{1}{1}}{z_{12}}; D_{2} = \theta_{2} + \frac{u + \frac{1}{2} + \frac{1}{3}}{z_{23}} + \frac{u + \frac{1}{3} + \frac{1}{2}}{z_{12}}$$
(3.0.5)
$$D_{3} = \theta_{3} = \frac{u + \frac{1}{2} + \frac{1}{3}}{z_{23}} = \frac{u + \frac{1}{2} + \frac{1}{3}}{z_{34}}; D_{4} = \theta_{4} = \frac{u + \frac{1}{1} + \frac{1}{4}}{z_{14}} + \frac{u + \frac{1}{1} + \frac{1}{4}}{z_{34}};$$

The s'(2)-generators are expressed in term s of D_i as follows:

$$S = \bigcup_{i=1}^{X^{4}} D_{i}; S = \bigcup_{i=1}^{X^{4}} z_{i}D_{i}; S^{+} = \bigcup_{i=1}^{X^{4}} z_{i}^{2}D_{i}; S = \bigcup_{i=1}^{X^{4}} S_{i}$$

The common solution of the system (3.0.4) up to overall normalisation is

R (z)
$$z_{12}^{u}^{1+\frac{1}{4}}z_{14}^{u}^{1+\frac{1}{4}}z_{32}^{u}^{2+\frac{1}{2}}z_{34}^{u+\frac{1}{2}}$$
 '3 $z_{34}^{u+\frac{1}{2}}$ '3 ; '3 = 1 '1; '4 = 1 '2:

3.1 Dierential equations

There exists the simple and elegant way to derive the equation of the rst order for the R-m atrix which is especially useful for the supersym metric generalisation. The similar trick was used in [16, 17, 18, 22] for the derivation of integrals of motion for $X \times X$ spin chains. The matrix R has very special structure

$$R = \begin{pmatrix} z & 1 \\ z^2 & z \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 2 & z \end{pmatrix} + \begin{pmatrix} z & 1 \\ z^2 & z \end{pmatrix} = \begin{pmatrix} 1 & z & 1 \\ z & z & 1 \end{pmatrix}$$

so that the matrix

$$Rf(z) = \begin{cases} z & 1 \\ z^2 & z \end{cases} f^0(z) + \begin{cases} 0 & 0 \\ 2 & z \end{cases} f(z)$$

has two evident eigenvectors

$$z = 1 Rf(z) = z = 1 f(z); Rf(z) = z = z = 1 f(z)$$

where f(z) is an arbitrary function. The derivative $f^0(z)$ disappears from the eigenvalues and we shall use this property to derive the rst order di erential equation for the kernel R(z). Let us multiply the equations (2.2.4), (2.2.5) to the vector z_4 1 from the left and to the vector $\frac{1}{z_2}$ from the right. Then we obtain the system of two rst-order di erential equations for the function R(z)

$$D_1R + D_3R = ('_2 \quad '_4)z_{24}R$$
; $\frac{u}{2} + '_2 D_1R + \frac{u}{2} + '_4 D_3R = \frac{u}{2}('_2 \quad '_4)z_{24}R$

which is equivalent to the system

$$D_1R = (u + '_4) z_{24}R$$
; $D_3R = (u + '_2) z_{24}R$: (3.1.6)

W e use the notation

$$D_k$$
 z_4 1 R_k $\frac{1}{z_2}$ = $z_{k2}z_{k4}\theta_k$ $k(z_{k2}+z_{k4})$; $k=1;3$

for the arising di erential operators. Let us multiply the equations (22.4), (22.5) to the vector z_1 1 from the left and to the vector $\frac{1}{z_3}$ from the right. In the same manner we obtain the mirror system of equations:

$$D_2R = (u + '_3) z_{13}R$$
; $D_4R = (u + '_1) z_{13}R$; (3.1.7)

w here

$$D_k$$
 z_1 1 R_k $\frac{1}{z_3}$ = $z_{k1}z_{k3}\theta_k$ $k(z_{k1}+z_{k3})$; $k=2;4$:

Finally, as consequence of the main system of dierential equations of the second order we obtain the system of dierential equations (3.1.6), (3.1.7) of the rst order. The eqs. (3.1.6), (3.1.7) are equivalent to

$$D_1R = 0$$
; $D_2R = 0$; $D_3R = 0$; $D_4R = 0$:

where the dierential operators D $_{\rm i}$ are dened in (3.0.5).

The next step is to prove the reverse, that the system of equations (3.0.2), (3.0.3) is the consequence of the system (3.0.4). It is easy to rewrite the s'(2)-generators and the equations of s'(2)-invariance (3.0.2) in term s of the introduced operators D_i:

$$S = \sum_{i=1}^{X^{4}} S_{i} = \sum_{i=1}^{X^{4}} D_{i}; S = \sum_{i=1}^{X^{4}} S_{i} = \sum_{i=1}^{X^{4}} z_{i}D_{i}; S^{+} = \sum_{i=1}^{X^{4}} S_{i}^{+} = \sum_{i=1}^{X^{4}} z_{i}^{2}D_{i};$$

$$S R = \sum_{i=1}^{X^{4}} D_{i}R = 0; SR = \sum_{i=1}^{X^{4}} z_{i}D_{i}R = 0; S^{+}R = \sum_{i=1}^{X^{4}} z_{i}^{2}D_{i} = 0$$

As explained above due to s'(2)-invariance we can restrict to one K $_{12}$ -equation in (22.7). This equation has the explicit form

$$z_{12}@_{1}@_{2}R (\mathbf{z}) + z_{34}@_{3}@_{4}R (\mathbf{z}) =$$

$$= \frac{u}{2} \quad \mathbf{z} \quad @_{1}R (\mathbf{z}) + \frac{u}{2} \quad \mathbf{z} \quad \mathbf{z} \quad & \frac{u}{2} \quad \mathbf{z} \quad & \frac{u}{2} \quad & \mathbf{z} \quad & \frac{u}{2} \quad & \mathbf{z} \quad &$$

and can be easily rewritten in terms of the operators D $_{
m i}$

$$z_{12}D_{1}D_{2}R + z_{34}D_{3}D_{4}R =$$

$$= \frac{u}{2} + \frac{1}{3} + (u + \frac{1}{2} + \frac{1}{3})\frac{z_{12}}{z_{23}} D_{1}R \qquad \frac{u}{2} + \frac{1}{4} (u + \frac{1}{1} + \frac{1}{4})\frac{z_{12}}{z_{14}} D_{2}R +$$

$$+ \frac{u}{2} + \frac{1}{1} (u + \frac{1}{1} + \frac{1}{4})\frac{z_{34}}{z_{14}} D_{3}R \qquad \frac{u}{2} + \frac{1}{2} + (u + \frac{1}{2} + \frac{1}{3})\frac{z_{34}}{z_{23}} D_{4}R :$$

We see that the system of equations (3.0.2), (3.0.3) is the consequence of the system (3.0.4) and nally, the system of the de ning equation (2.2.7) is equivalent to the system (3.0.4). The common solution of the system (3.0.4) up to an overall normalisation is

R (z)
$$z_{12}^{u}$$
 '1 + '4 z_{14}^{u} '1 '4 z_{32}^{u} '2 '3 z_{34}^{u+2} '3; '3 = 1 '1; '4 = 1 '2:

The operators D_i are obtained by the sim ple sim ilarity transform ation from the derivatives Q_i

so that by using the ansatz

R (z)
$$z_{12}^{u} \dot{z}_{14}^{u} \dot{z}_{14}^{u} \dot{z}_{14}^{u} \dot{z}_{32}^{u} \dot{z}^{u} \dot{z}_{34}^{u} \dot{z}_{34}^{u} \dot{z}_{3}$$
 R (z)

the system of equations D_iR (z) = 0 can be reduced to the system $\theta_i R(z) = 0$ with the evident solution R (z) = const.

3.2 The integral representation.

It remains to x the contours of integrations. The R-operator acts on the space of polynom ials $(z_1; z_2)$ which can be of arbitrary large degree. The requirement of convergence allows compact contours only and there remains the one possibility:

$$[R_{12}(u)](z_{1};z_{2}) = \begin{cases} Z_{23} & Z_{23} \\ dz_{3} & dz_{4}z_{21}^{1+u} & 2z_{41}^{2} & 2z_{41}$$

where the normalisation factor is chosen so that R:17 1,

$$= \frac{1}{('_1 \quad '_2 \quad u) \quad ('_2 \quad '_1 \quad u)} \frac{('_1 + '_2 \quad u)}{('_1 + '_2 + u)}; \tag{32.9}$$

It is easy to check a posteriori that the obtained integral operator is s'(2)-invariant and has the eigenvalues (2,3,2). The eigenvalues can be calculated by using twice the general form ula

Z $_{z_3}$ $dz_4z_{34}^a$ $^1z_{41}^b$ $^1=z_{31}^{a+b}$ 1 (a) (b) $\overline{}$:

The supersym metric extension: s'(2jl)-invariant R-matrix

The algebra s'(21) and lowest weights modules

Here we collect the main formulas about algebra s'(2]1) (for details see [12, 13, 14, 15, 9]). We represent the generators as sixt order dierential operators, acting on the space of polynomials (z; ;):

$$S = \emptyset; V = \emptyset + \frac{1}{2} \emptyset; W = \emptyset + \frac{1}{2} \emptyset; \qquad (4.1.1)$$

$$V^{+} = z \emptyset + \frac{1}{2} z \emptyset + \frac{1}{2} \emptyset \qquad ('b); W^{+} = z \emptyset + \frac{1}{2} z \emptyset + \frac{1}{2} \emptyset \qquad ('+b);$$

$$S^{+} = z^{2} \emptyset + z \emptyset + z \emptyset + z 'z b ; \qquad (4.1.2)$$

$$S = z \emptyset + \frac{1}{2} \emptyset + \frac{1}{2} \emptyset + \frac{1}{2} \emptyset + '; B = \frac{1}{2} \emptyset \quad \frac{1}{2} \emptyset + b:$$

The lowest weight s'(2jl)-module $V_{ib} = V_{r}$; r = (ib) is built on the lowest weight vector obeying:

$$V = 0; W = 0; S = 0; S = '; B = b$$

The module V_{κ} is a vector space spanned by the following basis [12], [14] with the even vectors

$$A_k = (S^+)^k$$
; $B_k = (S^+)^{k-1}W^+V^+$; $k \ 2 \ Z_+$

and the odd vectors

$$V_k = (S^+)^k V^+$$
; $W_k = (S^+)^k W^+$; k 2 Z₊

We shall use the above realization of the s'(2jl)-generators as the dierential operators of rst order acting on the in nite-dimensional (for generic') space V_{ij} of polynomials (z; ;) of variables z; ; .

The tensor product of two s'(2jl)-modules has the following direct sum decomposition [14,9]:

$$V_{1,b_{1}} V_{2,b_{2}} = V_{2,b} + 2 V_{1,b} + 2 V_{1,b_{1}} + V_{1,b_{2}} + V_{1,a_{1},b_{2}} + V_{1,a$$

Note that this formula is applicable in the generic situation $i_1 \in b_1$. All possible lowest weight vectors appearing in the tensor product V_{i_1,i_2} V_{i_2,i_2} are divided in two sets, the even lowest weights:

$$_{n}$$
 Z_{12} $\frac{1}{2}_{12}^{12}_{12}^{12}$; $D_{1}_{n} = 0$; $S_{n} = (n + ')_{n}$; $B_{n} = b_{n}$ (4.1.4)

and the odd lowest weights:

$$_{n}$$
 $_{12}Z_{12}^{n}$; $_{n}^{+}$ $_{12}Z_{12}^{n}$; $S_{n} = (n + ' + \frac{1}{2})_{n}$; $B_{n} = (b + \frac{1}{2})_{n}$ (4.1.5)

where

$$Z_{ij}$$
 z_i $z_j + \frac{1}{2}(i_{j} + i_{j}); i_{j}$ i_{j} ; i_{j} i_{j} ; i_{j} :

4.2 Yang-Baxter equation and general operator $R_{\tau_1\tau_2}$ (u)

Let V_{i,b_i} ; i = 1;2;3 be three lowest weight s'(2jl)-m odules. We shall use the short-hand notation:

$$f' = \frac{1}{2}; \frac{1}{2} ; r' = (';b) ; V_r = V_{';b}$$
:

Let us consider the three operators $R_{\gamma_i \gamma_j}(u)$ which are acting in $V_{\gamma_i} = V_{\gamma_j}$ and obey the Yang-Baxter equation in the space $V_{\gamma_i} = V_{\gamma_j} = [25]$:

$$R_{\gamma_{3}\gamma_{1}}(u)R_{\gamma_{3}\gamma_{2}}(v)R_{\gamma_{1}\gamma_{2}}(v-u) = R_{\gamma_{1}\gamma_{2}}(v-u)R_{\gamma_{3}\gamma_{2}}(v)R_{\gamma_{3}\gamma_{1}}(u) \tag{4.2.1}$$

To obtain the de ning relation for the general R-operator we consider the special case $^{\circ}_{3} = f^{\circ}$ in (4.2.1). Then one can choose the matrix realization [26, 7, 9] in V_{∞}

and the operators R_{f_1} , R_{f_2} are linear functions of spectral parameter u in this particular case

$$R_{f_{1}}(u) = u + R_{i}; R_{i} = \begin{pmatrix} 0 & & & & 1 \\ & S_{i} + B_{i} & W_{i} & S_{i} \\ & V_{i}^{+} & 2B_{i} & V_{i} & A ; i = 1; 2 \\ & S_{i}^{+} & W_{i}^{+} & B_{i} & S_{i} \end{pmatrix}$$

Now the general R-m atrix R $_{\tau_1\tau_2}$ (u) acting in the tensor product V_{τ_1} V_{τ_2} of arbitrary modules, is xed by the condition

$$R_{f_{3}}(u)R_{f_{3}}(v)R_{f_{3}}(v) = R_{f_{3}}(v)R_{f_{3}}(v)R_{f_{3}}(u)$$
 (4.2.2)

or equivalently:

$$(R_1 + R_2)R_{\gamma_1\gamma_2}(u) = R_{\gamma_1\gamma_2}(u) (R_1 + R_2)$$
 (4.2.3)

$$\frac{u}{2}(R_1 - R_2) + R_1R_2 - R_{\gamma_1\gamma_2}(u) = R_{\gamma_1\gamma_2}(u) - \frac{u}{2}(R_1 - R_2) + R_2R_1$$
 (4.2.4)

The rst equation (2.2.4) expresses the fact that R (u) has to be invariant with respect to the action of s'(2jl)-algebra and the second equation is the dening relation for the operator $R_{\gamma,\gamma}$ (u).

4.3 Eigenvalues of operator R, (u)

Due to s'(2jl)—invariance any eigenspace of the operator R $_{1,2}$ is a lowest weight s'(2jl)—module generated by some lowest weight eigenvector. As we have seen from direct sum decomposition (4.1.3) for every xed n the space of lowest weight vectors with eigenvalue b is two-dimensional and the ones with eigenvalues b $\frac{1}{2}$ are one-dimensional. Therefore the operator R $_{1,2}$ is diagonal on odd lowest weight vectors $_{n}^{+}$ and $_{n}^{-}$ but acts non-trivially on the two-dimensional subspace of even lowest weight vectors spanned on $_{n}^{+}$ and $_{n}^{-}$. In matrix form we have:

The m atrix relation (2.2.5) leads to a set of recurrence relations for the coe cients A $_n$; :::; E $_n$ with the general solution [9]:

$$A_{n}(u) = (1)^{n+1} \frac{(u + \mathbf{i}_{n})}{(u + \mathbf{i}_{n} + 1)}$$

$$\frac{(\mathbf{i}_{1} \ b_{1}) (\mathbf{i}_{1} + \mathbf{b}_{1}) (u \ b_{1} \ b_{2}) \ (u + \mathbf{b}_{1} + \mathbf{b}_{2}) (u \ b_{1} \ \mathbf{i}_{2}) (u \ b_{1} \ \mathbf{i}_{2})}{(\mathbf{i}_{2} \ b_{2}) (\mathbf{i}_{1} + \mathbf{b}_{1})}$$

$$B_{n}(u) = (1)^{n} \frac{(u + \mathbf{i}_{n})}{(u + \mathbf{i}_{n})}; C_{n}(u) = (1)^{n} \frac{(u + \mathbf{i}_{n} + 1)}{(u + \mathbf{i}_{n} + 1)} \frac{(\mathbf{i}_{2} + \mathbf{b}_{2}) (\mathbf{i}_{1} \ b_{1})}{(\mathbf{i}_{2} \ b_{2}) (\mathbf{i}_{1} + \mathbf{b}_{1})}$$

$$D_{n}(u) = (1)^{n+1} \frac{(u + \mathbf{i}_{n} + 1)}{(u + \mathbf{i}_{n})} \frac{u + b_{2} \ b_{1}}{(\mathbf{i}_{2} \ b_{2}) (\mathbf{i}_{1} + \mathbf{b}_{1})}$$

$$E_{n}(u) = (1)^{n} \frac{(u + \mathbf{i}_{n} + 1)}{(u + \mathbf{i}_{n} + 1)} \frac{(u + \mathbf{b}_{2} \ \mathbf{i}_{1}) (u + \mathbf{b}_{2} + \mathbf{i}_{1})}{(\mathbf{i}_{2} \ b_{2}) (\mathbf{i}_{1} + \mathbf{b}_{1})}$$

$$F_n(u) = (1)^n - \frac{(u + i_n + 1)}{(u + i_n + 1)} - \frac{(u b_1 i_2)(u b_1 + i_2)}{(i_2 b_2)(i_1 + b_1)}$$

where we used the notations:

$$n + 1 + 2 = u + b_2 + b_1$$

As usual the obtained general solution of the Yang-Baxter equation is xed up to overall normalization. We choose the normalization such that the R-matrix coincides with the permutation operator for u=0 and $\tilde{\gamma}_1=\tilde{\gamma}_2$. Note that the transformation from our basis vectors n to the ones diagonalizing the R-matrix depends on n and this does not result in simpler formulae. ¹

5 The operator R_{3,7} (u) as integral operator

Let us go to the derivation of the explicit representation for the R-m atrix as integral operator acting on polynomials ($Z_1;Z_2$) 2 V_{τ_1} V_{τ_2}

h i
$$Z Z$$
 Z $R_{\gamma_1 \gamma_2} (u) (Z_1; Z_2) = dZ_3 dZ_4 R (Z_1; Z_2; Z_3; Z_4) (Z_3; Z_4) (5.0.1)$

where Z fz; ; g and we use the standard superintegration

$$Z$$
 Z Z Z dz dz d d :

We suppose that it is possible to integrate by parts so that the equations (4.2.3), (4.2.4) give the set of dierential equations for the kernel R (\mathbb{Z}) = R (\mathbb{Z}_1 ; \mathbb{Z}_2 \mathbb{Z}_3 ; \mathbb{Z}_4)

$$(R_1 + R_2 + R_3 + R_4)R(Z) = 0$$
 (5.0.2)

$$\frac{u}{2} (R_1 - R_2) + R_1 R_2 - R (\tilde{Z}) = -\frac{u}{2} (R_4 - R_3) + R_4 R_3 - R (\tilde{Z})$$
 (5.0.3)

where the matrices R_i ; i = 3;4 arise after integration by parts

$$R_{i}(u) = \begin{pmatrix} 0 & & & & & 1 \\ & S_{i} + B_{i} & W_{i} & S_{i} & & & & \\ & V_{i}^{+} & 2B_{i} & V_{i} & A ; \Upsilon_{3} & \Upsilon_{1} = (\Upsilon_{1}; b_{1}); \Upsilon_{4} & \Upsilon_{2} = (\Upsilon_{2}; b_{2}) \\ & S_{i}^{+} & W_{i}^{+} & B_{i} & S_{i} & & & \end{pmatrix}$$

$$R_{x_1,x_2}(u \quad v)R_{x_1,x_2}(u)R_{x_2,x_2}(v) = R_{x_2,x_2}(v)R_{x_1,x_2}(u)R_{x_2,x_2}(u \quad v)$$

have choosen $^{\mathbf{q}}_{3} = \mathbf{f}^{\mathbf{r}}$ and obtained

$$R_{3,3}$$
 (u v) $R_{3,5}$ (u) $R_{3,5}$ (v) = $R_{3,5}$ (v) $R_{3,5}$ (u) $R_{3,5}$ (u v)

Then we have \autom atically "used the ansatz for R_{f} , instead of R_{f} . As a consequence the results for the matrix elements formulated there are referring actually to the operator R_{f} , (u)

$$R_{\gamma_2\gamma_1}\left(u\right) - PR_{\gamma_1\gamma_2}\left(u\right)P_{\gamma_1 \, , \, \gamma_2} \; ; \; P \quad (Z_1;Z_2) \qquad (Z_2;Z_1)$$

In the simplest s'(2)-case these matrices coincide $R_{i_1i_2}$ (u) = $R_{i_2i_1}$ (u): In the case of superalgebra s'(2]1) the situation is unusual because the ip-operator P acts nontrivially in the lowest weights basis we have used. The correct answer in selfconsistent notations is given in (4.3.2).

 $^{^{1}}$ N ote that we have adm itted some inconsistency in notations in our previous work [9]. We have started from the general Y ang-B axter equation

First we shall solve this system of dierential equations and then specify selfconsistently the contours of integration.

Proposition 2 The system of dierential equations (5.0.2), (5.0.3) of second order is equivalent to the system of dierential equations of rst order

$$D_{iR} = 0$$
; $D_{iR} = 0$; $i = 1;2;3;4$: (5.0.4)

where the operators D $_{i}$; D $_{i}$ are de ned in a following way

This system of twelve equations is invariant with respect the following discrete symmetry transform ations:

$$1 \$ 4; 2 \$ 3; (`1;b_1) \$ (`2; b_2);$$

 $1 \$ 2; 3 \$ 4; \$; D^+ \$ D ; (`1;b_1) \$ (`2; b_2):$

5.1 Dierential equations

We use the same trick as in the s'(2)-case and derive the simple rst order dierential equations for the kernel of R-operator. The matrix R has a very special structure

w here

$$D^{+} = 0 + \frac{1}{2} 0; D = 0 + \frac{1}{2} 0;$$

so that the matrix Rf(Z), where f(Z) is an even function, has two evident eigenvectors

$$z = \frac{1}{2}$$
 1 Rf = ('b)f $z = \frac{1}{2}$ 1

0 1 0 1

Rf @ A = ('+b)f @ A

 $z + \frac{1}{2}$ $z + \frac{1}{2}$

and all derivatives $(f;D^+f;D^-f)$ disappear from the relations. Moreover, there are two other vectors with the properties

1 0 Rf = 2bf 1 0 + z
$$\frac{1}{2}$$
 1 D + f
0 1 0 1 0 1 0 1
Rf@ 1 A = 2bf @ 1 A + @ A D f: $z + \frac{1}{2}$

We shall derive the rst order dierential equations for the kernel R(Z) by multiplying the equations (42.3), (42.4) by the appropriate vectors from the left and from the right. Let us start from the three variants for the choose of the such vectors. First we obtain the system of two rst-order dierential equations for the function R(Z)

$$D_{1}R + D_{3}R = ('_{2} b_{2} '_{4} + b_{4})Z_{24}^{+}R$$

$$\frac{u}{2} + '_{2} + b_{2} D_{1}R + \frac{u}{2} + '_{4} b_{4} D_{3}R = \frac{u}{2}('_{2} '_{4} + b_{2} + b_{4})Z_{24}^{+}R$$

which is equivalent to the system

$$D_1R = (u + '_4 b_4) Z_{24}^+ R$$
; $D_3R = (u + '_2 + b_2) Z_{24}^+ R$: (5.1.9)

W e use the notation

$$= Z_{k2}Z_{k4}^{+}\theta_{k} + Z_{k4}^{+} k_{2}D_{k} + Z_{k2}^{+} k_{4}D_{k}^{+} \quad (k + b_{k})Z_{k4}^{+} \quad (k + b_{k})Z_{k4}^{+} \quad (k + b_{k})Z_{k2}^{+} + 2b_{k}^{+} k_{2} ; k = 1;3$$

for the arising di erential operators. Secondly we obtain the system of di erential equations

$$B_{1}R + B_{3}R = (\mbox{$^{\prime}_{4}$} & b_{4} & 2b_{2})_{24}R + Z_{24}^{+}D_{2}R$$

$$\frac{u}{2} + 2b_{2} B_{1}R + \frac{u}{2} + \mbox{$^{\prime}_{4}$} & b_{4} B_{3}R = \frac{u}{2}(\mbox{$^{\prime}_{4}$} & b_{4} 2b_{2})_{24}R \frac{u}{2}Z_{24}^{+}D_{2}R \quad D_{1}D_{2}R$$
 which is equivalent to the system

$$B_1R = (u + v_4 b_4)_{24}R$$
; $B_3R = (u + 2b_2)_{24}R + Z_{24}^+D_2R$; (5.1.10)

by conditions (5.1.9) and where

=
$$Z_{k4}^{+}$$
 k_{2} θ_{k} Z_{k4}^{+} D_{k} k_{2} k_{4} D_{k}^{+} + ('k + bk) k_{2} ('k bk) k_{3}) k_{2} ; k = 1;3:

The last system is analogous to the previous ones:

$$C_1R + C_3R = (^{1}_{2} + b_2 + 2b_4)_{24}R + Z_{24}^{+}D_{4}^{+}R$$

$$\frac{u}{2} + \frac{u}{2} + b_2 + C_1 R + \frac{u}{2} + 1 + 2b_4 + C_3 R = \frac{u}{2} (^{1}_{2} + b_2 + 2b_4)_{24} R + \frac{u}{2} Z_{24}^{+} D_{4}^{+} R + D_{4}^{+} D_{3} R$$

and is equivalent to the system

$$C_3R = (u + v_2 + b_2)_{24}R$$
; $C_1R = (u 2b_4)_{24}R + Z_{24}^+D_4^+R$; (5.1.11)

by conditions (5.1.9). The operators C_k are

=
$$Z_{k2} k4 Q_k + k4 k2 D_k + Z_{k2} D_k^+$$
 ('k + bk) 24 ('k bk) k2; k = 1;3:

The obtained system of the six equations can be transformed to the equivalent form. The D $_{1;3}$ -equations (5.1.9) are equivalent to the equations D $_1R=0$; D $_3R=0$. The B $_1$ -equation from (5.1.10) can be transformed to the equation D $_1R=0$ by the condition D $_1R=0$. The C $_3$ -equation from (5.1.11) can be transformed to the equation D $_3R=0$ by the condition D $_3R=0$. The remaining B $_3$; C $_1$ -equations are equivalent to the more involved equations

$$Z_{34}^{+}$$
 1 $\frac{32 \ 32}{Z_{32}}$ $D_3 + (u + v_2) \frac{32}{Z_{32}}$ $(u + v_2) \frac{34}{Z_{34}}$ $R = Z_{24}^{+}D_2R + (v_2) \frac{b_2}{24}R$

$$Z_{12}$$
 1 + $\frac{14 14}{Z_{14}}$ D_1^+ + $(u + v_1^* v_2)\frac{14}{Z_{14}}$ $(u v_2 b_1)\frac{12}{Z_{12}}$ $R = Z_{24}^+D_4^+R$ $(v_2 + b_2)_{24}R$

by the conditions $D_1R = 0$; $D_3R = 0$.

It is possible to derive the m irror system of six equations in the same manner by choosing appropriate vectors for multiplication from the left and from the right. In fact, the resulting equations can be obtained from the present ones by the symmetry transformation

1! 4; 2! 3;
$$('_1;b_1)$$
! $('_4;b_4) = ('_2; b_2)$; $('_2;b_2)$! $('_3;b_3) = ('_1; b_1)$

In this way we obtain the equations $D_4R = 0$; $D_2R = 0$ and $D_4R = 0$; $D_3^+R = 0$ and the m irror partners of the m ore involved equations:

$$Z_{21}^{+}$$
 1 + $\frac{32 \ 32}{Z_{32}}$ D_2 + (u + '2 '1) $\frac{32}{Z_{32}}$ (u '1 + b2) $\frac{21}{Z_{21}}$ $R = Z_{31}^{+}D_3R$ ('1 b1) $_{31}R$

$$Z_{43}$$
 1 $\frac{14 \ 14}{Z_{14}}$ $D_4^+ + (u + '_1 \ '_2) \frac{14}{Z_{14}}$ $(u + '_1 + b_2) \frac{43}{Z_{43}}$ $R = Z_{31}^+ D_1^+ R + ('_1 + b_1)_{31} R$:

The system of four involved equations decomposes into two independent systems of two equations. It can be shown that the system which involves the covariant derivatives D_1^+ and D_4^+ is equivalent to the system of the equations $D_1^+R=0$; $D_4^+R=0$ and the system which involves the covariant derivatives D_2^- and D_3^- is equivalent to the system of the equations $D_2^-R=0$; $D_3^-R=0$.

The next step is to prove the reverse, that the system of equations (5.0.2), (5.0.3) is the consequence of the system (5.0.4). It is easy to rewrite the s'(2.jl)-generators and therefore the equations of s'(2)-invariance (5.0.2) in term s of introduced operators D $_{\rm i}$; D $_{\rm i}$:

$$S = X^{4} D_{i}; V = X^{4} D_{i} D_{i}; W = X^{4} D_{i}$$

The last step is absolutely analogous to the s'(2)-case. Due to s'(2jl)-invariance we can use one K $_{13}$ -equation in (5.0.3) and rewrite it in term s of operators D $_{i}$; D $_{i}$. We om it these form ulae for sim plicity.

5.2 Solution of the system of dierential equations

It is useful to transform the main system of equations (5.0.5)-(5.0.8) to a simpler hom ogeneous form. We shall use the following ansatz for this purpose

$$R (\mathcal{I}) = Z_{12}^{a} Z_{14}^{b} Z_{32}^{c} Z_{34}^{d} \qquad 1 + \frac{12 \ 12}{Z_{12}} \qquad 1 + \frac{14 \ 14}{Z_{14}} \qquad 1 + \frac{32 \ 32}{Z_{32}} \qquad 1 + \frac{34 \ 34}{Z_{34}} \qquad R (\mathcal{I})$$

$$a \quad u \quad '_{1} \quad '_{2}; b \quad u \quad '_{1} + '_{2}; c \quad u + '_{1} \quad '_{2}; d \quad u + '_{1} + '_{2}$$

$$= b_{1} \qquad ; \qquad = b_{2} \qquad ; \qquad = b_{2} \qquad b_{1} + :$$

Proposition 3 The system of dierential equations (5.0.5)-(5.0.8) for the function R (\mathbb{Z}) is transformed to the simpler system of dierential equations for the function R (\mathbb{Z})

$$D_{i}R(\tilde{Z}) = 0$$
; $D_{i}R(\tilde{Z}) = 0$; $i = 1;2;3;4$: (5.2.12)

where operators Di; D; are de ned in a following way

where

A
$$\frac{u + 1 + 2}{2}$$
 ; B $\frac{u + 1 + 2}{2} + b_2 + b_1 + 3$; A + B = u + b₂ b₁

The common solution of these twelve equations $D_iR = 0$; $D_iR = 0$ i = 1;2;3;4 is the manifestly superconform al-invariant function (see Appendix A):

$$R(Z) = 1 + A \frac{\frac{3}{12} \frac{3}{12}}{Z_{12}^{3}} \qquad 1 + B \frac{\frac{3}{14} \frac{3}{14}}{Z_{14}^{3}} \qquad (A + B) \frac{\frac{3}{14} \frac{3}{12}}{Z_{14}^{3}} = (52.14)$$

$$= 1 + \frac{\frac{3}{12} \frac{3}{12}}{Z_{12}^{3}} \qquad \frac{\frac{3}{14} \frac{3}{12}}{Z_{14}^{3}} \qquad 1 + \frac{\frac{3}{14} \frac{3}{14}}{Z_{14}^{3}} \qquad \frac{\frac{3}{14} \frac{3}{12}}{Z_{14}^{3}}$$

where

$$Z_{ij}^{k}$$
 $\frac{Z_{ij}}{Z_{ik}Z_{jk}}$; $\frac{k}{ij}$ $\frac{ik}{Z_{ik}}$ $\frac{jk}{Z_{jk}}$; $\frac{k}{ij}$ $\frac{ik}{Z_{ik}}$ $\frac{jk}{Z_{jk}}$

This proposition can be proved by direct calculations. It is remarkable that all involved operators D_i ; D_i can be obtained by the similarity transformation from the simplest derivatives θ_i ; D_i

$$D_i = R @_i R^{-1}$$
; $D_i = R D_i R^{-1}$; $R = R (Z)$:

5.3 The integral representation.

It rem ains to x the contours of integrations. The R-operator acts on the space of polynomials $(Z_1;Z_2)$ which can be of arbitrary large degree. The requirement of convergence allows the compact contours only. The nalrepresentation for the R-m atrix is:

h i
$$Z_{z_2}$$
 Z_{z_3}
 R_{γ_1,γ_2} (u) $(Z_1;Z_2) = dZ_3$ dZ_4R $(Z_1;Z_2;Z_3;Z_4)$ $(Z_3;Z_4)$ (5.3.15)

and

$$R(Z') = Z_{12}^{a} Z_{14}^{b} Z_{32}^{c} Z_{34}^{d} \qquad 1 + \frac{12 \cdot 12}{Z_{12}} \qquad 1 + \frac{14 \cdot 14}{Z_{14}} \qquad 1 + \frac{32 \cdot 32}{Z_{32}} \qquad 1 + \frac{34 \cdot 34}{Z_{34}}$$

$$1 + A \frac{\frac{3}{12 \cdot 12}}{Z_{12}^{3}} \qquad 1 + B \frac{\frac{3}{14 \cdot 14}}{Z_{14}^{3}} \qquad (A + B) \frac{\frac{3}{14 \cdot 12}}{Z_{14}^{3}} \qquad (5.3.16)$$

$$a \quad u \quad 1 \quad 2; \quad b \quad u \quad 1 + 2; \quad c \quad u + 1 \quad 2; \quad d \quad u + 1 + 2$$

$$= b_{1} \qquad ; \qquad = b_{2} \qquad ; \qquad = b_{2} \qquad b_{1} + :$$

The normalisation factor is chosen so that R:17 1

$$= \frac{(c+b+d)}{(b)(c)(d)} \frac{(u \quad '_1 + b_2)(u \quad '_2 \quad b_1)(u + '_1 + '_2)}{u + '_1 + '_2}$$
(5.3.17)

It is possible to check a posteriori that the obtained integral operator has the matrix elements (4.3.1). Using the formulae from Appendix B we derive the formulae for the even basis vectors: $\frac{1}{2}$

ctors:
$$Z_{z_2}$$
 Z_{z_3} Z_{z_3} Z_{z_4} Z_{z_2} Z_{z_3} Z_{z_4} Z_{z_4}

w here

$$R_{++} = (\mathbf{i}_1 + b_1)(\mathbf{i}_2 \quad b_2); R_{+} = \frac{u^3 + u^2(b_2 \quad b_1) \quad u(\mathbf{i}_1^2 + \mathbf{i}_2^2 + 2b_1b_2) + (b_1 + b_2)(\mathbf{i}_1^2 \quad \mathbf{i}_2^2)}{u + \mathbf{i}_1 + \mathbf{i}_2 + n}$$

and

w here

$$R_{+} = (u + b_{2} b_{1}) (u + '_{1} + '_{2} + n) ; R_{-} = \frac{('_{1} b_{1}) ('_{2} + b_{2}) (u + '_{1} + '_{2} + n)}{u + '_{1} + '_{2} + n}$$

$$R_{++} + R_{-+} \dot{j}_{h=0} = R_{+-} + R_{--} \dot{j}_{h=0} = \frac{u + \ \ \dot{j}_{h=0}}{(u - \ \ \dot{j}_{h} + b_{2})(u - \ \ \dot{j}_{h})(u + \ \dot{j}_{h} + \dot{j}_{2})}$$

The analogous formulae for the odd vectors are much simpler:

These formulae for the action of the R-matrix on the basis vectors coincide up to overall normalisation with the ones (4.3.1) obtained in our previous paper [9].

6 Conclusions

We have derived the integral operator form of the rational sl(2jl) sym metric solution of the Yang-Baxter equation acting on the tensor product of two general lowest weight representations $\gamma_1 = (i_1; b_1)$ and $\gamma_2 = (i_2; b_2)$. The lowest weight representations space is identified with the space of polynomials in one even (z) and two odd (;). The kernel of the Roperator is a superconformal four-point function of primary elds with weights $\gamma_1; \gamma_2$ and $\gamma_3 = \gamma_1; \gamma_4 = \gamma_2$. This function of the supercoordinates of four points arises as a simple sum of terms in powers of $Z_{ij}; i_{jj}; i_{jj}$. The simplicity of the result makes this integral operator form useful in applications and further investigations.

Following the well known procedure, the general R-operator is derived from the Yang-Baxter equation involving the latter and the known R-operator $R_{f;}$, acting in the tensor product of the fundamental representation (in matrix form) and an arbitrary representation. We have formulated the conditions involved in this Yang-Baxter relation as dierential equations on the kernel.

There is a remarkable reductions of these equations, being an overdeterm ined system of second order equations, to a simple set of rst order equations. In this way the obtained kernel expresses the similarity transformation relating this consistent system of rst order equations to the trivial one being the condition for a function to be constant in all variables z_i ; i; i. We have obtained this reduction by observing that the Lax matrix representing R_f ; has a peculiar structure implying that appropriate projections reduce the number of derivatives.

We have calculated the matrix elements of the obtained R-operator. The result is consistent with the one obtained earlier from recurrence relations in the framework of the matrix formulation. We have quoted the formulae to be used for calculating the action of the integral operator.

Them ethod of calculation relying on this reduction seems to be general enough to become useful for solving the Yang-Baxter equations with other symmetries. The structure on which the reduction relies deserves further study.

7 Acknowledgments

We thank GK orchem sky and AM anashov for the stimulating discussion and critical remarks. This work has been supported by Deutsche Forschungsgemeinschaft, grant NoKI

623/1-2 and by INTAS, grant No 96-524. One of us (D K.) is grateful to the Germ an M instry BMBF for support. The work of S.D. was partially supported by RFFI-grant No 00-01-005000.

Appendix A

In this Appendix we shall construct the superconform alinvariant N-point function with zero global U (1)-charge. The requirement of s'(2j)-invariance (5.0.2) results in the equations:

$$V R (Z) = 0; W R (Z) = 0; V^{+}R (Z) = 0; W^{+}R (Z) = 0$$

where $V = \bigcup_{i=1}^{P} V_i$ and so on. Note we show the independent equations only. The others follow from the commutation relations

$$fV : W q = S : fV : W q = S B:$$

Let us consider the general case of N-point functions with zero total U (1)-charge: $Z = (Z_1; :::; Z_N)$; $b_i = 0$. It is convenient to use the globals `(2jl)-transform ations which are generated by the lowering operators V; W

$$e^{V}$$
 $(z; ;) = z + \frac{1}{2}; + ; ; e^{W}$ $(z; ;) = z + \frac{1}{2}; ; + :$

and the rising operators V^+ ; W^+

$$e^{V^{+}}(z;;) = \frac{1}{(1+1)^{'b}} \frac{z}{1+\frac{z}{2}}; \frac{z}{1+\frac{z}{2}};;$$

$$e^{W^{+}}(z;;) = \frac{1}{(1+1)^{'b}} \frac{z}{1+\frac{z}{2}}; \frac{z}{1+\frac{z}{2}};$$

so that the requirem ent of superconform alinvariance results in the system of equations:

$$V : R (Z_{1} :: Z_{N}) = R \quad Z_{1} + \frac{1}{2}; \; _{1} + \; ; \; _{1} :: Z_{N} + \frac{N}{2}; \; _{N} + \; ; \; _{N}$$

$$W : R (Z_{1} :: Z_{N}) = R \quad Z_{1} + \frac{1}{2}; \; _{1}; \; _{1} + \; :: Z_{N} + \frac{N}{2}; \; _{N}; \; _{N} + \frac{N}{2}; \; _{N}; \; _{N}; \; _{N} + \frac{N}{2}; \; _{N}; \; _{N}; \; _{N} + \frac{N}{2}; \; _{N}; \; _{N};$$

Solutions of V - and W -equations are shift-invariant functions which depend on the superdi erences only

$$R (Z) = R (Z_{ij}; i_{j}; i_{j});$$

$$Z_{ij} \quad z_{i} \quad z_{j} + \frac{1}{2} (i_{j} + i_{j}); i_{j} \quad i_{j}; i_{j} \quad i_{j}; i_{j} \quad i_{j};$$

Let us x som e point k. There are N 1 independent even di erences Z_{ik} ; $i \notin k$ for x ed k and 2 (N 1) independent odd di erences i_k ; i_k ; $i \notin k$ that transform with respect of the rem aining transform ations as follows

$$V^+: Z_{ij}! \frac{Z_{ij}}{(1+\frac{-i}{2})(1+\frac{-j}{2})}; ij! \frac{ij}{(1+\frac{-i}{2})(1+\frac{-j}{2})}; ij! ij$$

W + :
$$Z_{ij}$$
! $\frac{Z_{ij}}{(1+\frac{-i}{2})(1+\frac{-j}{2})}$; ij ! $\frac{ij}{(1+\frac{-i}{2})(1+\frac{-j}{2})}$; ij ! ij :

It is useful to extract the representation dependent part from the function R (2)

$$R (\mathcal{Z}) = \sum_{i < j}^{Y} Z_{ij}^{a_{ij}} 1 + \frac{ij \ ij}{Z_{ij}} R (\mathcal{Z}):$$

U sing the form ulae

$$V^{+}: \frac{ij}{Z_{ij}}! \frac{ij}{Z_{ij}}$$
 ; $1 + \frac{ij \ ij}{Z_{ij}}! 1 + \frac{ij \ ij}{Z_{ij}}$ (1 ij

$$W^{+}: \frac{ij}{Z_{ij}}! \frac{ij}{Z_{ij}}$$
 ; $1 + \frac{ij \ ij}{Z_{ij}}! 1 + \frac{ij \ ij}{Z_{ij}}$ (1 + ij)

it is easily to check that under the conditions

$$X$$
 $a_{ij}=$ $2\mathbf{i}_j$; $a_{ij}=$ a_{ji} ; $b_{ij}=$ b_j ; $b_{ij}=$ b_{ji} if j

the equations for the function R (2) reads as follows

$$V^{+}: R(Z_{1}::Z_{N}) = R \quad \frac{Z_{1}}{1 + \frac{1}{2}}; \frac{1}{1 + \frac{1}{2}}; 1:::\frac{Z_{N}}{1 + \frac{N}{2}}; \frac{N}{1 + \frac{N}{2}}; N \quad ;$$

$$V^{+}: R(Z_{1}::Z_{N}) = R \quad \frac{Z_{1}}{1 + \frac{1}{2}}; 1; \frac{1}{1 + \frac{N}{2}}:::\frac{Z_{N}}{1 + \frac{N}{2}}; N; \frac{N}{1 + \frac{N}{2}}; N \quad ;$$

i.e. the function R (Z) is the general function of all possible superconform all invariants. Let us go to the construction of the superconform all invariants. For this purpose we x the two points $Z_i; Z_k$ and choose the new variables (j and k are xed)

$$Z_{ij}^{k} = \frac{Z_{ji}}{Z_{ik}Z_{jk}}$$
; $Z_{ik}^{k} = \frac{ik}{Z_{ik}}$ Z_{ik}^{jk} Z_{ik}^{k} Z_{ik}^{k} Z_{ik}^{jk}

with the simple transform ation properties

It is easy to construct the superconform al invariants using these variables. Let us x the three points Z_j ; Z_k ; Z_p . Then the set of invariants is the following:

$$\frac{Z_{ij}^{k}}{Z_{pj}^{k}} = \frac{Z_{ij}Z_{pk}}{Z_{ik}Z_{pj}}; \frac{Z_{ij}^{k}}{Z_{pj}^{k}}$$

Finally we obtain the general solution of conform alconstraints

$$R(\mathbf{z}) = \begin{cases} Y \\ z_{ij}^{a_{ij}} \end{cases} 1 + \frac{ij \ ij}{Z_{ij}} \qquad R \qquad \frac{Z_{ij}Z_{pk}}{Z_{ik}Z_{pj}}; \frac{k_{ij} \ nm}{Z_{pj}^{k}} \end{cases}$$

$$X$$

$$a_{ij} = 2 \cdot_{j}; a_{ij} = a_{ji}; \qquad b_{ij} = b_{j}; b_{ij} = b_{ji}$$

$$i \in_{j} \qquad i \in_{j}$$

$$(7.0.18)$$

where j;k;p are xed. In the case of four-point function we can choose the superconform al invariant ansatz in the form

$$R (Z) = Z_{12}^{a} Z_{14}^{b} Z_{32}^{c} Z_{34}^{d} \qquad 1 + \frac{12 \cdot 12}{Z_{12}} \qquad 1 + \frac{14 \cdot 14}{Z_{14}} \qquad 1 + \frac{32 \cdot 32}{Z_{32}} \qquad 1 + \frac{34 \cdot 34}{Z_{34}}$$

$$R \frac{Z_{12} Z_{34}}{Z_{13} Z_{24}}; \frac{\frac{3}{12} \cdot \frac{3}{12}}{\frac{12}{12}}; \frac{\frac{3}{14} \cdot \frac{3}{14}}{Z_{\frac{3}{14}}}; \frac{\frac{3}{14} \cdot \frac{3}{12}}{Z_{\frac{3}{14}}}; \frac{\frac{3}{14} \cdot \frac{3}{12}}{Z_{\frac{3}{12}}}$$

w here

a
$$u '_1 '_2; b u '_1 + '_2; c u + '_1 '_2; d u + '_1 + '_2$$

= $b_1 ; = b_2 ; = b_2 b_1 + :$

Appendix B

In this Appendix we collect all needed integration formulae. General formulae of integration

Form ulae of integration for even vectors

$$Z_{z_{3}} dZ_{4}Z_{34}^{d}Z_{41}^{b} 1 + \frac{14 \cdot 14}{Z_{14}} 1 + \frac{34 \cdot 34}{Z_{34}} 1 + B \frac{\frac{3}{14} \cdot \frac{3}{14}}{Z_{14}^{3}} Z_{34}^{n} 1 + J \frac{\frac{34 \cdot 34}{Z_{34}}}{Z_{34}} = \frac{(b) \cdot (d+n)}{(b+d+n)} Z_{31}^{b+d+n} C_{1} + C_{2} \frac{\frac{31 \cdot 31}{Z_{31}}}{Z_{31}}$$

$$C_{1} = B + \frac{(d+n) \cdot b(+J)}{b+d+n}; C_{2} = (+J) \cdot \frac{b(d+n)}{4} + B + J + \frac{(d+n) \cdot b(+J)}{b+d+n}$$

$$Z_{21}^{a} 1 + \frac{12 \ 12}{Z_{12}} = \frac{Z_{22}}{Z_{12}} dZ_{3}Z_{23}^{c} 1 + \frac{32 \ 32}{Z_{32}} = 1 + A \frac{\frac{31 \ 31}{2}}{Z_{14}^{3}} Z_{34}^{b+d+n} 1 + J \frac{31 \ 31}{Z_{31}} = \frac{(c) (b+d+n)}{(c+b+d+n)} Z_{21}^{a+b+c+d+n} C_{1} + C_{2} \frac{21 \ 21}{Z_{21}}$$

$$C_{1} = A + \frac{cJ (b+d+n)}{b+c+d+n};$$

$$C_{2} = J + \frac{c(b+d+n)}{4} \frac{cJ (b+d+n)}{b+c+d+n} + A J \frac{cJ (b+d+n)}{b+c+d+n}$$

$$Z_{21}^{a} 1 + \frac{12 \ 12}{Z_{12}} \frac{Z_{22}}{Z_{12}} dZ_{3}Z_{23}^{c} 1 + \frac{32 \ 32}{Z_{32}}$$

$$Z_{23}^{a} dZ_{4}Z_{34}^{d}Z_{41}^{b} 1 + \frac{14 \ 14}{Z_{14}} 1 + \frac{34 \ 34}{Z_{34}} \frac{\frac{3}{14} \ \frac{32}{12}}{\frac{32}{21}}Z_{34}^{n} 1 + J \frac{34 \ 34}{Z_{34}} = \frac{(b) (d+n)}{(c+b+d+n)} (c) (b+d+n) Z_{21}^{a+b+c+d+n} C_{1} 1 + C_{2} \frac{21 \ 21}{Z_{21}}$$

$$C_{1} = C_{1} + C_{2} + C_$$

Form ulae of integration for odd vectors

References.

- [1] N M Bogoliubov, A G Jzergin, V E K orepin Quantum inverse scattering method, correlation functions and algebraic Bethe ansatz,

 C am bridge Univ P ress 1992
- [2] Z M aassarani, J Phys A: 28 (1995) 1305
- [3] P.B.Ramos, M. J.Martins, Nucl. Phys. B 474, (1996) 678
- [4] F.E. ler, V. Korepin, Phys. Rev. B 46, (1992) 9147
- [5] A Foerster, M K arowski, NuclPhys. B 396, (1993) 611
- [6] M P.P. fannmuller and H. Frahm , Nucl. Phys. B 479, (1996) 575
- [7] H Frahm, M P P fannmuller and A M . T svelik, Phys. Rev. Lett. 81, (1998) 2116
- [8] A Foerster, M K arowski, NuclPhys. B 408, (1993) 512, A Foerster, JPhys. A 29 (1996) 7625
 - D A maudon JHEP 12 (1997) 006, J Links, A Foerster, J Phys. A 32 (1999) 147 A Foerster, J Links, A P. Tonel, Nucl Phys. B 552, (1999) 707
 - JAmbjorn, DK arakhanyan, MM inum yan, A. Sedrakyan Ferm ionisation of the Spin-SU im in-Lai-Sutherland Model: Generalisation of Supersymmetric t-JM odel to Spin-Scond-mat/9909432, NuclPhys.BFS](2001) in press
- [9] S D erkachov, D K arakhanyan, R K irschner Heisenberg spin chains based on s'(2jl) symmetry, NuclPhys.B 583, (2000) 691
- [10] E.K. Sklyanin Classical limits of the Yang-Baxter equation, J.Soviet M ath. 40, (1988) 93
- [11] S.E. Derkachov, V.B. Kuznetsov, E.K. Sklyanin Baxter's Q-operator for the generic integrable XXX magnetic chain, in preparation
- [12] M. Scheunert, W. Nahm and V. Rittenberg, J.M. ath Phys. 18 (1977) 155
- [13] P.D. Jarvis, H.S.G. reen , J.M. ath Phys. 20 (1979) 2115
- [14] M M arcu, JM ath Phys21 (1980) 1277, JM ath Phys21 (1980) 1284
- [15] D A maudon, C Chryssom alakos, L Frappat, JM ath Phys. 36 (1995) 5262 L. Frappat, P. Sorba, A. Sciarrino, D ICTIONARY ON LIE SUPERALGEBRAS, hep-th/9607161
- [16] L.N. Lipatov, High-energy asymptotics of multicolor QCD and exactly solvable lattice models, Padova preprint DFPD-93-TH-70B, JETP Lett. 59 (1994)596.
- [17] LD Faddeev and GPK orchem sky, PhysLett.B 342 (1995)311.
- [18] G P K orchem sky, Nucl. Phys. B 443 (1995)255
- [19] V Braun, S Derkachov and A M anashov, PhysRev Lett. 81, (1998) 2020 V Braun, S Derkachov, G K orchem sky and A M anashov, NuclPhys. B 553, (1999) 355

- [20] S D erkachov, G K orchem sky and A M anashov, Evolution equations for quark-gluon distributions in multi-color Q C D and open spin chains, hep-ph/9909539, Nucl. Phys. B 566 (2000) 203.
- [21] A Belitsky, hep-ph/9907420, hep-ph/9903512, PhysLett. B 453, (1999) 59
- [22] D K arakhanian and R K irschner, Conserved currents of the three-reggeon interaction, hep-th/9902147; High-energy scattering in gauge theories and integrable spin chains, hep-th/9902031, Fortschr. Phys. 48, (2000) 139
- [23] H Frahm, D oped Heisenberg chains: spin S generalizations of the supersymmetric t-J model, cond-mat/9904157, NuclPhysB 559, (1999) 613
- [24] P. Kulish, N. Yu. Reshetikhin and E.K. Sklyanin, Lett. Math. Phys. 5 (1981) 393-403, E.K. Sklyanin, "Quantum Inverse Scattering Method. Selected Topics", in "Quantum Group and Quantum Integrable Systems" (Nankai Lectures in Mathematical Physics), ed. Mo-Lin Ge, Singapore World Scientic, 1992, pp. 63-97; hep-th/9211111
- [25] P.P.Kulish and E.K. Sklyanin, Zap Nauchn Sem. LOM I 95 (1980) 129
- [26] P.P.Kulish, Zap Nauchn Sem. LOM I145 (1985) 140, J.Soviet. Math. 35 (1986) 1111, Yang-Baxter equation and re-ection equations in integrable models, Schladming lecture 1995, hep-th/9507070
- [27] V O Tarasov, L A Takhta an and L D Faddeev Theor M ath Phys 57 (1983) 163-181,
- [28] L.D. Faddeev, Les-Houches lectures 1995, hep-th/9605187