

PERTURBATION THEORY FOR LYAPUNOV EXPONENTS OF A TORAL MAP :
EXTENSION OF A RESULT OF SHUB AND WILKINSON .

by David Ruelle* .

Abstract. Starting from a hyperbolic toral automorphism , we obtain, for a small volume preserving perturbation, an exact and rigorous second order perturbation expansion of the Lyapunov exponents.

Keywords: Lyapunov exponent, toral automorphism , hyperbolicity.

* Mathematics Dept., Rutgers University, and I.H.E.S. 91440 Bures sur Yvette, France.
<ruelle@ihes.fr>

We consider volume preserving perturbations F of a diffeomorphism $F_0 = (\cdot; J)$ of $T^{m+1} = T^m \times T$, where \cdot is a hyperbolic automorphism of T^m , and J is a translation of T . Writing $F = F_0 + aF^0$, we shall show that the Lyapunov exponents for $(F; \text{volume})$ can be expanded to second order in a (Theorem 1). In particular, the central Lyapunov exponent λ^c of $(F; \text{volume})$, to second order in a , is generally $\neq 0$ (Corollary 11). For a special family of perturbations one obtains particularly simple formulae, first noted by Shub and Wilkinson [10]. We recover their result in Theorem 12. We deviate from [10] mostly in that we don't have differentiability of λ^c , only a second order expansion around $a = 0$. The ideas used here are largely those in Shub and Wilkinson [10], and can be appreciated in the background provided by Hirsh, Pugh and Shub [6], Burns and Wilkinson [3], Ruelle and Wilkinson [9]. For recent work related to Lyapunov exponents, see also Bonatti, Gomez-Mont and Viana [2], Avila and Bochi [1].

After completing the writing of this paper, the author received a preprint by D. Dolgopyat [4], which develops similar ideas in a more general setting, but without the specific formulae we obtain here.

1. Theorem .

Let \cdot be a hyperbolic automorphism of T^m , and $J : y \mapsto y + (\text{mod } 1)$ a translation of T . Define $F_0 = (\cdot; J)$, and let $F = F_0 + aF^0$ be a C^2 perturbation of F_0 , volume preserving to first order in a . (We take $F^0 : T^{m+1} \rightarrow T^{m+1}$ and $F_0 + aF^0(\cdot)$ has to be understood (mod 1) in each component). Let $\lambda_1 < \lambda_2 < \dots$ be the Lyapunov exponents of $(F_0; \text{volume})$ and m_1, m_2, \dots their multiplicities (the exponent $= 0$ occurs with multiplicity 1). Also let $\lambda_a^{(1)}, \lambda_a^{(2)}, \dots$ be the Lyapunov exponents of $(F; \text{volume})$ repeated according to multiplicity. Then we have the second order expansion

$$\lambda_a^{(i)} = m_{r-1} + a^2 L_r + o(a^2) \quad \text{for } i = m_1 + \dots + m_{r-1} + 1$$

If $m_r = 1$, and writing $\lambda_r = \lambda_a^{(0)}$, we have

$$\lambda_a^{(0)} = \lambda_0^{(0)} + a^2 L^{(0)} + o(a^2)$$

(this applies in particular to $\lambda^c = \lambda_a^{(0)}$ for $\lambda_0^{(0)} = 0$).

An explicit expression for L_r can be obtained (see Proposition 9). We do not assume ergodicity of $(F; \text{volume})$, and therefore we use integrated Lyapunov exponents (averaged over the volume), see however Remark 15 (a).

If one likes one may to $F_0 + aF^0$ add terms of higher order in a so that the sum F is exactly volume preserving. These higher order terms will not change our results, and are omitted in what follows.

2. Normal hyperbolicity.

As in [10], we invoke the theory of normal hyperbolicity of [6]. We start from the fact that F_0 is normally hyperbolic to the smooth foliation of T^{m+1} by circles $\text{fix } T$.

Taking some $k \geq 2$ we apply [6] Theorems (7.1), (7.2). Thus we obtain a C^1 neighborhood U of F_0 in the C^k diffeomorphisms of T^{m+1} such that, for $F \in U$, there is an equivariant fibration $\pi: T^{m+1} \rightarrow T^m$ with

$$F^* =$$

The fibers $\pi^{-1}(x)$ are C^k circles forming a continuous fibration of T^{m+1} (this fibration is in general not smooth). Furthermore there is a TF -invariant continuous splitting of TT^{m+1} into three subbundles:

$$TT^{m+1} = E^s + E^u + E^c$$

such that E^c is 1-dimensional tangent to the circles $\pi^{-1}(x)$, E^s is m^s -dimensional contracting and E^u is m^u -dimensional expanding for TF .

If $\mu_r < 0$ (and F is in a suitable C^1 -small neighborhood U of F_0), we can introduce a continuous vector subbundle E^r of TT^{m+1} which consists of vectors contracting under TF^n faster than $(\mu_r + \epsilon)^n$ where $\epsilon > 0$ and $\mu_r + \epsilon < \mu_{r+1}$. In fact E^r is a hyperbolic (attracting) fixed point for the action induced by TF^{-1} on the bundle of $m_1 + \dots + m_r$ dimensional linear subspaces of TT^{m+1} (over F^{-1} acting on T^{m+1}).

If $\mu_r > 0$, replacement of F by F^{-1} similarly yields a continuous subbundle E^r of $m_r + \dots$ dimensional subspaces.

3. Proposition.

Assume that F is of class C^k , $k \geq 2$, and that F is C^k close to F_0 . The bundles E^r , E^r , when restricted to a circle $\pi^{-1}(x)$ are of class C^{k-1} , continuously in x .

If G denotes the (Grassmannian) manifold of $m_1 + \dots + m_r$ dimensional linear subspaces of \mathbb{R}^{m+1} , we may identify the bundle of $m_1 + \dots + m_r$ dimensional linear subspaces of TT^{m+1} with $T^{m+1} \times G$. We denote by $E \subset G$ the spectral subspace of the matrix defining π corresponding to the smallest $m_1 + \dots + m_r$ eigenvalues (in absolute value, and repeated according to multiplicity).

If F_0 is the action defined by TF_0 on $TT^{m+1} \times G$, the circles $\pi^{-1}(x) \times \pi^{-1}(E)$ form an F_0 invariant fibration of $T^{m+1} \times E$, to which F_0 is normally hyperbolic. If F is C^k close to F_0 , the corresponding C^{k-1} action F is normally hyperbolic to a perturbed fibration where $\pi^{-1}(x) \times \pi^{-1}(E)$ is replaced by $E^r \times \pi^{-1}(x)$. According to [6] Theorem 7.4, Corollary (8.3) and the following Remark 2, we find that the C^{k-1} circle $E^r \times \pi^{-1}(x) \subset T^{m+1} \times G$ depends continuously on $x \in T^{m+1}$. Similarly for E^- . \square

Note that in [10], the C^r section theorem is used in a similar situation, giving estimates uniform in x . However, continuity in x (not just uniformity) will be essential for us in what follows.

4. Corollary.

The splitting $TT^{m+1} = E^s + E^u + E^c$ when restricted to a circle $\pi^{-1}(x)$ is of class C^{k-1} , continuously in x .

It is clear that $E^{c,j-1}fxg$ is of class C^{k-1} because it is the tangent bundle to the C^k circle ^{j-1}fxg . As to E^s, E^u , they are special cases of E^r, E^r . \square

Notation.

Remember that $F = F_0 + aF^0$, and $x \in F^0$. We shall use the notation E_a, E_a^r, \dots to indicate the a -dependence of E, E^r, \dots

5. Proposition.

For small $a > 0$ there is a continuous function $x \mapsto x$ from T^m to $C^k(T^m(\cdot; a) \rightarrow T^m)$ such that $x(y; 0) = 0$ and $^{j-1}fxg = f(x + x(y; a); y) : y \in Tg$.

To see this define $\tilde{F} : T^{m+1}(\cdot; a) \rightarrow T^{m+1}(\cdot; a)$ by $\tilde{F}(\cdot; a) = (F_0 + aF^0)(\cdot; a)$ and observe that \tilde{F} is normally hyperbolic to the 2-dimensional manifolds

$$[a_2(\cdot; a)](^{j-1}fxg; a)$$

and these are thus C^k 2-dimensional submanifolds of $T^{m+1}(\cdot; a)$. \square

We may in the same manner replace ^{j-1}fxg by $[a_2(\cdot; a)](^{j-1}fxg; a)$ in Proposition 3 and Corollary 4. Writing E_a for $E_a^r, E_a^r, E_a^s, E_a^u, E_a^c$, we obtain that $(\cdot; a) \mapsto E_a(\cdot)$, when restricted from $T^{m+1}(\cdot; a)$ to $[a_2(\cdot; a)](^{j-1}fxg; a)$ is of class C^{k-1} . We rephrase this as follows:

6. Proposition.

The map

$$x \mapsto f(y; a) \mapsto E_a(x + x(y; a); y)g$$

where E_a stands for $E_a^r, E_a^r, E_a^s, E_a^u, E_a^c$, is continuous $T^m \rightarrow C^{k-1}(T^m(\cdot; a) \rightarrow T^m)$ (Grassmannian of R^{m+1}) where we have used the identification $TT^{m+1} = T^{m+1} \rightarrow R^{m+1}$. \square

Notation.

From now on we write E_a for $E_a^r, E_a^r, E_a^s, E_a^u, E_a^c$. When $a = 0$, E_0 is a trivial subbundle of $TT^{m+1} = T^{m+1} \rightarrow R^{m+1}$, and we shall write $E_0 = T^{m+1} \rightarrow E$, denoting thus by E a spectral subspace of the matrix on R^{m+1} defining $(\cdot; 1)$. We denote by E^\perp the complementary spectral subspace.

Taking $k = 2$ we have then:

7. Corollary.

There are linear maps $G(x; y); R(x; y; a) : E \rightarrow E^\perp$ such that $G(x; y)$ depends continuously on $(x; y) \in T^m \rightarrow T, R(x; y; a)$ on $(x; y; a) \in T^m \rightarrow T(\cdot; a)$,

$$E_a(x + x(y; a); y) = fX + aG(x; y)X + R(x; y; a)X : X \in E_g$$

and $\|R(x; y; a)\|$ is $o(a)$ uniformly in $x; y$. \square

Notice now that, if $x = x_a(x; y)$, then $x = x + x(y; a)$, where $x(y; a) = O(a)$. Now

$$E_a(x; y) = E_a(x + x(y; a); y) = fX + aG(x; y)X + R(x; y; a)X : X \in E_g$$

differs from

$$E_a(x + x(y; a); y) = fX + aG(x; y)X + R(x; y; a)X : X \in E_g$$

by the replacement $x \rightarrow x$ in the right-hand side, and since $\text{dist}(x; x) = O(a)$, we find that $\text{dist}(E_a(x; y); E_a(x + x(y; a); y)) = o(a)$. Therefore, changing the definition of R , we can again write:

8. Corollary.

There are linear maps $G(x; y); R(x; y; a) : E \rightarrow E^2$, depending continuously on their arguments, such that

$$E_a(x; y) = fX + aG(x; y)X + R(x; y; a)X : X \in E_g$$

and $\|R(x; y; a)\|$ is $o(a)$ uniformly in $x; y$. \square

We may write $TF = T(F_0 + aF^0) = D_0 + aD^0(\cdot)$ where D_0 does not depend on a and preserves the decomposition $T M = E + E^2$. If we apply TF to an element $X + aGX + RX$ of E_a (as in Corollary 8) we obtain $X_1 +$ element of $E^2 \subset E_a$, with $X_1 \in E$:

$$X_1 = D_0X + aD^0X + a^2D^0GX + aD^0RX \quad \text{projected on } E \quad (1)$$

Under $(TF)^\wedge$, the volume element in $E_a(\cdot)$ is multiplied by a factor $M(\cdot; a)$, and the projection in E of $(TF)^\wedge$ is equal to the projection in E of \cdot multiplied by a factor $N(\cdot; a)$ such that

$$M(\cdot; a) = N(\cdot; a) + \frac{1}{a}(\cdot) - \frac{1}{a}(F(\cdot))$$

for suitable $\frac{1}{a}$. We may compute N from (1):

$$N(\cdot; a) = N_{(0)} + aN_{(1)}(\cdot) + a^2N_{(2)}(\cdot) + o(a^2)$$

To proceed we take now $E_a = E_a^r$, and assume $e_r < 0$. We have then, writing d for the volume element in T^{m+1} ,

$$\begin{aligned} L_a &= \int_{x=1}^{m_1+x \dots + m_r} \int_a^{(')} = \int_a^Z d \log M(\cdot; a) = \int_a^Z d \log N(\cdot; a) \\ &= L_{(0)} + aL_{(1)}(\cdot) + a^2L_{(2)}(\cdot) + o(a^2) \end{aligned} \quad (2)$$

More precisely, we shall prove

9. Proposition.

If $r < 0$, we have

$$\sum_{k=1}^{m_1+\dots+m_r} \frac{X^k}{a^k} = \sum_{k=1}^r m_k \lambda_k + a^2 L + o(a^2)$$

where

$$L = \frac{1}{2} \sum_{n=1}^{\infty} \text{Tr} (D_0^{-1} D^0 (F_0^n)) \text{Tr} (D_0^{-1} D^0 (F_0^n)) = 0$$

and E is the spectral subspace of the matrix defining corresponding to the smallest $m_1 + \dots + m_r$ eigenvalues (in absolute value, and repeated according to multiplicity).

The proof that $L = 0$ is postponed to Remark 15(b).

The proposition is obtained by comparing formula (2) with the formula (5) below, which we shall obtain by a second order perturbation calculation.

To first order in a we have

$$F^n = (F_0 + aF_0^n)^n = F_0^n + a \sum_{j=1}^n F_0^{n-j} F_0^j$$

hence

$$TF^n = D_0^n + a \sum_{j=1}^n D_0^{n-j} D^0(F^{j-1}) D_0^{j-1}$$

If we apply TF^n to $X + aGX + RX \in E_a$ we obtain $X_n \in E_a$, with $X_n \in E$. To zero-th order in a , $X_n = D_0^n X$, so we may write to first order $X_n = D_0^n X + aY_n(\cdot)$. Therefore, to first order in a ,

$$D_0^n X + aY_n(\cdot) + aG(F^n)D_0^n X = D_0^n X + a \sum_{j=1}^n D_0^{n-j} D^0(F^{j-1}) D_0^{j-1} X + aD_0^n G(\cdot)X$$

and, taking the components along E^\perp ,

$$G(F^n)D_0^n X = \sum_{j=1}^n D_0^{n-j} D_?^0(F^{j-1}) D_0^{j-1} X + D_0^n G(\cdot)X$$

where $D_?^0(\cdot)$ is $D^0(\cdot)$ followed by taking the component along E^\perp , or

$$\sum_{j=1}^n D_0^{n-j} D_?^0(F^{j-1}) D_0^{j-1} X + G(\cdot)X = D_0^n G(F^n)D_0^n X$$

When $n \rightarrow 1$, the right-hand side tends to zero (exponentially fast, remember that $X \in E$, $G \in E^2$). Therefore

$$G(\alpha)X = \sum_{j=1}^{\infty} D_0^{-j} D_0^0 (F^{j-1}) D_0^{j-1} X$$

which we shall use in the form

$$G(\alpha)X = \sum_{n=0}^{\infty} D_0^{n-1} D_0^0 (F_0^n) D_0^n X \quad (3)$$

where we have written F_0^n instead of F^n since G is evaluated to order 0 in α . (The right-hand side is an exponentially convergent series).

Returning to (1) we see that, to second order in α ,

$$\begin{aligned} X_1 &= D_0 X + \alpha D_0^0(\alpha)X + \alpha^2 D_0^0(\alpha)G(\alpha)X \quad \text{projected on } E \\ &= D_0 (1 + \alpha D_0^{-1} D_0^0(\alpha) + \alpha^2 D_0^{-1} D_0^0(\alpha)G(\alpha))X \quad \text{projected on } E \end{aligned}$$

Let now $(u^{(i)})$ and $(u^{(i)\dagger})$ be conjugate bases of E . Also let λ_i for $i = 1, \dots, m_1 + \dots + m_r$ be the eigenvalues of D_0 restricted to E . Then, to second order in α ,

$$N(\alpha; a) \sim \prod_{i=1}^{m_1 + \dots + m_r} u^{(i)}$$

is, up to a factor of absolute value 1,

$$\begin{aligned} & \left(\prod_{i=1}^{m_1 + \dots + m_r} u^{(i)} \right) \left[1 + \alpha \sum_{i=1}^{m_1 + \dots + m_r} (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) \right. \\ & \quad + \alpha^2 \sum_{i < j}^X ((u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) (u^{(j)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(j)})) \\ & \quad \left. (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(j)}) (u^{(j)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) + \alpha^2 \sum_i^X (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) G(\alpha) u^{(i)}) \right] \sim u^{(\alpha)} \end{aligned}$$

so that

$$\begin{aligned} N(\alpha; a) &= \left(\prod_{i=1}^{m_1 + \dots + m_r} u^{(i)} \right) \left[1 + \alpha \sum_i^X (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) \right. \\ & \quad + \alpha^2 \sum_{i < j}^X ((u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) (u^{(j)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(j)})) \\ & \quad \left. (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(j)}) (u^{(j)\dagger}; D_0^{-1} D_0^0(\alpha) u^{(i)}) + \alpha^2 \sum_i^X (u^{(i)\dagger}; D_0^{-1} D_0^0(\alpha) G(\alpha) u^{(i)}) g \right] \end{aligned}$$

Since $\log j^{(i)} = \log j_0^{(i)}$ we obtain, to second order in a ,

$$L_a = \sum_i \log N(\cdot; a) = m_{1-1} + \dots + m_{r-r} + \sum_i \int \frac{a^2}{2} \left(\sum_i (u^{(i)})^2; D_0^{-1} D^0(\cdot) u^{(i)} \right)^2 \Big]$$

where \int has the same meaning as above. Write

$$\sum_i (u^{(i)})^2; D_0^{-1} F^0(\cdot) u^{(i)})$$

The first term of \int is

$$a^2 \sum_i \int (u^{(i)})^2; D_0^{-1} T F^0(\cdot) u^{(i)}) = a^2 \sum_i \int \frac{\partial}{\partial x_i}$$

which vanishes because $\int \frac{\partial}{\partial x_i} = 0$. The next term in \int is

$$a^2 \sum_{i < j} \int \left(\frac{\partial}{\partial x_i} \right) \left(\frac{\partial}{\partial x_j} \right) \left(\frac{\partial}{\partial x_j} \right) \left(\frac{\partial}{\partial x_i} \right) = a^2 \sum_{i < j} \int \left(\frac{\partial}{\partial x_i} \left(\frac{\partial}{\partial x_j} \right) \right) \left(\frac{\partial}{\partial x_j} \left(\frac{\partial}{\partial x_i} \right) \right)$$

which vanishes as above. Thus we are left with

$$L_a = (m_{1-1} + \dots + m_{r-r}) = a^2 \sum_i \int \left((u^{(i)})^2; D_0^{-1} D^0(\cdot) G(\cdot) u^{(i)} \right) - \frac{1}{2} \sum_i \int (u^{(i)})^2; D_0^{-1} D^0(\cdot) u^{(i)})^2 \Big] \quad (4)$$

and we may write, using (3),

$$\begin{aligned} \sum_i (u^{(i)})^2; D_0^{-1} D^0(\cdot) G(\cdot) u^{(i)} &= \sum_{n=0}^{\infty} \sum_i (u^{(i)})^2; D_0^{-1} D^0(\cdot) D_0^{n-1} D^0(F_0^n) D_0^n u^{(i)} \\ &= \sum_{n=0}^{\infty} \sum_i \sum_j (u^{(i)})^2; D_0^{-1} D^0(\cdot) u^{(j)}) (u^{(j)})^2; D_0^{n-1} D^0(F_0^n) D_0^n u^{(i)} \end{aligned}$$

where we have introduced conjugate bases $(u^{(j)})$, $(u^{(j)})^*$ of E , indexed by $j = m_1 + \dots + m_r + 1; \dots; m_1 + \dots + m_r + 1$, and i is over $i = m_1 + \dots + m_r + 1$, j is over $j > m_1 + \dots + m_r + 1$. The above expression is also

$$= \sum_{n=0}^{\infty} \sum_i \sum_j \frac{\partial}{\partial x_j} (u^{(i)})^2; D_0^{-1} F^0(\cdot) u^{(i)}) \frac{\partial}{\partial x_i} (u^{(j)})^2; D_0^{n-1} F^0(F_0^n) u^{(i)})$$

and integration by part gives thus

$$\sum_i \int (u^{(i)})^2; D_0^{-1} D^0(\cdot) G(\cdot) u^{(i)})$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} d \sum_i \frac{\partial}{\partial_i} (u^{(i)}; D_0^{-1} F^0(\cdot, u^{(i)})) \sum_j \frac{\partial}{\partial_j} (u^{(j)}; D_0^{n-1} F^0(F_0^n \cdot, u^{(j)})) \\
&= \sum_{n=0}^{\infty} d \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{E}^?}(D_0^{n-1} D^0(F_0^n) D_0^n) \\
&= \sum_{n=0}^{\infty} d \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{E}^?}(D_0^{-1} D^0(F_0^n))
\end{aligned}$$

The fact that $F = F_0 + aF^0$ is volume preserving is expressed by $\operatorname{Tr}_{\mathbb{R}^{m+1}}(D_0^{-1} D^0(\cdot)) = 0$ hence

$$\begin{aligned}
&d \sum_i (u^{(i)}; D_0^{-1} D^0(\cdot)) G(\cdot) u^{(i)} \\
&= \sum_{n=0}^{\infty} d \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(F_0^n))
\end{aligned}$$

and introducing this in (4) yields

$$\begin{aligned}
&L_a = (m_{1-1} + \dots + m_{r-r}) \\
&= a^2 \left[\sum_{n=1}^{\infty} d \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(F_0^n)) + \frac{1}{2} \sum_{n=1}^{\infty} d (\operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)))^2 \right] \\
&= \frac{a^2}{2} \sum_{n=1}^{\infty} d \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{E}}(D_0^{-1} D^0(F_0^n)) \tag{5}
\end{aligned}$$

where the last step used the invariance of d under F_0^n . \square

10. Proof of Theorem 1.

We use Proposition 9, the corresponding result with F replaced by F^{-1} , and the fact that $\sum_{i=1}^m a^{(i)} = 0$ (because F is volume preserving). This gives an estimate of all the sums of $a^{(i)}$ that occur in Theorem 1. \square

11. Corollary.

In the situation of Theorem 1, the central Lyapunov exponent is

$$\begin{aligned}
c &= \frac{a^2}{2} \sum_1^{\infty} d [\operatorname{Tr}^u(D_0^{-1} D^0(\cdot)) \operatorname{Tr}^u(D_0^{-1} D^0(F_0^n)) - \operatorname{Tr}^s(D_0^{-1} D^0(\cdot)) \operatorname{Tr}^s(D_0^{-1} D^0(F_0^n))] \\
&= \frac{a^2}{2} \sum_1^{\infty} d [\operatorname{Tr}^s(D_0^{-1} D^0(\cdot)) - \operatorname{Tr}^u(D_0^{-1} D^0(\cdot))] \operatorname{Tr}^c(D_0^{-1} D^0(F_0^n))
\end{aligned}$$

where $\operatorname{Tr}^s, \operatorname{Tr}^u, \operatorname{Tr}^c$ denote the traces over the spectral subspaces E^s, E^u, E^c of D_0 corresponding to eigenvalues $< 1, > 1$, or $= 1$ in absolute value (E^c is one dimensional).

Since F preserves the volume, the sum of all Lyapunov exponents vanishes. Therefore λ^c is minus the sum of the negative Lyapunov exponents, given by (5), minus the sum of the positive Lyapunov exponents. Note that replacing F by F^{-1} , E^s by E^u (and, to the order considered, $D^0(\cdot)$ by $-D^0(\cdot)$) replaces the sum of the negative Lyapunov exponents by minus the sum of the positive exponents. This gives the first formula for λ^c .

To obtain the second formula, express $\text{Tr}^u \text{Tr}^u - \text{Tr}^s \text{Tr}^s$ in terms of $\text{Tr}^u - \text{Tr}^s$, and remember that (because F preserves the volume) $\text{Tr}^s + \text{Tr}^u + \text{Tr}^c = 0$ when applied to $D_0^{-1} D^0(\cdot)$. \square

The above formula (5) takes a particularly simple form in a special case described in the next theorem.

12. Theorem.

Let h be a hyperbolic automorphism of T^m , with stable and unstable dimensions m^s and $m^u = m - m^s$, and with entropy h_0^u . Let $J : y \mapsto y + a(y) \pmod{1}$ be a translation of T , and $\gamma : T^m \rightarrow T$ a morphism $\neq 0$. Finally let $\phi : T \rightarrow \mathbb{R}^m$ be a nullhomotopic C^2 function.

Define $h_a, g_a : T^m \rightarrow T^m$ by

$$h_a \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ Jy + \gamma(x) \end{pmatrix}; \quad g_a \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x + a(y) \\ y \end{pmatrix} \pmod{1}$$

and let $f_a = g_a \circ h$.

Denote by λ_a^s (resp. λ_a^u) the sum of the smallest m^s (resp. the largest m^u) Lyapunov exponents for $(f_a; \text{volume})$. Also let $\lambda_a^c = \lambda_a^s + \lambda_a^u$ be the "central exponent". Then $\lambda_a^s, \lambda_a^u, \lambda_a^c$ have expansions of order 2 in a :

$$\begin{aligned} \lambda_a^s &= \lambda_0^u + \frac{a^2}{2} \int_T dy ((r^-)^{0s}(y))^2 + o(a^2) \\ \lambda_a^u &= \lambda_0^u - \frac{a^2}{2} \int_T dy ((r^-)^{0u}(y))^2 + o(a^2) \\ \lambda_a^c &= \frac{a^2}{2} \int_T dy [((r^-)^{0u}(y))^2 - ((r^-)^{0s}(y))^2] + o(a^2) \end{aligned}$$

Here $(r^-)^{0s}(y)$ and $(r^-)^{0u}(y)$ are the components of the derivative $D\phi(y) \in \mathbb{R}^m$ in the stable and unstable subspaces E^s and E^u for h . Also, we have used $r^- : \mathbb{R}^m \rightarrow \mathbb{R}$ to denote the derivative of the map $\gamma : T^m \rightarrow T$ with the obvious identifications.

This theorem is a simple (but nontrivial) extension of the result proved by Shub and Wilkinson [10]. In the situation that they consider $\gamma = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$, $J = \text{identity}$, $\phi = (1; 0)$, $\lambda_0^u = \lambda_0^u$. [Remark that, in the notation of [10], $u_0 = ((1; 1) : v_0) = (m - 1) = ((1; 0) : v_0)$ so that the formula given in Proposition II of [10] agrees with our result above].

Notation.

We shall henceforth omit the (mod 1). We shall keep r to denote the derivative in T^m . With obvious abuses of notation, the reader may find it convenient to think of r as an $m \times m$ matrix (with integer entries and determinant 1), and r as a row m -vector (with integer entries not all zero).

13. Reformulation of the problem.

Note that $f_a^{-1} = h^{-1} g_a^{-1}$ where h^{-1}, g_a^{-1} are obtained from h, g_a by the replacements $J; ; \rightarrow -1; J^{-1}; ;$. These replacements also interchange the stable and unstable subspaces for r and replace s, u by u, s . Therefore the formula for u in the theorem follows from the formula for s . And the formula for $c = s - u$ also follows. To complete the proof of the theorem we turn now to the formula for s .

Define

$$\hat{x} = y + x$$

then

$$F_0 \frac{x}{y} = h^{-1} \frac{\hat{x}}{y} = \frac{x}{Jy}$$

$$g_a \frac{x}{y} = g_a^{-1} \frac{\hat{x}}{y} = \frac{x + a(y + x)}{y - a(r)(y + x)}$$

so that

$$F \frac{x}{y} = h^{-1} f_a \frac{\hat{x}}{y} = g_a F_0 \frac{x}{y} = \frac{x + a(Jy + x)}{Jy - a(r)(Jy + x)}$$

Finally, $F = F_0 + aF^0$ with

$$F_0 \frac{x}{y} = \frac{x}{Jy} ; \quad F^0 \frac{x}{y} = \frac{(Jy + x)}{(r)(Jy + x)}$$

Since F is conjugate (linearly) to f_a , we may compute s from F instead of f_a .

14. Proof of Theorem 12.

Write $R^{m+1} = E^s + E^u + R$. We shall apply Proposition 9 with $E = E^s, E^? = E^u + R$. Using $\cdot = (x; y)$ and $X \in E^s, Y \in E^u, Z \in R$ we may write

$$D_0 \frac{X + Y}{Z} = \frac{(r)(X + Y)}{Z}$$

$$D^0(\cdot) \frac{X + Y}{Z} = \frac{{}^0(Jy + x)((r)(X + Y) + Z)}{(r)({}^0(Jy + x)((r)(X + Y) + Z))}$$

where 0 denotes the derivative of \cdot . Therefore

$$\text{Tr}_E(D^0(\cdot)D_0^{-1}) = (r) {}^0s(Jy + x)$$

and (5) contains the integrals

$$\begin{aligned} & \int d \operatorname{Tr} \left(D_0^{-1} D^0(\cdot) \right) \operatorname{Tr} \left(D_0^{-1} D^0(F_0^n) \right) \\ &= \int d \left[(r)^{-\alpha} (Jy + x) \right] \left[(r)^{-\alpha} (J^{n+1}y + x^{n+1}) \right] \end{aligned}$$

Performing a change of variables $x = x, y = Jy + x$ we find that this is

$$= \int dx dy \left[(r)^{-\alpha} (y) \right] \left[(r)^{-\alpha} (J^n y + x^n x) \right]$$

We claim that this last integral vanishes unless $n = 0$. This is because, if $n \neq 0$,

$$\int dx \left(J^n y + x^n x \right) = 0$$

Indeed, $x^n x$ is a linear combination with integer coefficients of the components x_1, \dots, x_m of x , and the coefficients do not all vanish because $x^n =$ is impossible (is hyperbolic and $\neq 0$). Integrating the derivative 0 with respect to a variable x_j really occurring in $x^n x$ gives zero as announced.

Returning to (5) we have thus

$$\begin{aligned} \frac{s}{a} + \frac{u}{o} &= \frac{a^2}{2} \int d \left(\operatorname{Tr} \left(D_0^{-1} D^0(\cdot) \right) \right)^2 \\ &= \frac{a^2}{2} \int dy \left((r)^{-\alpha} (y) \right)^2 \end{aligned}$$

which is the formula given for $\frac{s}{o}$ in Theorem 12. And according to Section 13 this completes our proof. \square

15. Final remarks.

(a) Shub and Wilkinson [10] showed that close to a diffeomorphism (hyperbolic automorphism of T^2) (identity on T) there is a C^1 open set of ergodic volume preserving C^2 diffeomorphisms of T^3 with central Lyapunov exponent $\lambda^c > 0$. They remark that their result extends to the situation where ϕ is a hyperbolic automorphism of T^m with one-dimensional expanding eigenspace. More generally, if ϕ is any hyperbolic automorphism of T^m , Theorem 12 gives close to $(\phi; \text{rotation of } T)$ in $C^2(T^{m+1})$ a diffeomorphism F with $\lambda^c > 0$. Since λ^c is given by an integral over the volume of a local "central" stretching exponent, we have $\lambda^c > 0$ in a C^1 neighborhood of F . But by a result of Dolgopyat and Wilkinson [5] (Corollary 0.5), stable ergodicity is here C^1 open and dense in the C^2 volume preserving diffeomorphisms: we have center bunching and stable dynamical coherence because we consider perturbations of $(\phi; \text{rotation of } T)$ for which the center foliation is C^1 , see [6], [7]. In conclusion, close to (hyperbolic automorphism of T^m) (rotation on T) there is a C^1 open set V of ergodic volume preserving C^2 diffeomorphisms of T^{m+1} with

central Lyapunov exponent $\lambda^c > 0$ (or also with $\lambda^c < 0$). In particular, if $F \in V$, the conditional measures of the volume on the circles $\pi^{-1}(x)$ are atomic, as discussed in [9].

(b) The coefficient L in Proposition 9 is 0. Consider indeed the unitary operator U defined by $U = F$ on $L^2(T^{m+1}; \text{volume})$, and let $E(\cdot)$ be the corresponding spectral measure, so that

$$U = \int_T e^{2\pi i \lambda} E(d\lambda)$$

If $\mu(\cdot) = \text{Tr} (D_0^{-1} D^0(\cdot))$ we have a measure μ on T defined by $\mu(d\lambda) = \mu(\cdot; E(d\lambda))$ and the Fourier coefficients

$$c_n = \int_T e^{2\pi i n \lambda} \mu(d\lambda) = \int_T d \text{Tr} (D_0^{-1} D^0(\cdot)) (D_0^{-1} D^0(F_0^n \cdot))$$

of this measure tend to zero exponentially. Therefore $\mu(d\lambda) = \rho(\lambda) d\lambda$ has a smooth density and

$$L = \frac{1}{2} \sum_{n=1}^{\infty} \lambda_n^2, \quad c_n = \frac{1}{2} \lambda_n^2 \quad (0) \quad 0$$

(c) Suppose now that F is not necessarily a volume preserving perturbation of F_0 . We may still hope that F has an SRB measure μ_a . If F_0 were hyperbolic, we would have an expansion

$$\mu_a = \mu_0 + a + o(a)$$

(see [8]) with $\mu_0 = \text{Lebesgue measure}$ and a a distribution. For smooth μ , $\mu(\cdot)$ is given (because μ_0 is Lebesgue measure) by the simple formula (see [8])

$$\mu(\cdot) = \int_0^1 \mu_0((\cdot - F_0^n) \text{div}(F^0 - F_0^{-1}))$$

Similarly (replacing F by F^{-1} , hence $F_0, D_0^{-1} D^0(\cdot)$ by $F_0^{-1}, D^0(F_0^{-1}) D_0^{-1}$) we see that the anti-SRB state has an expansion

$$\mu_a = \mu_0 + a + o(a)$$

with

$$\begin{aligned} \mu(\cdot) &= \sum_{n=1}^{\infty} \int_T d(F_0^{-n}) \text{Tr}_{\mathbb{R}^{m+1}} (D^0(F_0^{-1}) D_0^{-1}) \\ &= \sum_{n=0}^{\infty} \int_T d(F_0^{-n}) \text{Tr}_{\mathbb{R}^{m+1}} (D_0^{-1} D^0(\cdot)) \end{aligned}$$

We can now estimate the Lyapunov exponents for $(F; \mu_a)$ to second order in a even though we are not sure of the existence of the SRB measure μ_a . We simply assume that we can use the formula for $\mu(\cdot)$. Going through the proof of Proposition 9 we have to replace

R
 $d \log N(\cdot; a)$ by $a(\log N(\cdot; a))$ and (to second order in a) this adds to the right-hand side of (4) a term

$$a^2 \sum_{n=1}^Z d \operatorname{Tr}(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{K}_{m+1}}(\mathbb{D}_0^{-1} D^0(\cdot))$$

Taking into account the integrations by part we obtain now instead of (5) the formula

$$L_a(m_{1-1} + \dots + m_{r-r}) = \frac{a^2}{2} \sum_{n=1}^Z d \operatorname{Tr}(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$a^2 \sum_{n=1}^Z d \operatorname{Tr}(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{K}_{m+1}}(\mathbb{D}_0^{-1} D^0(F_0^n \cdot)) \quad (6)$$

Let $a^2 L^s, a^2 L^u, a^2 L^c$ be the a^2 contributions to the sum of the noncentral negative, noncentral positive, and the central Lyapunov exponents for the SRB measure. We obtain $a^2 L^s$ from (6) when $n_r = n^s$. A similar calculation gives $a^2 L^u$ (it is convenient here to work via the anti-SRB measure, then replace F by F^{-1}). Estimating the average expansion coefficient gives $a^2(L^s + L^u + L^c) = a(\log \det(\mathbb{D}_0 + a D^0(\cdot)))$, hence $L^s + L^u + L^c$, hence L^c . The results are

$$L^s = \frac{1}{2} \sum_{n=1}^Z d \operatorname{Tr}^s(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}^s(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$\sum_{n=1}^Z d \operatorname{Tr}^s(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{K}_{m+1}}(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$L^u = \frac{1}{2} \sum_{n=1}^Z d \operatorname{Tr}^u(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}^u(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$L^c = \frac{1}{2} \sum_{n=1}^Z d \operatorname{Tr}^c(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}^c(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$\sum_{n=1}^Z d \operatorname{Tr}^c(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}^u(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

$$L^s + L^u + L^c = \frac{1}{2} \sum_{n=1}^Z d \operatorname{Tr}_{\mathbb{K}_{m+1}}(\mathbb{D}_0^{-1} D^0(\cdot)) \operatorname{Tr}_{\mathbb{K}_{m+1}}(\mathbb{D}_0^{-1} D^0(F_0^n \cdot))$$

which can be rewritten variously.

The existence of second order expansions for the Lyapunov exponents gives added interest to the question whether the SRB measure a really exists for (small) finite a .

(d) The author has not looked seriously into possible extensions of the results presented here. Generalizations are thus left for the reader to formulate, and to prove.

References.

- [1] A. Avila and J. Bochi. \A formula with some applications to the theory of Lyapunov exponents." Preprint
- [2] C. Bonatti, X. Gomez-Mont and M. Viana. \G nericit  d'exposants de Lyapunov non-nuls des produits d'endomorphismes de matrices." Preprint
- [3] K. Burns and A. Wilkinson. \Stable ergodicity of skew products." Ann. Sci.  cole Norm. Sup. 32,859-889 (1999).
- [4] D. Dolgopyat. \On differentiability of SRB states." Preprint.
- [5] D. Dolgopyat and A. Wilkinson. \Stable accessibility is C^1 dense." Preprint
- [6] M. Hirsch, C.C. Pugh, and M. Shub. Invariant manifolds. Lect. Notes in Math. 583 Springer, Berlin, 1977.
- [7] C. Pugh, and M. Shub. \Stable ergodicity and julienne quasi-conformality." J. Eur. Math. Soc. 2,1-52 (2000).
- [8] D. Ruelle. \Differentiation of SRB states." Commun. Math. Phys. 187,227-241 (1997).
- [9] D. Ruelle and A. Wilkinson. \Absolutely singular dynamical foliations." Commun. Math. Phys., to appear.
- [10] M. Shub and A. Wilkinson. \Pathological foliations and removable exponents." Inventiones Math. 139,495-508 (2000).