

# On Relations of Hyperelliptic Weierstrass al Functions

SHIGEKI MATSUTANI

## Abstract

We study relations of the Weierstrass's hyperelliptic al-functions over a non-degenerated hyperelliptic curve  $y^2 = f(x)$  of arbitrary genus  $g$  as solutions of sine-Gordon equation using Weierstrass's local parameters, which are characterized by two ramified points. Though the hyperelliptic solutions of the sine-Gordon equation had already obtained, our derivations of them is simple; they need only residual computations over the curve and primitive matrix computations.

**2000 MSC:** 14H05, 14K20, 33E99, 30F30

**Key Words and Phrases:** Hyperelliptic Functions,  
sine-Gordon equation, Weierstrass al Functions

## §1. Introduction

The sine-Gordon equation is a famous nonlinear integrable differential equations. For a hyperelliptic curve  $C_g$  ( $y^2 = f(x) = (x - b_1) \cdots (x - b_{2g+1})$ ) of genus  $g$ , the hyperelliptic solutions of the sine-Gordon equation were formulated in [Mu 3.241] in terms of Riemann theta functions. In [Mu],  $(U, V, W)$  representation of symmetric product space of the  $g$  curves  $Symm^g C_g$  is defined; especially,  $U$  is defined by  $U(z) := (x_1 - z) \cdots (x_g - z)$  a for a point  $((x_1, y_1), \cdots, (x_g, y_g))$  in  $Symm^g C_g$ . (In this article, we will denote  $U$  by  $F(z)$  on later following the conventions in [Ba1, 2, 3, Ma].) Using the relation between  $U$  and the Riemann theta functions in [Mu 3.113], the solutions [Mu 3.241] can be rewritten as,

$$\frac{\partial}{\partial t_P} \frac{\partial}{\partial t_Q} \log([2P - 2Q]) = A([2P - 2Q] - [2Q - 2P]), \quad (1-3)$$

where  $P$  and  $Q$  are ramified points of  $C_g$ ,  $A$  is a constant number,  $[D]$  is a meromorphic function over  $Symm^g(C_g)$  with a divisor  $D$  for each  $C_g$  and  $t_{P'}$  is a coordinate in the Jacobi

variety such that it is identified with a local parameter at a branch point  $P'$  up to constant. In other words, for a finite branch point  $(b_i, 0)$   $U(b_i)$  is identified with  $[2(b_i, 0) - 2\infty]$  up to constant factor.

In the formulations in [Mu], local parameters  $t_{P'}$  were not concretely treated. In this article, we will give more explicit representations of (1-3) using concrete local parameters in [Ba2, W2, 3] and present simpler derivations of (1-3) without using any  $\theta$ -function. This article is an application of a scheme developed in [Ma] to the sine-Gordon equation, which is based upon [Ba3].

In [W1, W2], Weierstrass defined  $\text{al}$  function by  $\text{al}_r = \gamma_r \sqrt{U(b_r)}$  using a constant factor  $\gamma_r$ . In Theorem 3.1, we will give

$$\frac{\partial^2}{\partial u_1^{(r)} \partial u_g^{(r)}} \log \text{al}_r(u^{(r)}) = \frac{1}{2} \left( \frac{\text{al}_r^2(u^{(r)})}{\gamma_r^2} - \frac{f'(b_r) \gamma_r^2}{\text{al}_r^2(u^{(r)})} \right), \quad (1-4)$$

in terms of a coordinate system  $u^{(r)}$ 's defined in (2-5).  $\text{al}_r(u)$  has the single order zero at  $(b_r, 0)$  and a singularity of the single order at  $\infty$  as a function of  $x_i \in C_g$ .

Further we give another representation in Theorem 4.1 in terms of  $v$ 's defined in (2-6) ( $a_1 := b_r, a_2 := b_s$ ) [W2, 3],

$$\frac{\partial^2}{\partial v_1 \partial v_2} \log \frac{\text{al}_r(v)}{\text{al}_s(v)} = \frac{1}{2(b_r - b_s)} \left( f'(b_s) \frac{\gamma_s^2 \text{al}_r(v)^2}{\gamma_r^2 \text{al}_s(v)^2} + f'(b_r) \frac{\gamma_r^2 \text{al}_s(v)^2}{\gamma_s^2 \text{al}_r(v)^2} \right). \quad (1-5)$$

The function  $\text{al}_s(v)/\text{al}_r(v)$  vanishes with order one when  $x_i$  is at  $(b_s, 0)$  whereas it diverges with order one if  $x_i$  approaches to  $(b_r, 0)$ . As they were discovered by Weierstrass [W2, 3] and they play the essential roles in the investigation in [W2, 3] and in §4. Thus we have called them *Weierstrass parameters*.

In these proofs, we will use only residual computations using the data of curve itself without any  $\theta$  functions as the derivation of hypereilliptic solutions of the modified Korteweg-de Vries equations in [Ma]. The curve is sometimes given by an affine equation with special coefficients. Then it might be important to study the relation between the properties of line-bundle over the curve and these coefficients. As (1-4) and (1-5) can be explicitly expressed by data of curve  $C_g$ , the author believes that they have some advantage as relations of special functions.

## §2. Differentials of a Hyperelliptic Curve

In this section, we will give the conventions and notations of the hyperelliptic functions in this article. We denote the set of complex numbers by  $\mathbb{C}$  and the set of integers by  $\mathbb{Z}$ .

**2.1 Hyperelliptic Curve.** We deal with a hyperelliptic curve  $C_g$  of genus  $g$  ( $g > 0$ ) given by the affine equation,

$$\begin{aligned} y^2 &= f(x) \\ &= \lambda_{2g+1}x^{2g+1} + \lambda_{2g}x^{2g} + \cdots + \lambda_2x^2 + \lambda_1x + \lambda_0 \\ &= (x - b_r)h_r(x), \end{aligned} \tag{2-1}$$

where  $\lambda_{2g+1} \equiv 1$  and  $\lambda_j$ 's are complex numbers. We use the expressions,

$$\begin{aligned} f(x) &:= (x - b_1)(x - b_2) \cdots (x - b_{2g})(x - b_{2g+1}) \\ &= P(x)Q(x), \\ P(x) &:= (x - a_1)(x - a_2) \cdots (x - a_g), \\ Q(x) &:= (x - c_1)(x - c_2) \cdots (x - c_g)(x - c), \end{aligned} \tag{2-2}$$

where  $b_j$ 's ( $b_i = a_i, b_{g+i} = c_i$ ) are complex numbers.

It is noted that the permutation group acts on these  $\{b_r\}$  and  $\{a_r\}$ .

**2.2 Definition** [Ba1 , Ba2, W2, 3].

- (1) For a point  $(x_i, y_i) \in C_g$ , the unnormalized differentials of the first kind are defined by,

$$du_1^{(r,i)} := \frac{dx_i}{2y_i}, \quad du_2^{(r,i)} := \frac{(x_i - b_r)dx_i}{2y_i}, \quad \dots, \quad du_g^{(r,i)} := \frac{(x_i - b_r)^{g-1}dx_i}{2y_i}. \tag{2-3}$$

$$\begin{aligned} dv_1^{(i)} &:= \frac{P(x_i)dx_i}{2P'(a_1)(x_i - a_1)y_i}, \quad dv_2^{(i)} := \frac{P(x_i)dx_i}{2P'(a_2)(x_i - a_2)y_i}, \quad \dots, \\ dv_g^{(i)} &:= \frac{P(x_i)dx_i}{2P'(a_g)(x_i - a_g)y_i}. \end{aligned} \tag{2-4}$$

- (2) Let us define the Abel maps for  $g$ -th symmetric product of the curve  $C_g$ ,

$$u^{(r)} := (u_1^{(r)}, \dots, u_g^{(r)}) : \text{Sym}^g(C_g) \longrightarrow \mathbb{C}^g,$$

$$\left( u_k^{(r)}((x_1, y_1), \dots, (x_g, y_g)) := \sum_{i=1}^g \int_{\infty}^{(x_i, y_i)} du_k^{(r, i)} \right), \quad (2-5)$$

$$v := (v_1, \dots, v_g) : \text{Sym}^g(C_g) \longrightarrow \mathbb{C}^g,$$

$$\left( v_k((x_1, y_1), \dots, (x_g, y_g)) := \sum_{i=1}^g \int_{\infty}^{(x_i, y_i)} dv_k^{(i)} \right). \quad (2-6)$$

These coordinates are universal covering of the related Jacobian  $\mathcal{J}$ . The definition (2-6) [Ba2 p.382] is due to Weierstrass [W2, 3] and we call (2-6) *Weierstrass parameter*, though we choose different constant factor from the original one [W2, 3]. This parameterization is a key of the second solutions mentioned in §4.

### 2.3 Definition.

(1) Hyperelliptic *al* function is defined by [Ba2 p.340, W2, 3],

$$\text{al}_r(u) := \gamma_r \sqrt{F(b_r)}, \quad (2-7)$$

where  $\gamma_r := \sqrt{-1/P'(b_r)}$  and

$$\begin{aligned} F(x) &:= (x - x_1) \cdots (x - x_g) \\ &= (x - b_r - x_1 + b_r) \cdots (x - b_r - x_g + b_r). \end{aligned} \quad (2-8)$$

On the choice of  $\gamma_r$ , we will employ the convention of Baker [Ba2] instead of original one [W2, 3]. We note that  $\text{al}_r$ 's have mutually algebraic relations.

For later convenience, a polynomial associated with  $F(x)$  is introduced by

$$\begin{aligned} \pi_i^{(r)}(x) &:= \frac{F(x)}{x - x_i} \\ &= \chi_{i, g-1}^{(r)}(x - b_r)^{g-1} + \chi_{i, g-2}^{(r)}(x - b_r)^{g-2} + \cdots + \chi_{i, 1}^{(r)}(x - b_r) + \chi_{i, 0}^{(r)}. \end{aligned}$$

Then we have  $\chi_{i, g-1}^{(r)} \equiv 1$  and  $\chi_{i, 0}^{(r)} = F(b_r)/(x_i - b_r)$ . Further we introduce  $g \times g$ -matrices,

$$\mathcal{W}^{(r)} := \begin{pmatrix} \chi_{1,0}^{(r)} & \chi_{1,1}^{(r)} & \cdots & \chi_{1,g-1}^{(r)} \\ \chi_{2,0}^{(r)} & \chi_{2,1}^{(r)} & \cdots & \chi_{2,g-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots \\ \chi_{g,0}^{(r)} & \chi_{g,1}^{(r)} & \cdots & \chi_{g,g-1}^{(r)} \end{pmatrix}, \quad \mathcal{Y} = \begin{pmatrix} y_1 & & & \\ & y_2 & & \\ & & \ddots & \\ & & & y_g \end{pmatrix},$$

$$\begin{aligned}
\mathcal{M} &:= \begin{pmatrix} \frac{1}{x_1 - a_1} & \frac{1}{x_2 - a_1} & \cdots & \frac{1}{x_g - a_1} \\ \frac{1}{x_1 - a_2} & \frac{1}{x_2 - a_2} & \cdots & \frac{1}{x_g - a_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{x_1 - a_g} & \frac{1}{x_2 - a_g} & \cdots & \frac{1}{x_g - a_g} \end{pmatrix}, \\
\mathcal{P} &= \begin{pmatrix} \sqrt{\frac{P(x_1)}{Q(x_1)}} & & & \\ & \sqrt{\frac{P(x_2)}{Q(x_2)}} & & \\ & & \ddots & \\ & & & \sqrt{\frac{P(x_g)}{Q(x_g)}} \end{pmatrix}, \\
\mathcal{A} &= \begin{pmatrix} P'(a_1) & & & \\ & P'(a_2) & & \\ & & \ddots & \\ & & & P'(a_g) \end{pmatrix}, \quad \mathcal{F}' = \begin{pmatrix} F'(x_1) & & & \\ & F'(x_2) & & \\ & & \ddots & \\ & & & F'(x_g) \end{pmatrix},
\end{aligned}$$

where  $F'(x) := dF(x)/dx$ .

**2.3 Lemma.** *For these matrices, following relations hold:*

- (1) *The inverse matrix of  $\mathcal{W}^{(r)}$  is given by  $\mathcal{W}^{(r)-1} = \mathcal{F}^{(r)'}{}^{-1} \mathcal{V}^{(r)}$ , where  $\mathcal{V}^{(r)}$  is Vandermonde matrix,*

$$(2) \quad \mathcal{V}^{(r)} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ (x_1 - b_r) & (x_2 - b_r) & \cdots & (x_g - b_r) \\ (x_1 - b_r)^2 & (x_2 - b_r)^2 & \cdots & (x_g - b_r)^2 \\ \vdots & \vdots & \ddots & \vdots \\ (x_1 - b_r)^{g-1} & (x_2 - b_r)^{g-1} & \cdots & (x_g - b_r)^{g-1} \end{pmatrix}.$$

$$\det \mathcal{M} = \frac{(-1)^{g(g-1)/2} P(x_1, \dots, x_g) P(a_1, \dots, a_g)}{\prod_{k,l} (x_k - a_l)},$$

where

$$(3) \quad P(z_1, \dots, z_g) := \prod_{i < j} (z_i - z_j).$$

$$(\mathcal{M}\mathcal{P})^{-1} \mathcal{A} = \left[ \left( \frac{2y_i F(a_j)}{F'(x_i)(a_j - x_i)} \right)_{i,j} \right]. \quad (2-9)$$

*Proof.* (1) is obtained by direct computations. (2) is a well-known result [T]. Since the zero and singularity in the left hand side give the right hand side as

$$CP(x_1, \dots, x_g)P(a_1, \dots, a_g)/\prod_{k,l}(x_k - a_l),$$

for a certain constant  $C$ . In order to determine  $C$ , we multiply  $\prod_{k,l}(x_k - a_l)$  both sides and let  $x_1 = a_1, x_2 = a_2, \dots$ , and  $x_g = a_g$ . Then  $C$  is determined as above. (3) is obtained by the Laplace formula using the minor determinant for the inverse matrix. ■

Then we have following corollary.

**2.5 Corollary.** Let  $\partial_{u_i}^{(r)} := \partial/\partial u_i^{(r)}$ ,  $\partial_{v_i} := \partial/\partial v_i$ , and  $\partial_{x_i} := \partial/\partial x_i$ .

$$\begin{pmatrix} \partial_{u_1}^{(r)} \\ \partial_{u_2}^{(r)} \\ \vdots \\ \partial_{u_g}^{(r)} \end{pmatrix} = 2\mathcal{Y}\mathcal{F}'^{-1} \cdot {}^t\mathcal{W}^{(r)} \begin{pmatrix} \partial_{x_1} \\ \partial_{x_2} \\ \vdots \\ \partial_{x_g} \end{pmatrix}, \quad \begin{pmatrix} \partial_{v_1} \\ \partial_{v_2} \\ \vdots \\ \partial_{v_g} \end{pmatrix} = 2(\mathcal{MP})^{-1}\mathcal{A} \begin{pmatrix} \partial_{x_1} \\ \partial_{x_2} \\ \vdots \\ \partial_{x_g} \end{pmatrix}. \quad (2-10)$$

### §3. Relations between Hyperelliptic al Functions $(b_r, \infty)$ -type

In this section, we will give the first relation of hyperelliptic al function using the parameters  $u_1^{(r)}$  and  $u_g^{(r)}$  in (2-5).

#### 3.1 Theorem.

$$\frac{\partial}{\partial u_1^{(r)}} \frac{\partial}{\partial u_g^{(r)}} \log \text{al}_r = \frac{1}{2} \left( \frac{\text{al}_r^2}{\gamma_r^2} - \frac{f'(b_r)\gamma_r^2}{\text{al}_r^2} \right). \quad (3-1)$$

Here we will give a comment on Theorem 3.1. Let us fix the parameters  $x_2, \dots, x_g$  and regard  $\text{al}_r$  as a function of a parameter related to  $x_1$  over  $C_g$ . Then its divisor is  $(\text{al}_r) = (b_r, 0) - \infty$ . Further by letting  $t^2 = (x_i - b_r)$  around  $(b_r, 0)$ , the definition (2-3) shows,

$$du_1^{(r,i)}|_{(b_r,0)} = \frac{2}{\sqrt{f'(b_r)}} dt,$$

6

while for  $s^2 = 1/x$  around  $\infty$ ,

$$du_g^{(r,i)}|_{(\infty)} = -2ds.$$

Hence (3-1) can be regarded as an explicit representation of (1-3).

*Proof.* Instead of (3-1), we will prove following formula (3-2) in remainder in this section.

$$\frac{\partial}{\partial u_1^{(r)}} \frac{\partial}{\partial u_g^{(r)}} \log F(b_r) = F(b_r) - \frac{f'(b_r)}{F(b_r)}. \quad (3-2)$$

The strategy is essentially the same as [Ba3, Ma]. First we translate the words of the Jacobian into those of the curves; we rewrite the differentials  $u^{(r)}$ 's in terms of the differentials over curves as in (3-3). We count the residue of an integration and use a combinatorial trick. Then we will obtain (3-2).

From (2-10), we will express  $u^{(r)}$ 's by the affine coordinates  $x_i$ 's,

$$\begin{aligned} \frac{\partial}{\partial u_g^{(r)}} &= \sum_{i=1}^g \frac{2y_i}{F'(x_i)} \frac{\partial}{\partial x_i}, \\ \frac{\partial}{\partial u_1^{(r)}} &= \sum_{i=1}^g \frac{2y_i \chi_{i,0}^{(r)}}{F'(x_i)} \frac{\partial}{\partial x_i} = F(b_r) \sum_{i=1}^g \frac{2y_i}{(x_i - b_r) F'(x_i)} \frac{\partial}{\partial x_i}. \end{aligned} \quad (3-3)$$

Hence the right hand side of (3-2) becomes

$$-\frac{\partial^2}{\partial u_1^{(r)} \partial u_g^{(r)}} \log F(b_r) = F(b_r) \sum_{j=1, i=1}^g \frac{2y_j}{(x_i - b_r)^2 F'(x_j)} \frac{\partial}{\partial x_j} \frac{2y_i}{F'(x_i)(x_i - b_r)^2}. \quad (3-4)$$

Here we will note the derivative of  $F(x)$ , which is shown by direct computations.

$$\frac{\partial}{\partial x_k} \left( \left[ \frac{\partial}{\partial x} F(x) \right]_{x=x_k} \right) = \frac{1}{2} \left[ \frac{\partial^2}{\partial x^2} F(x) \right]_{x=x_k}.$$

Then (3-4) can be written as,

$$\begin{aligned} -\frac{\partial}{\partial u_1^{(r)}} \frac{\partial}{\partial u_g^{(r)}} \log F(b_r) &= F(b_r) \sum_{i=1}^g \frac{1}{F'(x_i)} \left[ \frac{\partial}{\partial x} \left( \frac{f(x)}{(x - b_r) F'(x)} \right) \right]_{x=x_i} \\ &\quad - F(b_r) \sum_{k,l,k \neq l} \frac{4y_k y_l}{(b_r - x_k)(b_r - x_l)(x_k - x_l) F'(x_k) F'(x_l)}. \end{aligned}$$

The proof of Theorem 3.1 finishes due to the following lemma. ■

**3.2 Lemma.** *Following relations hold:*

$$\sum_{k=1}^g \frac{1}{F'(x_k)} \left[ \frac{\partial}{\partial x} \left( \frac{f(x)}{(x-b_r)^2 F'(x)} \right) \right]_{x=x_k} = 1 - \frac{f'(b_r)}{F(b_r)^2}. \quad (3-5)$$

$$\sum_{k,l,k \neq l} \frac{2y_k y_l}{(b_r - x_k)(b_r - x_l)(x_k - x_l)F'(x_k)F'(x_l)} = 0. \quad (3-6)$$

*Proof.* : (3-5) will be proved by the following residual computations: Let  $\partial C_g^o$  be the boundary of a polygon representation  $C_g^o$  of  $C_g$ ,

$$\oint_{\partial C_g^o} \frac{f(x)}{(x-b_r)^2 F(x)^2} dx = 0. \quad (3-7)$$

The divisor of the integrand of (3-7) is given by,

$$\left( \frac{f(x)}{(x-b_r)^2 F(x)^2} dx \right) = 3 \sum_{i=1, b_i \neq b_r}^{2g+1} (b_i, 0) - (b_r, 0) - 2 \sum_{i=1}^g (x_i, y_i) - 2 \sum_{i=1}^g (x_i, -y_i) - \infty.$$

We check these poles: First we consider the contribution around  $\infty$  point. Noting that the local parameter  $t$  at  $\infty$  is  $x = 1/t^2$ ,

$$\text{res}_{\infty} \frac{f(x)}{(x-b_r)^2 F(x)^2} dx = -2.$$

Since the local parameter  $t$  at  $(x_k, \pm y_k)$  is  $t = x - x_k$ , we have

$$\text{res}_{(x_k, \pm y_k)} \frac{f(x)}{(x-b_r)^2 F(x)^2} dx = \frac{1}{F'(x_k)} \left[ \frac{\partial}{\partial x} \left( \frac{f(x)}{(x-b_r)^2 F'(x)} \right) \right]_{x=x_k}.$$

For each branch point  $(b_r, 0)$ , the local parameter  $t$  is  $t^2 = x - b_r$  and thus

$$\text{res}_{(b_r, 0)} \frac{f(x)}{(x-b_r)^2 F(x)^2} dx = 2 \frac{f'(b_r)}{F(b_r)^2}.$$

By arranging them, we obtain (3-5).

On the other hand, (3-6) can be proved by using a trick: for  $i \neq j$ ,

$$\frac{1}{(b_r - x_k)(b_r - x_l)(x_k - x_l)} = \frac{1}{(x_k - x_l)^2} \left( \frac{1}{(b_r - x_k)} - \frac{1}{(b_r - x_l)} \right).$$

■



#### §4. Relations between Hyperelliptic al Functions: $(a_1, a_2)$ -type

In the previous section, we have a solution with a duality between a finite ramified point and  $\infty$ -point. In this section, we will give a relation between hyperelliptic al functions using the Weierstrass parameter (2-6). The relation has a duality between finite ramified points  $(a_r, 0)$  and  $(a_s, 0)$ .

**4.1 Theorem.** *For  $r \neq s$ , we obtain*

$$\frac{\partial}{\partial v_r} \frac{\partial}{\partial v_s} \log \frac{\text{al}_r}{\text{al}_s} = \frac{1}{2(a_r - a_s)} \left( f'(a_r) \frac{\gamma_r^2 \text{al}_s^2}{\gamma_s^2 \text{al}_r^2} + f'(a_s) \frac{\gamma_s^2 \text{al}_r^2}{\gamma_r^2 \text{al}_s^2} \right). \quad (4-1)$$

Before we prove it, we will give some comments: Let us fix the parameter  $x_2, \dots, x_g$  and regard  $\text{al}_r/\text{al}_s (\propto \sqrt{F(a_r)/F(a_s)})$  as a function of  $x_1$  over  $C_g$ . Then its divisor is  $(\text{al}_r/\text{al}_s) = (a_r, 0) - (a_s, 0)$ . By letting  $t_r^2 = (x_i - a_r)$  around  $(a_r, 0)$ , infinitesimal value of Weierstrass parameter (2-4) is given,

$$dv_r^{(i)}|_{(a_r, 0)} = \frac{1}{\sqrt{f'(a_r)}} dt_r.$$

Thus (4-1) is also a concrete expression of (1-3).

*Proof.* Similar to the proof of Theorem 3.1, let us prove the theorem. Without loss of generality, we will prove the following relation instead of (4-1):

$$\frac{\partial}{\partial v_1} \frac{\partial}{\partial v_2} \log \frac{F(a_1)}{F(a_2)} = \frac{F(a_1)F(a_2)}{(a_1 - a_2)} \left( \frac{f'(a_1)}{F(a_1)^2} + \frac{f'(a_2)}{F(a_2)^2} \right). \quad (4-2)$$

From (2-9) and (2-10), the derivative  $v$ 's are expressed by the affine coordinate  $x_i$ 's,

$$\frac{\partial}{\partial v_r} = F(a_r) \sum_{j=1}^g \frac{2y_j}{F'(x_j)(x_j - a_r)} \frac{\partial}{\partial x_j}.$$

The right hand side of (4-2) becomes,

$$\frac{\partial^2}{\partial v_1 \partial v_2} \log \frac{F(a_1)}{F(a_2)} = F(a_1) \sum_{j=1, i=1}^g \frac{2y_j}{(x_i - a_1)F'(x_j)} \frac{\partial}{\partial x_j} \frac{2y_i F(a_2)}{F'(x_i)(x_i - a_2)} \frac{(a_1 - a_2)}{(x_i - a_1)(x_i - a_2)}. \quad (4-3)$$

The right hand side of (4-3) is

$$F(a_1)F(a_2) \sum_{i=1}^g \frac{1}{F'(x_i)} \left[ \frac{\partial}{\partial x} \left( \frac{f(x)(a_2 - a_1)}{(x - a_1)^2(x - a_2)^2 F'(x)} \right) \right]_{x=x_i} \\ - F(a_1)F(a_2) \sum_{k,l,k \neq l} \frac{2y_k y_l (a_2 - a_1)}{F'(x_k)F'(x_l)(x_l - a_1)(x_k - a_2)(x_k - a_1)(x_l - a_2)(x_l - x_k)}.$$

Then the proof of Theorem 4.1 is completely done due to the following lemma. ■

**4.2 Lemma.** *Following relations hold:*

$$\sum_{i=1}^g \frac{1}{F'(x_i)} \left[ \frac{\partial}{\partial x} \left( \frac{f(x)}{(x - a_1)^2(x - a_2)^2 F'(x)} \right) \right]_{x=x_i} = \frac{1}{(a_1 - a_2)^2} \left( \frac{f'(a_1)}{F(a_1)^2} - \frac{f'(a_2)}{F(a_2)^2} \right). \quad (4-4)$$

$$\sum_{k,l,k \neq l} \frac{2y_k y_l (a_2 - a_1)}{F'(x_k)F'(x_l)(x_l - a_1)(x_k - a_2)(x_k - a_1)(x_l - a_2)(x_l - x_k)} = 0. \quad (4-5)$$

*Proof.* : Similar to Lemma 3-2, we consider an integral,

$$\oint_{\partial C_g^o} \frac{f(x)}{(x - a_1)^2(x - a_2)^2 F(x)^2} dx = 0. \quad (4-6)$$

As the divisor of the integrand of (4-6) is

$$\left( \frac{f(x)}{(x - a_1)^2(x - a_2)^2 F(x)^2} dx \right) \\ = 3 \sum_{i=1, b_i \neq a_1, a_2}^{2g+1} (b_i, 0) - (a_1, 0) - (a_2, 0) - 2 \sum_{i=1}^g (x_i, y_i) - 2 \sum_{i=1}^g (x_i, -y_i) + 3\infty, \quad (4-9)$$

we count residual contributions from each terms as in the proof of Lemma 3-2 and obtain (4-4). Considering the symmetry, (4-5) is easily obtained. ■

## REFERENCES

- [Ba1] Baker, H. F., *Abelian functions – Abel’s theorem and the allied theory including the theory of the theta functions* –, Cambridge Univ. Press, 1897, republication 1995.
- [Ba2] ———, *On the hyperelliptic sigma functions*, Amer. J. of Math. **XX** (1898), 301-384.
- [Ba3] ———, *On a system of differential equations leading to periodic functions*, Acta math. **27** (1903), 135-156.
- [Ma] Matsutani, S., *Explicit Hyperelliptic Solutions of Modified Korteweg-de Vries Equation: Essentials of Miura Transformation*, J. Phys. A. Math. & Gen. **35** (2002), 4321-4333.
- [Mu] Mumford, D., *Tata Lectures on Theta, vol II*, Birkhäuser, Boston, 1984.

- [T] Takagi, T., *Daisuu-Gaku-Kougi (Lecture of Algebra)*, Kyouritsu, Tokyo, 1930. (japanese)
- [W1] Weierstrass, K., *Mathematische Werke I*, Mayer und Müller, Berlin, 1894.
- [W2] Weierstrass, K., *Beitrage zur Theorie der Abel'schen Functionen*, Beilage zum Jahresbericht über des Gymnasium zu Braunsberg in dem Schuljahre 1848-1849, in [W1] (1849).
- [W3] Weierstrass K., *Zur Theorie der Abel'schen Functionen*, in [W1], Aus dem Crelle'schen Journal **47** (1854).

address:8-21-1 Higashi-Linkan Sagamihara 228-0811 Japan

e-mail:RXB01142@nifty.ne.jp