Proof of the generalized Lieb-Wehrl conjecture for integer indices larger than one

Ayumu Sugita

Yukawa Institute for Theoretical Physics, K itashirakawa-0 iwakecho, K yoto, K yoto 606-8502, Japan

(Dated: March 28, 2024)

# Abstract

G nutzm ann and Zyczkowski have proposed the RenyiW ehrl entropy as a generalization of the W ehrl entropy, and conjectured that its m in im um is obtained for coherent states. We prove this conjecture for the Renyi index q = 2;3; ::: in the cases of compact sem is imple Lie groups. A general formula for the minimum value is given.

PACS num bers: 0220Qs, 05.45Mt

#### I. INTRODUCTION

The Husim i function [1] is a useful tool to investigate dynam ical properties of quantum systems. We had proposed a classical entropy [2] using the Husim i function and conjectured that the minimum of the entropy is obtained for coherent states. His conjecture was soon proved by Lieb [3]. However, Lieb's analogous conjecture for spin coherent state in the same paper has remained unproved for more than 20 years.

The Husim i function for the density operator ^ is de ned as

where j i is a coherent state. The generalized coherent states de ned by Perelom ov [4] can also be used instead of the standard coherent state. G nutzm ann and Zyczkowski [5] have recently generalized the Wehrlentropy by analogy with the well-known Renyi entropy [6]. The Renyi-Wehrlentropy is de ned as

$$S^{(q)}[^{]} = \frac{1}{1 - q} \log M^{(q)}[^{]};$$
 (2)

H ere, M  $^{(q)}$  is the m om ent of the H usim i function

$$M^{(q)}[^] = c d () f () g^q$$
 (3)

where d ( ) is the Haar m easure and c is a normalization constant. In the  $\lim it q! 1$ , it reproduces the usual form of the entropy

$$\lim_{\substack{\text{cl. } 1}} S^{(q)}[^{\uparrow}] = S[^{\uparrow}] \qquad c \quad d \quad ( ) \quad ( ) \log \quad ( ) : \tag{4}$$

The generalized Lieb-Wehrl conjecture formed in [5] is the following.

C on jecture 1 The minimum value of the Renyi-W ehrlentropy for q > 0 is obtained for coherent states.

We prove this conjecture for integer indices q=2 in the cases of the coherent states of compact sem is in ple Lie groups. The case of spin (SU (2)) coherent states has already been proved [5, 7]. In the case of the standard coherent state, q=1 corresponds to the original Wehrl's conjecture proved in [3], and theorem 3 therein gives the proof for arbitrary q>1.

### II. GENERALIZED COHERENT STATES

Before coming to the proof, let us review the denition and some properties of the generalized coherent states [4]. Let G be a compact semisimple Lie group. The Lie algebra of G can be written in so-called Cartan basis fH ; E g. An irreducible representation of G is characterized by the lowest (or the highest) weight. Let j i be the lowest weight state with the weight . We denote the irreducible representation space specified by j i as D . Coherent states in D are obtained by the action of the group G on j i, which can be written explicitly as [11]

$$j i = N () exp(E) j i$$
; (5)

Here, N ( ) is the normalization constant and runs over all positive roots. The expansion of the exponential function becomes a nite series because D is nite dimensional. Therefore

$$^{\sim}() \frac{h j i}{N ()}$$
 (6)

is a polynomial of for any state ji. We will use this fact later to prove lem ma 1.

The \resolution of unity"

$$I_D = \dim D \qquad d \ ( \ )j \ ih \ j \qquad \qquad (7)$$

is valid, and hence the normalization constant in (3) should be taken as

$$G_D = \dim D ; (8)$$

in order to satisfy the normalization condition  $M^{(1)} = 1$ .

# III. PROOF OF THE MAIN RESULT

Turning now to the proof, we rst note the following lem ma.

Lem m a 1 If  $^{n}$  m in im izes S  $^{(q)}$   $^{(q)}$  for som e q > 0,  $^{n}$  m ust be a pure state.

This is a generalization of lem m a 2 in  $[\beta]$ , and can be proved in the same way (see the appendix). Hereafter we concentrate on pure states.

Let us consider the sim plest case q=2. We will see later that the discussion in this case can be easily generalized. We have to prove that the maximum of the second moment of the Husim i function

$$M_{\tilde{I}i}^{(2)} c_{D} d() \hat{J} \tilde{J} i \hat{J}$$

$$(9)$$

is given by coherent states.

Let us rewrite (9) as

$$M_{\vec{j},i}^{(2)} = (h'j h'j) c_{b} \quad d () (ji ji) (h j h j) (j'i j'i):$$
 (10)

The key observation is that the tensor product of the coherent states j i j i is a coherent state in the irreducible representation space D<sub>2</sub> whose lowest weight state is j i j i. The same idea is used in [7] for SU (2). It can be shown explicitly as

$$(H_{j} I + I H_{j})j i j i = 2_{j}j i j i$$
 (11)

$$(E I + I E )j i j i = 0 (12)$$

$$ji ji = N()^{2} exp[(E I + I E)]j i j i$$
 (13)

where runs over positive roots. Therefore  $^{R}$  d ( ) (j i j i) (h j h j) is proportional to the projection operator to D  $_{2}$  in D D , i.e.

$$c_{D}$$
 d ( ) (j i j i) (h j h j) =  $\frac{\text{dim } D}{\text{dim } D_{2}} P_{D_{2}}$ : (14)

Here the proportional constant is determined from (7). Hence

$$M_{\mathring{J}\dot{i}}^{(2)} = \frac{\dim D}{\dim D_{2}} (h' \dot{j} h' \dot{j}) P_{D_{2}} (\mathring{J}\dot{i} \mathring{J}\dot{i})$$

$$\tag{15}$$

and we have

$$\vec{J} \text{ i } \vec{J} \text{ i 2 D}_{2} \quad \text{( )} \quad M_{\vec{J} \text{ i}}^{(2)} = M_{\text{max}}^{(2)} \quad \frac{\dim D}{\dim D_{2}} :$$
(16)

Then we should prove

$$\vec{j} i : \text{coherent state} () \quad \vec{j} i \quad \vec{j} i 2 D_2 :$$
(17)

LHS) RHS is obvious, because j i j i is obtained by the action of G on j i j i which belongs to D<sub>2</sub> by de nition.

Then we prove LHS ( RHS.Let us consider the quadratic Casim ir operator  $C_2$  of the group G. By taking an orthonorm albasis  $fT_ag$  of the Lie algebra, it can be written as

$$C_2 = T^2$$
  $X_a$ : (18)

In the tensor product space D D, the quadratic C asim ir operator is written as

$$C_2^{(2)}$$
  $X$   $(T_a$   $1+1$   $T_a)^2 = T^2$   $1+1$   $T^2+2$   $T_a$   $T_a$ : (19)

T herefore

$$(h' j h' j) C_2^{(2)} (j i j i) = 2 < T^2 > +2 < T >^2$$
 (20)

where the symbol <> denotes the expectation value for j i.

Here we note the lem ma proved in [8]:

Lem m a 2 (Delbourgo and Fox) The minimum of the uncertainty (T) $^2$  < T $^2$  > < T > $^2$  is obtained for coherent states

$$(T)^2$$
 is m in im um  $()$  j i : coherent state: (21)

Since  $C_2 = T^2$  is a constant in D , < T > 2 is maximum when if i is a coherent state, and hence  $C_2^{(2)}$  takes the maximum value at D<sub>2</sub> because it contains the tensor product of the coherent states. Therefore

$$7i \quad 7i2 \quad D_2 =$$
 < T > 2 is maximum = )  $7i :$  coherent state: (22)

This completes the proof of (17).

The generalization to the higher m om ents is straightforward. By considering the tensor product space D  $^q$  D D ::: D , one can easily show

$$M_{Ji}^{(q)} = \frac{\dim D}{\dim D_q} h' j^q P_{D_q} J' i^q$$
 (23)

and

$$h' j^q C_2^{(q)} j' i^q = q < T^2 > + q(q 1) < T >^2$$
 (24)

in obvious notation. From these two formulae we can conclude

$$M_{Ji}^{(q)} = M_{max}^{(q)}$$
 ()  $Ji^{q} 2 D_{q}$  ()  $Ji^{z}$  coherent state: (25)

Combining this and Lemma 1, we obtain the main result of this paper:

Theorem 1 Let  $S^{(q)}$  be the Renyi-W ehrlentropy with an integer index q-2 of a compact sem is imple Lie group. The minimum value of  $S^{(q)}$  is obtained for coherent states

$$S^{(q)}$$
 [^] is m in im um () ^= j ih' j; j' i : coherent state: (26)

The minimum value  $S_{m\ in}^{\ (q)}$  is given by

$$S_{m \text{ in}}^{(q)} = \frac{1}{1 + q} \log \frac{\dim D}{\dim D_q}!$$
(27)

where D is the irreducible representation space in which S (q) is de ned.

## IV. M IN IM UM VALUES FOR SOME SPECIAL CASES

Finally, let us exam ine the general formula for the minima (27) in some special cases. Cases (i), (ii), and (iii) in the following can be used to investigate the dynamics of bosons, fermions and distinguishable particles, respectively [12] [9].

(i) G = SU(N),  $D = [1^m]$  (sym m etric product of m fundam ental representations)

$$\dim D_{q} = \dim [1^{qm}] = \frac{(N + qm - 1)!}{(N - 1)! (qm)!}$$
 (28)

$$S_{m \text{ in}}^{(q)} = \frac{1}{1 + q} \log \frac{(N + m) (qm + 1)}{(N + qm) (m + 1)};$$
 (29)

Assuming that (29) holds also for non-integer q, we obtain

$$\lim_{\alpha!} S_{m \text{ in}}^{(q)} = m f (N + m) (m + 1)g$$
 (30)

where (z)  $\frac{d}{dz} \log$  (z) is the digamma function. This result agrees with the value of S  $^{(1)}$  for coherent states given in [10], except for the shift by  $\ln N$  due to another normalization of the coherent states.

(ii) G = SU (N), D = [m] (anti-sym m etric product of m fundam ental representations)

$$\dim D_{q} = \dim [m^{q}] = \prod_{j=0}^{m_{1}} \frac{(N+q-j-1)!j!}{(N-j-1)!(q+j)!}$$
(31)

$$S_{m in}^{(q)} = \frac{1}{1 q} \sum_{j=0}^{n_{X}} \log \frac{(N+1 j) (q+j+1)}{(N+q j) (j+2)};$$
 (32)

In this case, the minimum value for q = 1 is expected to be

$$\lim_{q!} S_{m \text{ in}}^{(q)} = \int_{j=0}^{m_{X}} f(N + j + 1) \qquad (j+2)g;$$
(33)

(iii) 
$$G = SU(N)$$
 :::  $SU(N), D = [1]^m$ 

$$\dim D_q = \dim [1^q]^m = \frac{(N+q-1)!}{(N-1)!q!}^! m$$
(34)

$$S_{m \text{ in}}^{(q)} = \frac{m}{1 + q} \log \frac{(N + 1) (q + 1)}{(N + q)};$$
 (35)

The minimum value for q = 1 is expected to be

$$\lim_{\alpha!} S_{m \text{ in}}^{(q)} = m f (N + 1)$$
 (2)g: (36)

A cknow ledgm ents

The author thanks Professor Gnutzmann for letting him know about reference [7]. He also thanks Professor Zyczkowski for helpful discussions.

## APPENDIX A: PROOF OF LEMMA 1

Here we prove lem m a 1: if  $^m in im izes S^{(q)} [^n]$  for som e q > 0,  $^m ust be a pure state.$ 

We consider three cases (i) 0 < q < 1, (ii) q = 1 and (iii) 1 < q separately. ^ m in im izes M (q) and S in cases (i) and (ii), respectively, and maximizes M (q) in case (iii). Let us decompose the density operator as  $\hat{p}$  where  $\hat{p}$  is  $\hat{p}$  and  $\hat{p}$  is  $\hat{p}$  in  $\hat{p}$  a pure state. Since  $m^{(q)}(x) = x^q (0 < q < 1)$  and  $s(x) = x \log x$  are concave, and  $m^{(q)}(x) = x^{(q)}(q > 1)$  is convex,

$$M^{(q)}[^{\uparrow}]$$
  ${}^{X}_{i}M^{(q)}[^{\uparrow}_{i}]$   $(0 < q < 1)$  (A1)

$$X^{i}$$

$$S [^{]} \qquad _{i}S [^{]} \qquad (A 2)$$

The equality must hold in at least one of the three inequalities. Then  $_{i}() = _{j}()$ , and hence  $\tilde{i}() = \tilde{j}()$  in the notation of (6), almost everywhere for all i; j. Since  $\tilde{i}$  is a polynomial,  $i = a_j$ , and hence  $j_i = a_j$ , with  $j_i = 1$ . Therefore  $j_i = i$  for all i; j, which means 'is a pure state.

<sup>[1]</sup> K. Husim i, Proc. Phys. M ath. Soc. Japan 22 264 (1940).

- [2] A.W ehrl, Reps. Math. Phys. 16, 353 (1979).
- [3] E.H.Lieb, Commun.Math.Phys. 62, 35 (1978).
- [4] A. Perelom ov, Generalized Coherent States and Their Applications, Springer-Verlag Berlin Heidelberg (1986).
- [5] S.G nutzm ann and K. Zyczkowski, J. Phys. A 34 (2001) 10123.
- [6] A.Renyi, Probability Theory, North-Holland, Amsterdam (1970).
- [7] P.Schupp, Commun.Math.Phys.207 (1999) 481.
- [8] R.Delbourgo and J.R.Fox, J.Phys. A 10 (1977) L233.
- [9] A. Sugita, arX iv: nlin CD /0112042
- [10] W .Slom czynski and K .Zyczkowski, Phys. Rev. Lett. 80 (1998) 1880.
- [11] In fact, (5) cannot represent the coherent states orthogonal to j i. However, it does not matter when we consider entropies such as (2) because the measure of the set of such states is zero.
- [12] U (N), instead of SU (N), is used in [9] for bosons and ferm ions. U (N) is not sem is imple.

  However, U (N) U (1) SU (N), and the Abelian subgroup U (1) corresponds to the total phase which is irrelevant for the Husim i functions.