

Hierarchy of Chaotic maps with an invariant measure and their coupling.

M .A .Jafarizadeh^{a,b,c}, S.Behnia^{d,e,b}.

^aDepartment of Theoretical Physics and Astrophysics, Tabriz University, Tabriz 51664, Iran.

^bInstitute for Studies in Theoretical Physics and Mathematics, Teheran 19395-1795, Iran.

^cPure and Applied Science Research Center, Tabriz 51664, Iran.

^dPlasma Physics Research Center, IAU, Teheran 14835-159, Iran.

^eDepartment of Physics, IAU, Urmia, Iran.

March 28, 2024

E-mail: jafarizadeh@ark.tabrizu.ac.ir

Abstract

Hierarchy of one-parameter families of chaotic maps with an invariant measure have been introduced, where their appropriate coupling has lead to the generation of some coupled chaotic maps with an invariant measure. It is shown that these chaotic maps (also the coupled maps) do not undergo any period doubling or period- n -tupling cascade bifurcation to chaos, but they have either single fixed point attractor at certain values of the parameters or they are ergodic in the complementary region. Using the invariant measure or Sinai-Ruelle-Bowen measure the Kolmogorov-Sinai entropy of the chaotic maps (coupled maps) have been calculated analytically, where the numerical simulations support the results.

Keywords: coupled chaotic map, chaos, Invariant measure, Kolmogorov-sinai entropy, Lyapunov characteristic exponent, Ergodic dynamical systems.

PACS Index: 05.45.Ra, 05.45.Jn, 05.45.Tp

INTRODUCTION

In recent years chaos or more properly dynamical systems have become an important area of research activity. One of the landmarks in it was the introduction of the concept of Sinai-Ruelle-Bowen (SRB) measure or natural invariant measure. This is roughly speaking a measure that is supported on an attractor and it describes the statistics of the long time behavior of the orbits for almost every initial condition in the corresponding basin of attractor. This measure can be obtained by computing the fixed points of the so called Frobenius-Perron (FP) operator which can be viewed as a differential-integral operator, hence, exact determination of invariant measure of dynamical systems is rather a nontrivial task, such that invariant measure of few dynamical systems such as one-parameter families of one-dimensional piecewise linear maps like the Baker and tent maps [1, 2, 3, 4] or unimodal maps like logistic map for certain values of its parameter (Ulam-Von Neumann map), one-dimensional cubic and quintic maps [5, 6] can be derived analytically. In most of cases only numerical algorithms, as an example Ulam's method [7, 8, 9] are used for computation of fixed points of FP operator.

Here in this article, we give a hierarchy of one-parameter families $\mathcal{F}_N(\cdot; x)$ of maps of interval $[0; 1]$ with an invariant measure, such that Ulam-Von Neumann map, one-dimensional cubic and quintic maps correspond to three first maps of this hierarchy for particular value of their parameter. Also the map appearing in the study of field driven thermodynamic systems [4] is topologically conjugate to the first map (generalized Ulam-Von Neumann) of our hierarchy of family of one parameter chaotic maps. Then, by an appropriate coupling of these maps, we generate some coupled chaotic maps with an invariant measure. We have been able to derived analytically their invariant

measure. Using this measure, we have calculated analytically, Kolmogorov-Sinai(KS) entropy or equivalently positive Lyapunov characteristic exponent of these maps, where the numerical simulations support the analytic calculations. Also it is shown that these maps have another interesting property, that is, they bifurcate to chaotic regime without having any period doubling or period-n-tupling scenario.

The paper organized as follows. In section II the hierarchy of one parameter families of chaotic maps have been introduced. Section III is devoted to the appropriate coupling of these maps. In section IV we talk about their invariant measure. In section V, using the invariant measure, we give their KS entropy. Finally in section VI we try to support the analytic calculations with the numerical simulations. The paper ends with brief conclusion and two appendices which contains all algebraic calculations and proofs.

II Hierarchy of one-parameter families of chaotic maps

One of the exactly solvable one dimensional maps or the map with invariant measure is the well known Ulam-Von Neumann map [10]

$f(x) = x(1-x)$ (Logistic map for $r=1$). In reference [5, 6] the Ulam-Von Neumann map with KS-entropy $\ln(2)$ has been generalized to exactly solvable one dimensional cubic and quintic maps with KS-entropy $\ln(3)$ and $\ln(5)$, respectively. Here we generalize these maps to a Hierarchy of one parameter families of maps with invariant measure, where the map of Reference [4] is topologically conjugate to the first map (generalized Ulam-Von Neumann) of this hierarchy.

These one-parameter families of chaotic maps of the interval $[0;1]$ with an invariant measure can be defined as the ratio of polynomials of degree N :

$$\begin{aligned} f_N^{(1)}(x; \lambda) &= \frac{(\lambda^2 - 1) + (\lambda^2 - 1)^N {}_2F_1\left(-N; N; \frac{1}{2}; x\right)}{(\lambda^2 + 1) + (\lambda^2 - 1)(\lambda^2 - 1)^N {}_2F_1\left(-N; N; \frac{1}{2}; x\right)} \\ &= \frac{(\lambda^2 - 1) (T_N(\sqrt{x}))^2}{1 + (\lambda^2 - 1) (T_N(\sqrt{x}))^2} ; \end{aligned} \quad (\text{II-1})$$

where N is an integer greater than one. Also

$${}_2F_1\left(-N; N; \frac{1}{2}; x\right) = (\lambda^2 - 1)^N \cos(2N \arccos \sqrt{x}) = (\lambda^2 - 1)^N T_{2N}(\sqrt{x})$$

is the hypergeometric polynomials of degree N and T_N are Chebyshev polynomials of type I [11], respectively. Obviously these map the unit interval $[0;1]$ into itself.

$f_N(\lambda; x)$ is $(N-1)$ -model map, that is it has $(N-1)$ critical points in unit interval $[0;1]$ (see Figures 1,2), since its derivative is proportional to derivative of hypergeometric polynomial ${}_2F_1\left(-N; N; \frac{1}{2}; x\right)$ which is itself a hypergeometric polynomial of degree $(N-1)$, hence it has $(N-1)$ real roots in unit interval $[0;1]$. Defining Schwarzian derivative $S(f_N(x; \lambda))$ as [12]:

$$S(f_N(x; \lambda)) = \frac{f_N''(x; \lambda)}{f_N'(x; \lambda)} - \frac{3}{2} \left(\frac{f_N''(x; \lambda)}{f_N'(x; \lambda)} \right)^2 = \frac{f_N''(x; \lambda)}{f_N'(x; \lambda)} - \frac{1}{2} \left(\frac{f_N''(x; \lambda)}{f_N'(x; \lambda)} \right)^2 ;$$

with a prime denoting a single differential, one can show that:

$$S(f_N(x; \lambda)) = S({}_2F_1(-N; N; \frac{1}{2}; x)) \leq 0;$$

since $\frac{d}{dx}({}_2F_1(-N; N; \frac{1}{2}; x))$ can be written as:

$$\frac{d}{dx}({}_2F_1(-N; N; \frac{1}{2}; x)) = A \prod_{i=1}^{N-1} (x - x_i);$$

with $0 \leq x_1 < x_2 < x_3 < \dots < x_{N-1} \leq 1$, then we have:

$$S({}_2F_1(-N; N; \frac{1}{2}; x)) = \frac{1}{2} \sum_{j=1}^{N-1} \frac{1}{(x - x_j)^2} - \frac{1}{2} \sum_{j=1}^{N-1} \frac{1}{(x - x_j)^2} \leq 0;$$

Therefore, the maps $T_N(x; \lambda)$ have at most $N + 1$ attracting periodic orbits [12]. As we will show at the end of this section, these maps have only a single period one stable fixed points or they are ergodic.

Denoting n -composition of functions $T(x; \lambda)$ by $T^{(n)}$, it is straightforward to show that the derivative of $T^{(n)}$ at its possible n periodic points of an n -cycle:

$$x_2 = T(x_1; \lambda); x_3 = T(x_2; \lambda); \dots; x_n = T(x_{n-1}; \lambda)$$

$$j \frac{d}{dx} T^{(n)} = j \frac{d}{dx} T^{(1,2)} \circ T^{(1,2)} \circ \dots \circ T^{(1,2)} = \prod_{k=1}^n j \frac{d}{dx} T^{(1,2)}(x_k; \lambda) \quad (II-2)$$

since for $x_k \in [0; 1]$ we have:

$$j \frac{d}{dx} T^{(1,2)}(x_k; \lambda) = j \frac{d}{dx} (1 - \lambda^2 x_k) = -2\lambda^2 x_k$$

therefore,

$$j \frac{d}{dx} T^{(n)} = (-2\lambda^2)^n \prod_{k=1}^n x_k$$

Hence the above expression is definitely greater than one for $\frac{1}{N} < \lambda < N$, that is: maps do not have any kind of n -cycle or periodic orbits for $\frac{1}{N} < \lambda < N$, actually they are ergodic for this interval of parameter. From (II-2) it follows the values of $j \frac{d}{dx} T^{(n)}$ at n periodic points of the n -cycle belongs to interval $[0, 1]$, varies between $(N)^n$ and $(\frac{N}{\lambda})^n$ for $\lambda < \frac{1}{N}$ and between $(\frac{N}{\lambda})^n$ and $(N)^n$ for $\lambda > N$, respectively.

From the definition of these maps, we see that for odd N , both $x = 0$ and $x = 1$ belong to one of the n -cycles, while for even N , only $x = 1$ belongs to one of the n -cycles of $T_N(x; \lambda)$.

For $\lambda < (\frac{1}{N})$ ($\lambda > N$), the formula (II-2) implies that for those cases in which $x = 0$ ($x = 1$) belongs to one of n -cycles we will have $j \frac{d}{dx} T^{(n)} < 1$, hence the curve of $T^{(n)}$ starts at $x = 0$ ($x = 1$) beneath the bisector and then crosses it at the next (previous) periodic point with slope greater than

one, since the formula (II-2) implies that the slope of fixed points increases with the increasing (decreasing) of $|x_k|$, therefore at all periodic points of n -cycles except for $x = 0$ ($x = 1$) the slope is greater than one that is they are unstable, this is possible only if $x = 0$ ($x = 1$) is the only period one fixed point of these maps.

Hence all n -cycles except for possible period one fixed points $x = 0$ and $x = 1$ are unstable, where for $\alpha \in [0; \frac{1}{N}]$, the fixed point $x = 0$ is stable in maps $f_N(x; \alpha)$ (for odd integer values of N), while for $\alpha \in [\frac{1}{N}; 1)$ the $x = 1$ is stable fixed point in maps (see Figures 3,4 for $N = 2, 3$).

Below we also introduce their conjugate or isomorphic maps which will be very useful in derivation of their invariant measure and calculation of their KS-entropy in the next sections. Conjugacy means that the invertible map $h(x) = \frac{1-x}{x}$ maps $I = [0; 1]$ into $[0; 1)$ and transform maps $f_N(x; \alpha)$ into $\tilde{f}_N(x; \alpha)$ defined as:

$$\tilde{f}_N(x; \alpha) = h \circ f_N(x; \alpha) \circ h^{-1} = \frac{1}{2} \tan^2(N \arctan^{\frac{1}{2}} \frac{x}{1-x}) \quad (\text{II-3})$$

As an example we give below some of these maps:

$$f_2(x; \alpha) = \frac{\alpha^2 (2x-1)^2}{4x(1-x) + \alpha^2 (2x-1)^2}; \quad (\text{II-4})$$

$$f_3(x; \alpha) = \frac{\alpha^2 x(4x-3)^2}{2x(4x-3)^2 + (1-x)(4x-1)^2}; \quad (\text{II-5})$$

$$f_4(x; \alpha) = \frac{\alpha^2 (1-8x(1-x))^2}{2(1-8x(1-x))^2 + 16x(1-x)(1-2x)^2}; \quad (\text{II-6})$$

$$f_5(x; \alpha) = \frac{\alpha^2 x(16x^2-20x+5)^2}{2x(16x^2-20x+5)^2 + x(16x^2-20x+5)^2 + 1}; \quad (\text{II-7})$$

III Hierarchy of coupled map

Using the hierarchy of families of one-parameter chaotic maps (II-1), we can generate new hierarchy of coupled maps with an invariant measure. Hence,

by introducing a new parameter g as a coupling parameter we form coupling among the above mentioned maps, where they can be coupled through α and β functions defined as:

$$f(x) = \frac{q}{(1 + gx)^2}; \quad (\text{III-1})$$

$$f_N(x) = \frac{B_N\left(\frac{1}{x}\right)}{A_N\left(\frac{1}{x}\right)} \quad (\text{III-2})$$

with $B_N(x)$ and $A_N(x)$ defined as:

$$A_N(x) = \sum_{k=0}^{\lfloor \frac{N}{2} \rfloor} C_{2k}^N x^k; \quad (\text{III-3})$$

$$B_N(x) = \sum_{k=0}^{\lfloor \frac{N-1}{2} \rfloor} C_{2k+1}^N x^k; \quad (\text{III-4})$$

Now, the hierarchy of the coupled maps can be defined as:

$$\begin{aligned} X_1 &= f_{N_1}(x_1; f_{N_2}(x_2)) \\ X_2 &= f_{N_2}(x_2; f_{N_1}(x_1)); \end{aligned} \quad (\text{III-5})$$

Their conjugate or isomorphic maps are defined as:

$$\begin{aligned} X_{N_1}(x_1; f_{N_2}(x_2)) &= h_{N_1}^{-1} \circ f_{N_1} \circ h_{N_1} = \frac{1}{f_{N_1}(x_2)} \tan^2(N_1 \arctan^P \sqrt{x_1}) \\ X_{N_2}(x_2; f_{N_1}(x_1)) &= h_{N_2}^{-1} \circ f_{N_2} \circ h_{N_2} = \frac{1}{f_{N_2}(x_1)} \tan^2(N_2 \arctan^P \sqrt{x_2}); \end{aligned} \quad (\text{III-6})$$

Below we give some of coupled map as an illustration

$$\begin{aligned} X_1 &= \frac{f_1(x_2)(2x_1 - 1)^2}{4x_1(1 - x_1) + f_1(x_2)(2x_1 - 1)^2} \\ X_2 &= \frac{f_2(x_1)(2x_2 - 1)^2}{4x_2(1 - x_2) + f_2(x_1)(2x_2 - 1)^2} \end{aligned} \quad (\text{III-7})$$

and

$$\begin{aligned} X_1 &= \frac{f_1(x_2)(2x_1 - 1)^2}{4x_1(1 - x_1) + f_1(x_2)(2x_1 - 1)^2} \\ X_2 &= \frac{f_2(x_1)(4x_2 - 3)^2}{f_2(x_1)(4x_2 - 3)^2 + (1 - x_2)(4x_2 - 1)^2} \end{aligned} \quad (\text{III-8})$$

with

$$T_1(x_2) = \frac{2(x_2)}{1 + (x_2)} \sqrt{\frac{(x_2)}{(x_2)}};$$

$$S(x) = \left(\frac{1}{2} + gx \right)^2;$$

One can show the map $T_{2;3}$, is ergodic, for $0 < x_1 < 2$ and $\frac{1}{3} < x_2 < 3$, while for $2 < x_1 < 1$ and $0 < x_2 < \frac{1}{3}$ ($3 < x_2 < 1$) it has fixed point at $[0;0]$ ($[0;1]$) (see Fig. 5).

$$T_{3;2} = \begin{cases} X_1 = \frac{\frac{2}{1}(x_2)x_1(4x_1-3)^2}{\frac{2}{1}(x_2)x_1(4x_1-3)^2 + (1-x_1)(4x_1-1)^2} \\ X_2 = \frac{\frac{2}{2}(2x_2-1)^2}{4x_2(1-x_2) + \frac{2}{2}(2x_2-1)^2} \end{cases} \quad (\text{III-9})$$

Similarly, one can show that the map $T_{3;2}$, is ergodic for $\frac{1}{3} < x_1 < 3$ and $0 < x_2 < 2$, while for $0 < x_1 < \frac{1}{3}$ ($3 < x_1 < 1$) and $0 < x_2 < 2$ ($2 < x_2 < 1$) it has fixed point at $[0;0]$ ($[1;0]$) (see Fig. 6).

$$T_{3;3} = \begin{cases} X_1 = \frac{\frac{2}{1}(x_2)x_1(4x_1-3)^2}{\frac{2}{1}(x_2)x_1(4x_1-3)^2 + (1-x_1)(4x_1-1)^2} \\ X_2 = \frac{\frac{2}{2}x_2(4x_2-3)^2}{\frac{2}{2}x_2(4x_2-3)^2 + (1-x_2)(4x_2-1)^2} \end{cases} \quad (\text{III-10})$$

with

$$T_1(x_2) = \frac{3(x_2) + 1}{3 + (x_2)} \sqrt{\frac{(x_2)}{(x_2)}};$$

$$S(x) = \left(\frac{1}{3} + gx \right)^2;$$

Also the map $T_{3;3}$, is ergodic for $\frac{1}{3} < x_1 < 3$ and $\frac{1}{3} < x_2 < 3$, while for $0 < x_1 < \frac{1}{3}$ ($3 < x_1 < 1$) and $0 < x_2 < \frac{1}{3}$ ($3 < x_2 < 1$) it has fixed point at $[0;0]$ ($[1;1]$) (see Fig. 7).

$$T_{4;3} = \begin{cases} X_1 = \frac{\frac{2}{1}(x_1)(1-8x_1(1-x_1))^2}{\frac{2}{1}(x_1)(1-8x_1(1-x_1))^2 + 16x_1(1-x_1)(1-2x_2)^2} \\ X_2 = \frac{\frac{2}{2}x_2(4x_2-3)^2}{\frac{2}{2}x_2(4x_2-3)^2 + (1-x_2)(4x_2-1)^2} \end{cases} \quad (\text{III-11})$$

and

$$\begin{aligned} X_1 &= \frac{x_1^2 (1 - 8x_1(1 - x_1))^2}{x_1^2 (1 - 8x_1(1 - x_1))^2 + 16x_1(1 - x_1)(1 - 2x_2)^2} \\ X_2 &= \frac{x_2^2 (1 - 8x_2(1 - x_2))^2}{x_2^2 (1 - 8x_2(1 - x_2))^2 + 16x_2(1 - x_2)(1 - 2x_2)^2} \end{aligned} \quad (\text{III-12})$$

with

$$\begin{aligned} x_1(x_2) &= \frac{x_2^2(x_2) + 6x_2(x_2) + 1}{(x_2)(4 - (x_2) + 4)}; \\ x(x) &= \frac{4 - 6x_1 - 4x_1^2 + 2x_1^2}{2(1 - 4)} + gx^2: \end{aligned}$$

The maps $x_{2,2}$, $x_{4,3}$ and $x_{4,4}$ are also either ergodic or they only single fixed points at $[0;0]$ or $[0;1]$.

IV Invariant measure

Dynamical systems, even apparently simple dynamical systems which are described by maps of an interval can display a rich variety of different asymptotic behavior. On measure theoretical level these type of behavior are described by SRB [13, 14] or invariant measure describing statistically stationary states of the system. The probability measure on $[0;1]$ is called an SRB or invariant measure of the map $y = x_N(x; \cdot)$ given in (II-1), if it is $x_N(x; \cdot)$ -invariant and absolutely continuous with respect to Lebesgue measure. For deterministic system such as $x_N(x; \cdot)$ -map, the $x_N(x; \cdot)$ -invariance means that, its invariant measure $\mu(x)$ fulfills the following formal FP integral equation

$$\mu(y) = \int_0^{x_1} \mu(x_N(x; \cdot)) \mu(x) dx:$$

This is equivalent to:

$$\mu(y) = \int_{x_2}^x \mu(x) \frac{dx}{dy}; \quad (\text{IV-1})$$

defining the action of standard FP operator for the map $T_N(x; \mu)$ over a function as:

$$P_N f(y) = \int_{T_N^{-1}(y)} f(x) \frac{dx}{dy} : \quad (IV-2)$$

We see that, the invariant measure $\rho(x)$ is actually the eigenstate of the FP operator P_N corresponding to largest eigenvalue 1.

As it is proved in Appendix A, the invariant measure $\rho_N(x; \mu)$ has the following form :

$$\rho(x) = \frac{1}{\pi} \frac{\mu}{x(1-x)(1+\mu x)} \quad (IV-3)$$

with $\mu > 0$, is the invariant measure of the maps $T_N(x; \mu)$ provided that, we choose the parameter μ in the following form :

$$\mu = \frac{P_{\lfloor \frac{N-1}{2} \rfloor} C_{2k+1}^N}{P_{\lfloor \frac{N}{2} \rfloor} C_{2k}^N} ; \quad (IV-4)$$

where the symbol $\lfloor \cdot \rfloor$ means greatest integer part.

As we see, the above measure is defined only for $\mu > 0$, hence from the relations (IV-3), it follows that the maps $T_N(x; \mu)$ have invariant measure only for $\mu \geq (\frac{1}{N}; N)$ for odd value of N , and for even value $\mu \geq (N; 1)$. For other values of μ these maps have single attractive fixed points, which is the same as the prediction of the previous section.

Similarly the probability measure for coupled chaotic maps T_{N_1, N_2} given in (III-4) fulfills the following form of FP integral equation

$$\rho(y_1; y_2) = \int_0^1 dx_1 \int_0^1 dx_2 \rho(x_1; x_2) T_{N_1}(x_1; y_1) T_{N_2}(x_2; y_2) ;$$

which is equivalent to:

$$\rho(y_1; y_2) = \int_{x \in T_{N_1, N_2}^{-1}(y_1, y_2)} J(x_1; x_2) \rho(x_1; x_2) ; \quad (IV-5)$$

where $J(x_1; x_2)$ is the Jacobian of the transformation $N_1; N_2$ which is equal to

$$J(x_1; x_2) = \frac{\partial(x_1; x_2)}{\partial(y_1; y_2)};$$

again defining the action of standard FP operator for the map $N_1; N_2$ over a function as:

$$P_{N_1; N_2} f(y) = \int_{N_1; N_2} f(x) J(x_1; x_2) dx \quad (IV-6)$$

We see that, the invariant measure $(x_1; x_2)$ is also the eigenstate of the FP operator $P_{N_1; N_2}$ corresponding to largest eigenvalue 1.

As it is proved in Appendix B the invariant measure $N_1; N_2(x_1; x_2)$ has the following form :

$$\frac{1}{\int_{x_2(1-x_2)(\alpha_2 + (1-x_2)x_2)} P_{\alpha_2}} = \frac{1}{\int_{x_1(1-x_1)(\alpha_1(x_2) + (1-\alpha_1(x_2))x_1)} \alpha_1(x_2)}; \quad (IV-7)$$

with $\alpha_2 > 0$ and $\alpha_1(x) > 0$ given in (III-1).

V Kolmogorov-Sinai entropy

KS-entropy or metric entropy [13] measures how chaotic a dynamical system is and it is proportional to the rate at which information about the state of dynamical system is lost in the course of time or iteration. Therefore, it can also be defined as the average rate of information loss for a discrete measurable dynamical system $(N(x);)$. By introducing a partition $\mathcal{A} = \{A_i\}$ of the interval $[0;1]$ into individual laps A_i one can define the usual entropy associated with the partition by:

$$H(\mathcal{A}; N) = - \sum_{i=1}^K p_i(\mathcal{A}) \ln p_i(\mathcal{A});$$

where $m(A_i) = \int_{A_i} \rho(x) dx$ is the invariant measure of A_i . By defining n -th refining (n) of A_i as:

$$A_i^{(n)} = \bigcup_{k=0}^{n-1} (A_i + k \cdot \Delta x)$$

then entropy per unit step of refining is defined by :

$$h(\rho; A_i^{(n)}) = \lim_{n \rightarrow \infty} \frac{1}{n} H(\rho; A_i^{(n)})$$

Now, if the size of individual laps of $(A_i^{(n)})$ tends to zero as n increases, then the above entropy is known as K S-entropy, that is:

$$h(\rho; A_i^{(n)}) = h(\rho; \Delta x; A_i):$$

K S-entropy, which is a quantitative measure of the rate of information loss with the refining, may also be written as:

$$h(\rho; \Delta x; A_i) = - \int_{A_i} \rho(x) dx \ln \left| \frac{d}{dx} f^{(n)}(x) \right| \quad (V-1)$$

which is also a statistical mechanical expression for the Lyapunov characteristic exponent, that is, the mean divergence rate of two nearby orbits. The measurable dynamical system $(f; \rho)$ is chaotic for $h > 0$ and predictive for $h = 0$. As it is proved in Appendix C, the $h(\rho; \Delta x; A_i)$ has the following form

$$h(\rho; \Delta x; A_i) = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \left| \frac{f^{(n)}(x_1) - f^{(n)}(x_2)}{x_1 - x_2} \right| \quad (V-2)$$

Similarly the K S-entropy of coupled chaotic maps can be written as

$$h(\rho; \Delta x_1, \Delta x_2) = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \left| \frac{f^{(n)}(x_1, x_2) - f^{(n)}(x_1', x_2')}{(x_1 - x_1')(x_2 - x_2')} \right| \quad (V-3)$$

In Appendix D we have calculated K S-entropy of coupled chaotic maps (f_1, f_2) for $N_1 = 2, 3, 4$ and 5, where below we give the results.

$$\begin{aligned}
h(\cdot; {}_2N_2) = & \ln \frac{N_2(1 + \frac{p}{2} + 2^{\frac{p}{2}})^{N_2-1}}{\left(\prod_{k=0}^{\lfloor \frac{N_2}{2} \rfloor} C_{2k}^{N_2-k} \right) \left(\prod_{k=0}^{\lfloor \frac{N_2-1}{2} \rfloor} C_{2k+1}^{N_2-k} \right)} \\
& + \ln \frac{r \frac{q}{1 + \frac{q}{1_0} + \frac{g}{2}}}{q \frac{p}{1 + \frac{q}{1_0} + \frac{g}{2}} + \frac{r \frac{p}{2g \frac{1_0}{2} + \frac{2g}{2}}}{q \frac{p}{1 + \frac{q}{1_0} + \frac{g}{2}}}}; \quad (V-4)
\end{aligned}$$

$$\begin{aligned}
h(\cdot; {}_3N_2) = & \ln \frac{N_2(1 + \frac{p}{2} + 2^{\frac{p}{2}})^{N_2-1}}{\left(\prod_{k=0}^{\lfloor \frac{N_2}{2} \rfloor} C_{2k}^{N_2-k} \right) \left(\prod_{k=0}^{\lfloor \frac{N_2-1}{2} \rfloor} C_{2k+1}^{N_2-k} \right)} \\
& + \ln 3 \left(1 + \frac{r \frac{q}{1_0} + \frac{s \frac{g}{2}}{2}}{2} \right)^8 = \\
& \frac{B}{C} \left(\frac{q}{3 + \frac{q}{1_0} + \frac{g}{2}} + \frac{\frac{v}{u} \frac{q}{2g \frac{1_0}{2} + \frac{2g}{2}}}{(3 + \frac{q}{1_0})} \right)^{1/2} \\
& \frac{B}{C} \left(\frac{q}{1 + 3 \frac{q}{1_0} + \frac{p}{2}} + \frac{\frac{v}{u} \frac{q}{6g \frac{1_0}{2} + \frac{2g}{2}}}{3(1 + 3 \frac{q}{1_0})} \right)^{1/2^3} \frac{7}{5}; \quad (V-5)
\end{aligned}$$

$$\begin{aligned}
h(\cdot; {}_4N_2) = & \ln \frac{N_2(1 + \frac{p}{2} + 2^{\frac{p}{2}})^{N_2-1}}{\left(\prod_{k=0}^{\lfloor \frac{N_2}{2} \rfloor} C_{2k}^{N_2-k} \right) \left(\prod_{k=0}^{\lfloor \frac{N_2-1}{2} \rfloor} C_{2k+1}^{N_2-k} \right)} \\
& + \ln 4 \left(1 + \frac{r \frac{q}{1_0} + \frac{s \frac{g}{2}}{2}}{2} \right)^{12} = \\
& \frac{B}{C} \left(\frac{q}{3 + 2 \frac{p}{2} + \frac{q}{1_0} + \frac{g}{2}} + \frac{\frac{v}{u} \frac{q}{2g \frac{1_0}{2} + \frac{2g}{2}}}{(3 + 2 \frac{p}{2} + \frac{q}{1_0})} \right)^{1/2} \\
& \frac{B}{C} \left(\frac{q}{3 + 2 \frac{p}{2} + \frac{q}{1_0} + \frac{g}{2}} + \frac{\frac{v}{u} \frac{q}{2g \frac{1_0}{2} + \frac{2g}{2}}}{(3 + 2 \frac{p}{2} + \frac{q}{1_0})} \right)^{1/2} \\
& \frac{B}{C} \left(\frac{q}{4 + 4 \frac{q}{1_0} + \frac{4g}{2}} + \frac{\frac{v}{u} \frac{q}{8g \frac{1_0}{2} + \frac{2g}{2}}}{(4 + 4 \frac{q}{1_0})} \right)^{1/2^3} \frac{7}{5}; \quad (V-6)
\end{aligned}$$

and

$$\begin{aligned}
h(\cdot; \mathcal{M}_2) &= \ln \frac{N_2 (1 + \frac{1}{2} + 2^{\frac{1}{2}})^{N_2 - 1}}{\left(\prod_{k=0}^{\lfloor \frac{N_2}{2} \rfloor} C_{2k}^{N_2 - k} \right) \left(\prod_{k=0}^{\lfloor \frac{N_2 - 1}{2} \rfloor} C_{2k+1}^{N_2 - k} \right)} \\
&\quad + \ln \left(1 + \frac{1}{10} + \frac{g}{2} \right)^{16} = \\
&\quad \ln \left(\frac{1}{5} + \frac{1}{2} + \frac{1}{10} + \frac{g}{2} + \frac{\sqrt{2g}}{2} + \frac{2g}{2} \left(\frac{1}{5} + \frac{1}{2} + \frac{1}{10} \right) \right) \\
&\quad \ln \left(\frac{1}{5} + \frac{1}{2} + \frac{1}{10} + \frac{g}{2} + \frac{\sqrt{2g}}{2} + \frac{2g}{2} \left(\frac{1}{5} + \frac{1}{2} + \frac{1}{10} \right) \right) \\
&\quad \ln \left(1 + \frac{2}{5} + \frac{1}{10} + \frac{g}{2} + \frac{\sqrt{2g}}{2} + \frac{2g}{2} \left(1 + \frac{2}{5} + \frac{1}{10} \right) \right) \\
&\quad \ln \left(1 + \frac{2}{5} + \frac{1}{10} + \frac{g}{2} + \frac{\sqrt{2g}}{2} + \frac{2g}{2} \left(1 + \frac{2}{5} + \frac{1}{10} \right) \right)^3 :
\end{aligned}
\tag{V-7}$$

VI Simulation

Here in this section we try to calculate Lyapunov characteristic exponent of maps (II 3;4) and (III 6;7) in order to investigate these maps numerically. In fact, Lyapunov characteristic exponent is the characteristic exponent of the rate of average magnification of the neighborhood of an arbitrary point $\mathbf{x}_0 = (x_{10}; x_{20})$ and it is denoted by $\lambda(\mathbf{x}_0)$ which is written as [14]:

$$\begin{aligned}
\lambda(\mathbf{x}_0) &= \lim_{n \rightarrow \infty} \frac{1}{n} \ln \left| \frac{\partial \mathbf{x}_n}{\partial \mathbf{x}_0} \right| \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} \ln \left| \frac{\partial X_1(\mathbf{x}_k; 2)}{\partial x_{10}} : \frac{\partial X_2(\mathbf{x}_k; 1)}{\partial x_{20}} \right|
\end{aligned}
\tag{VI-1}$$

where $x_k = \frac{z}{\dots} \Big| \frac{1}{\dots} \Big\{ \dots \Big\}$. It is obvious that $(\lambda_0) < 0$ for an attractor, $(\lambda_0) > 0$ for a repeller and $(\lambda_0) = 0$ for marginal situation. Also the Liapunov number is independent of initial point x_0 , provided that the motion inside the invariant manifold is ergodic, thus (λ_0) characterizes the invariant manifold of as a whole. For the values of parameter , such that the map be measurable, Birkhoff ergodic [15] theorem implies the equality of KS-entropy and Liapunov number, that is:

$$h(\lambda; \mu) = (\lambda_0; \mu); \quad (VI-2)$$

Comparison of analytically calculated KS-entropy of maps (see Figures 8-11) with the corresponding Lyapunov characteristic exponent obtained by the simulation, indicates that in chaotic region, these maps are ergodic as Birkhoff ergodic theorem predicts. In non chaotic region of the parameter, Lyapunov characteristic exponent is negative definite, since in this region we have only single period fixed points without bifurcation. Therefore, combining the analytic discussion of section V with the numerical simulation we deduce that, these maps are ergodic in certain values of their parameters as explained above and in complementary interval of parameters they have only a single period one attractive fixed point, such that in contrary to the most of usual one-dimensional one-parameter family of maps they have only bifurcation from a period one attractive fixed point to chaotic state or vice versa. A comparison of these results with those of obtained in section V, verified the accuracy of implemented computation methods.

V II Conclusion

We have given hierarchy of exactly solvable one-parameter families of one-dimensional chaotic maps with an invariant measure, that is measurable dynamical system with an interesting property of being either chaotic (proper to say ergodic) or having stable period one fixed point and they bifurcate from a stable single periodic state to chaotic one and vice-versa without having usual period doubling or period- n -tupling scenario. Also by coupling these maps in an appropriate method, we have obtained some coupled chaotic maps with invariant measure, where they also bifurcate without period- n -tupling scenario to chaos. Since the invariant measure plays an important role in analytic calculation of important equations such as K S-entropy, therefore it would be interesting to find the other measurable one parameter maps, many parameters, coupled and higher dimensional maps with invariant measure, which are under investigation.

Appendix A : Derivation of measure of one-parameter families of chaotic maps

In order to prove that measure (IV-3) satisfies equation (IV-1), it is rather convenient to consider the conjugate map.

$$\tilde{\mu}_N(x; y) = \frac{1}{2} \tan^2 \left(N \arctan \sqrt{\frac{y}{1+y}} \right) \quad (\text{A-1})$$

with measure $\tilde{\mu}_N$ related to the measure μ_N with the following relation:

$$\tilde{\mu}_N(x) = \frac{1}{(1+x)^2} \mu_N\left(\frac{1}{1+x}\right);$$

Denoting $\tilde{\mu}_N(x; y)$ on the left hand side of (A-1) by y and inverting it, we get :

$$x_k = \tan^2 \left(\frac{1}{N} \arctan \sqrt{\frac{y}{1+y}} + \frac{k}{N} \right) \quad k = 1, \dots, N : \quad (\text{A-2})$$

Then, taking derivative of x_k with respect to y , we obtain:

$$y \frac{dx_k}{dy} = \frac{1}{N} \sqrt{\frac{y}{1+y}} \frac{1}{\sqrt{1+y}} : \quad (\text{A-3})$$

Substituting the above result in equation (IV-1), we get:

$$\tilde{\mu}_N(y) \sqrt{\frac{y}{1+y}} = \frac{1}{N} \sum_{k=1}^N \sqrt{\frac{y}{1+y}} \tilde{\mu}_N(x_k) ; \quad (\text{A-4})$$

Now, by considering the following ansatz for the invariant measure $\tilde{\mu}_N(y)$:

$$\tilde{\mu}_N(y) = \frac{1}{\sqrt{1+y}} ; \quad (\text{A-5})$$

the above equation reduces to:

$$\frac{1 + \frac{y}{1+y}}{1 + \frac{y}{1+y}} = \frac{1}{N} \sum_{k=1}^N \frac{1 + x_k}{1 + x_k}$$

which can be written as:

$$\frac{1 + \frac{y}{1+y}}{1 + \frac{y}{1+y}} = - + \frac{1}{2} \frac{d}{dy} \left(\ln \left(\prod_{k=1}^N (1 + x_k) \right) \right) : \quad (\text{A-6})$$

To evaluate the second term in the right hand side of above formulas we can write the equation in the following form :

$$\begin{aligned}
 0 &= {}^2y \cos^2(N \arctan \sqrt[N]{x}) - \sin^2(N \arctan \sqrt[N]{x}) \\
 &= \frac{(1)^N}{(1+x)^N} \sum_{k=0}^{\lfloor \frac{N-1}{2} \rfloor} {}^2y \left(C_{2k}^N (1)^N x^k \right)^2 - x \sum_{k=0}^{\lfloor \frac{N-1}{2} \rfloor} C_{2k+1}^N (1)^N x^k \left(\frac{1}{A} \right)^N ; \\
 &= \frac{\text{constant}}{(1+x)^N} \prod_{k=1}^N (x - x_k) ;
 \end{aligned}$$

where x_k are the roots of equation (A-1) and they are given by formula (A-2). Therefore, we have:

$$\begin{aligned}
 & \frac{\partial}{\partial x} \ln \prod_{k=1}^N (1 + x_k) \\
 &= \frac{\partial}{\partial x} \ln \left((1)^N \left({}^2y \cos^2(N \arctan \sqrt[N]{x}) - \sin^2(N \arctan \sqrt[N]{x}) \right) \right) \\
 &= \frac{N}{1+x} + \frac{N(1+{}^2y)A(\frac{1}{x})}{(A(\frac{1}{x}))^2 {}^2y + (B(\frac{1}{x}))^2} ; \quad (A-7)
 \end{aligned}$$

In deriving of above formulas we have used the following identities

$$\begin{aligned}
 \cos(N \arctan \sqrt[N]{x}) &= \frac{A(\frac{1}{x})}{(1+x)^{\frac{N}{2}}} ; \\
 \sin(N \arctan \sqrt[N]{x}) &= \sqrt[N]{x} \frac{B(\frac{1}{x})}{(1+x)^{\frac{N}{2}}} ; \quad (A-8)
 \end{aligned}$$

inserting the results (A-7) in (A-6), we get:

$$\frac{1+{}^2y}{1+y} = \frac{1+{}^2y}{\frac{B(\frac{1}{x})}{A(\frac{1}{x})} + (\frac{A(\frac{1}{x})}{B(\frac{1}{x})})y} ;$$

Hence to get the final result we have to choose the parameter as:

$$= \frac{B(\frac{1}{x})}{A(\frac{1}{x})} ;$$

Appendix B : Derivation of measure of coupled chaotic maps.

In order to prove that measure (IV 7) satisfies equation (IV 5), it is rather convenient to consider the conjugate map.

Now, denoting X_1 by y_1 and X_2 by y_2 and inverting the conjugate map \sim_{N_1, N_2}

we get:

$$\begin{aligned} x_{k_1 k_2} &= \tan^2 \left(\frac{1}{N_1} \arctan \sqrt{\frac{g_1(x_2)}{y_1^2}} + \frac{k_1}{N_1} \right) \\ x_{k_2} &= \tan^2 \left(\frac{1}{N_2} \arctan \sqrt{\frac{g_2(x_2)}{y_2^2}} + \frac{k_2}{N_2} \right); \end{aligned} \quad (B-1)$$

then taking the derivative of (B 1) with respect to y_1, y_2 :

$$\begin{aligned} \frac{\partial x_{k_1 k_2}}{\partial y_1} &= \frac{g_1(x_{k_2})}{N_1} \cdot \frac{\sqrt{\frac{g_1(x_{k_2})}{y_1^2}} (1 + x_{k_1 k_2})}{y_1 (1 + g_1^2(x_{k_2}) y_1)} \\ \frac{\partial x_{k_2}}{\partial y_2} &= \frac{g_2(x_{k_2})}{N_2} \cdot \frac{(1 + x_{k_2})}{(1 + \frac{g_2^2(x_{k_2})}{y_2^2})} \cdot \frac{\sqrt{\frac{g_2(x_{k_2})}{y_2^2}}}{y_2}; \end{aligned} \quad (B-2)$$

and substituting the above results in equation (IV 1), we obtain:

$$\begin{aligned} \sim(y_1; y_2) &= \sum_{k_1, k_2} \frac{\partial x_{k_2}}{\partial y_2} \cdot \frac{\partial x_{k_1 k_2}}{\partial y_1} \cdot \sim(x_{k_1 k_2}; x_{k_2}) \\ &= \sum_{k_1, k_2} \frac{g_1(x_{k_2})}{N_1} \cdot \frac{\sqrt{\frac{g_1(x_{k_2})}{y_1^2}} (1 + x_{k_1 k_2})}{y_1 (1 + g_1^2(x_{k_2}) y_1)} \cdot \frac{g_2(x_{k_2})}{N_2} \cdot \frac{(1 + x_{k_2})}{(1 + \frac{g_2^2(x_{k_2})}{y_2^2})} \cdot \frac{\sqrt{\frac{g_2(x_{k_2})}{y_2^2}}}{y_2} \cdot \sim(x_{k_1 k_2}; x_{k_2}); \end{aligned} \quad (B-3)$$

Now, considering the following ansatz for invariant measure \sim_{N_1, N_2}

$$\sim(x_1; x_2) = \frac{1}{N_1} \cdot \frac{g_1(x_2)}{(1 + \frac{g_1^2(x_2)}{x_1^2})}; \quad (B-4)$$

we come to the following equation:

$$\sim(y_1; y_2) = \frac{1}{N_1} \sum_{k_2} \frac{\partial x_{k_2}}{\partial y_2} \cdot \frac{1}{y_1} \cdot \frac{g_1(x_{k_2})}{(1 + \frac{g_1^2(x_{k_2})}{y_1^2})} \cdot \sum_{k_1} \frac{g_1(x_{k_2})}{N_1} \cdot \frac{(1 + x_{k_1 k_2})}{(1 + \frac{g_1^2(x_{k_2})}{y_1^2})}; \quad (B-5)$$

using the relation (III 3) in the last part of above relation, it reads

$$\begin{aligned} \prod_{k_1} \frac{g_1(x_{k_2})}{N_1} \cdot \frac{(1 + \prod_{k_1} x_{k_1 k_2})}{(1 + \prod_{k_1} (x_{k_2}) x_{k_1 k_2})} &= \frac{g_1(x_{k_2}) A_{N_1}(\prod_{k_1} x_{k_2}) (1 + g_1^2(x_{k_2}) y_1)}{B_{N_1}(\prod_{k_1} x_{k_2})} \\ &\frac{1}{(1 + \prod_{k_1} (x_{k_2}) (\frac{g_1(x_{k_2}) A_{N_1}(\prod_{k_1} x_{k_2})}{B_{N_1}(\prod_{k_1} x_{k_2})}))}; \end{aligned}$$

therefore, the relation (B-5) reduces to:

$$\begin{aligned} \sim(y_1; y_2) &= \prod_{k_2} \frac{(x_{k_2})}{P_{Y_1}} \cdot \frac{g_1(x_{k_2}) A_{N_1}(\prod_{k_1} x_{k_2})}{B_{N_1}(\prod_{k_1} x_{k_2})} \\ &\frac{1}{(1 + \prod_{k_1} (x_{k_2}) (\frac{g_1(x_{k_2}) A_{N_1}(\prod_{k_1} x_{k_2})}{B_{N_1}(\prod_{k_1} x_{k_2})})^2)} \cdot \frac{\partial x_{k_2}}{\partial y_2} \frac{C}{A}; \end{aligned} \quad (B-6)$$

On the otherhand, we know that the measure $\sim(y_1; y_2)$ is

$$\sim(y_1; y_2) = \frac{1}{(1 + \frac{(y_2)}{(y_2) y_1})} \frac{(y_2)}{P_{Y_1}}; \quad (B-7)$$

which is possible if $\prod_{k_1} (x_{k_2})$ and $\prod_{k_1} (y_2)$ to be related with

$$\prod_{k_1} (y_2) = \frac{g_1(x_{k_2}) A_{N_1}(\prod_{k_1} x_{k_2})}{B_{N_1}(\prod_{k_1} x_{k_2})} \prod_{k_1} (x_{k_2}) \quad (B-8)$$

and $\prod_{k_1} (y_2)$ can be given in terms of (x_k) as:

$$\prod_{k_2} (y_2) = \prod_{k_2} \frac{(x_{k_2})}{\prod_{k_1} (x_{k_2})} \cdot \frac{\partial x_{k_2}}{\partial y_2} \frac{C}{A}(x_{k_2});$$

or it can be written as

$$\frac{(y_2)}{\prod_{k_1} (y_2)} = \prod_{k_2} \frac{\partial x_{k_2}}{\partial y_2} \cdot \frac{(x_{k_2})}{\prod_{k_1} (x_{k_2})};$$

Therefore, $\prod_{k_1} \frac{(y_2)}{(y_2)}$ satisfies the PF equation of map N , hence it is proportional to its invariant measure, that is

$$\prod_{k_1} \frac{(y_2)}{(y_2)} = \frac{1}{(1 + \frac{P_2}{2 y_2})}; \quad (B-9)$$

Appendix C : Derivation of entropy of one-parameter families of chaotic maps

In order to calculate the KS-entropy of the maps $T_N(x; \lambda)$, it is rather convenient to consider their conjugate maps given by (II-2), since it can be shown that KS-entropy is a kind of topological invariant, that is, it is preserved under conjugacy map. Hence we have:

$$h(\lambda; T_N) = h(\tilde{\lambda}; \tilde{T}_N):$$

The integral (V-1) can be written as

$$h(\lambda; T_N(x; \lambda)) = \int_0^1 \tilde{\rho}(x) dx \ln \left| \frac{d}{dx} \left(\frac{1}{2} \tan^2 (N \arctan \sqrt{\frac{p}{x}}) \right) \right|$$

we have

$$h(\lambda; T_N(x; \lambda)) = \int_0^1 \tilde{\rho}(x) dx \ln \left| \frac{N}{2} \frac{1}{\sqrt{x}(1+x)} \frac{\sin N (\arctan \sqrt{\frac{p}{x}})}{\cos^3 N (\arctan \sqrt{\frac{p}{x}})} \right|;$$

using the relation (A-8), we get

$$h(\lambda; T_N(x; \lambda)) = \frac{1}{2} \int_0^1 \frac{p dx}{\sqrt{x}(1+x)} \ln \left| \frac{N}{2} \frac{(1+x)^{N-1} B(x)}{(A(x))^3} \right|; \quad (C-1)$$

We see that polynomials appearing in the numerator (denominator) of integrand appearing on the right hand side of equation (C-1), have $\frac{[N-1]}{2}$ ($\frac{[N-1]}{2}$) simple roots, denoted by x_k^B ($k = 1, \dots, \frac{[N-1]}{2}$) (x_k^A ($k = 1, \dots, \frac{[N-1]}{2}$)) in the interval $[0, 1]$. Hence, we can write the above formula in the following form :

$$h(\lambda; T_N(x; \lambda)) = \frac{1}{2} \int_0^1 \frac{p dx}{\sqrt{x}(1+x)} \ln \left| \frac{N}{2} \frac{(1+x)^{N-1} \prod_{k=1}^{\frac{[N-1]}{2}} (x - x_k^B)}{\prod_{k=1}^{\frac{[N-1]}{2}} (x - x_k^A)} \right|;$$

Now making the following change of variable $x = \frac{1}{2} \tan^2 \frac{\theta}{2}$, and taking into account that degree of numerators and denominator are equal for both even and odd values of N , we get

$$h(\boldsymbol{\mu}; \mathbf{N}(\mathbf{x}; \boldsymbol{\mu})) = -\frac{1}{2} \sum_{j=1}^N \left[\ln \left(\frac{N}{2} \right) + (N-1) \ln |j| + 1 + (-1)^j \cos j \right] \\ + \sum_{k=1}^{\lfloor \frac{N-1}{2} \rfloor} \ln |1 - x_k^B| + (1 + x_k^B) \cos j - 3 \sum_{k=1}^{\lfloor \frac{N-1}{2} \rfloor} \ln |1 - x_k^A| + (1 + x_k^A) \cos j : \quad (C-1)$$

Using the following integrals:

$$\frac{1}{2} \int_0^{\pi} \ln |a + b \cos \theta| d\theta = \begin{cases} \ln \left(\frac{a + \sqrt{a^2 - b^2}}{2} \right) & \text{if } |a| \geq |b| \\ \ln \left(\frac{|b|}{2} \right) & \text{if } |a| < |b| \end{cases}$$

we get

$$h(\boldsymbol{\mu}; \mathbf{N}(\mathbf{x}; \boldsymbol{\mu})) = \ln \frac{1}{2} \frac{N (1 + 2^{p-N-1})}{\left(\prod_{k=0}^{\lfloor \frac{N-1}{2} \rfloor} C_{2k}^N \right) \left(\prod_{k=0}^{\lfloor \frac{N-1}{2} \rfloor} C_{2k+1}^N \right)} : \quad (C-2)$$

Appendix D : Derivation of entropy of coupled chaotic maps.

The K S-entropy of $_{2N_2}$ given in (V 3) can be written as

$$h(\sim; \sim_{2N_2}) = \int_0^1 \sim(x_1) dx_1 \int_0^1 \sim(x_2) dx_2 \left(\ln \int \frac{\partial X_2}{\partial x_1} j + \ln \int \frac{\partial X_{N_2}}{\partial x_2} j \right) : \quad (D-1)$$

Now, using the formulas (III 5) the second integral reads

$$= \int_0^1 \sim(x_1) dx_1 \int_0^1 \sim(x_2) dx_2 \ln \int \frac{\partial}{\partial x_1} \left(\frac{1}{2} : \tan^2 (N_2 \arctan \sqrt{x_2}) \right) j$$

or

$$= \int_0^1 \sim(x_1) dx_1 \int_0^1 \sim(x_2) dx_2 \ln \int \frac{N_2}{2} \frac{1}{\sqrt{x_2} (1+x_2)} \frac{\sin N_2 (\arctan \sqrt{x_2})}{\cos^3 N_2 (\arctan \sqrt{x_2})} j ;$$

where using the relation (A 8), we get

$$= \frac{1}{2} \int_0^1 \frac{\sqrt{x_2} dx_2}{\sqrt{x_2} (1+x_2)} \ln \int \frac{N_2}{2} \frac{(1+x_2)^{N_2-1} B(\sqrt{x_2})}{x_2^2 (A(\sqrt{x_2}))^3} j \quad (D-2)$$

Finally, calculating the above integral with the prescription of Appendix C,

we obtain:

$$= \ln \frac{N_2 (1 + \frac{1}{2} + 2^{\frac{1}{2}})^{N_2-1} \frac{C}{A}}{\left(\sum_{k=0}^{\lfloor \frac{N_2}{2} \rfloor} C_{2k}^{N_2-k} \right) \left(\sum_{k=0}^{\lfloor \frac{N_2-1}{2} \rfloor} C_{2k+1}^{N_2-k} \right)} ; \quad (D-3)$$

Now, using the relation (II 3) together with the invariance of measure, the

first integral in (D 1) can be

$$\begin{aligned} &= \frac{1}{2} \int_0^1 \sim(x_1) dx_1 \int_0^1 \sim(x_2) dx_2 \ln \int \frac{\partial}{\partial x_1} \frac{(x_2)}{(X_2)} \frac{A_2^2 \left(\frac{1}{(x_2)} \right)}{B_2^2 \left(\frac{1}{(x_2)} \right)} \\ &\quad \frac{\sin^2 (2 \arctan \sqrt{x_1})}{\cos^2 (2 \arctan \sqrt{x_1})} j \\ &= \frac{1}{2} \int_0^1 \sim(x_1) dx_1 \int_0^1 \sim(x_2) dx_2 \ln \int \frac{(x_2)}{(X_2)} \frac{(1 + \sqrt{x_2})^2}{A_2(\sqrt{x_2}) B_2(\sqrt{x_2})} j \quad (D-4) \end{aligned}$$

Due to the invariance of measure the integral of the expression $\frac{(x_2)}{(X_2)}$ vanishes, since according to Appendix E, we have

$$\int_0^1 dx (x) \ln ((X_2)) = \int_0^1 dx (x) \ln (((x))) = \int_0^1 dx (x) \ln ((x_2)) \quad (D-5)$$

Making a change of variable $x = \frac{1}{2} \tan^2 \frac{\theta}{2}$ in the first integral and omitting the expression $\frac{(x_2)}{(x_2)}$, it reduces to

$$= \frac{2}{\pi} \int_0^{\frac{\pi}{2}} d\theta \left(\ln \left(\left(\frac{1}{2} (1 + \cos \theta) \right) + g \right) + \left(\frac{1}{2} (1 + \cos \theta) + g \right) \cos \theta \right)$$

$$\ln \left(\frac{1}{2} + \frac{1}{2} \cos \theta \right) + \ln \left(\frac{3}{4} + \cos \theta + \frac{1}{4} \cos 2\theta \right) + \ln (A + B \cos \theta + C \cos 2\theta)$$

Which

$$\begin{aligned} A &= \frac{1}{8} \left(3(1 + \cos \theta) + \frac{2g}{2} \cos \theta + 3 \left(\frac{g}{2} \right)^2 \right) \\ B &= \frac{1}{2} \left(1 + \cos \theta - \left(\frac{g}{2} \right)^2 \right) \\ C &= \frac{1}{8} \left((1 + \cos \theta) + \frac{2g}{2} \cos \theta + \left(\frac{g}{2} \right)^2 \right); \end{aligned}$$

The above expression can be calculated by using the following integral:

$$\begin{aligned} \frac{1}{\pi} \int_0^{\frac{\pi}{2}} d\theta \ln (A + B \cos \theta + C \cos 2\theta) &= 2 \ln 4 \\ 4 &= \frac{1}{\pi} \int_0^{\frac{\pi}{2}} \frac{A - 3C + \sqrt{(A + C)^2 - B^2}}{2} + \frac{B}{2} \frac{1}{A + B + C} + \frac{C}{2} \frac{1}{A + B + C} : \end{aligned} \quad (D-6)$$

where the above integral has been evaluated by using the well known mean values theorem of analytic function

$$\frac{1}{\pi} \int_0^{\frac{\pi}{2}} d\theta \ln |f(z_0 + R \exp i \theta)| = \ln |f(z_0)|$$

by choosing $f(z) = \frac{1}{2} + \exp i \theta + \exp 2i \theta$. Therefore, the first integral in

(D-1) reads

$$\begin{aligned} &= \ln 2 + 4 \ln \frac{1 + \frac{r}{1 + \frac{q}{1_0}} + \frac{s}{\frac{g}{2}}}{2} \\ &2 \ln \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} + \frac{v}{u} \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} + \frac{2g}{2} \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} + \frac{2g}{2} \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} \ln 2 \\ &= \ln \frac{1 + \frac{r}{1 + \frac{q}{1_0}} + \frac{q}{\frac{g}{2}}}{2} + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2} + \frac{2g}{2} \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} + \frac{2g}{2} \frac{1 + \frac{q}{1 + \frac{q}{1_0}} + \frac{g}{2}}{2} \ln 2; \end{aligned} \tag{D-7}$$

Now, combining it with (D-3) we get the expression (V-4).

Appendix E : Proof of the integral identity (D-5)

Denoting x_k by y and inverting it we get

$$x_k = x_k^{-1}(y); \quad k = 1, 2, \dots, n$$

with n as the number of branch, we can write the lefthand side of relation

(D-5) as

$$\int_0^{x_{f-1}} dx \, f(x) = \sum_{k=1}^n \int_{x_{i_k}}^{x_{f_k}} dx_k \, f(x_k); \quad (E-1)$$

where x_{i_k} (x_{f_k}) denote the end coordinate of the k -th branch) denote its initial coordinate of the k -th branch now, making the change of variable for x_k to y using the relations, $1 = x_{f_k}$, $0 = x_{i_k}$ and $f(y) = f(x_k)$, together with the FP equation we obtain

$$\begin{aligned} \int_0^{x_{f-1}} f(x) dx &= \sum_{k=1}^n \int_0^{x_{f_k}} f(x_k(y)) \frac{dx_k(y)}{dy} dy \\ &= \int_0^{x_{f-1}} dy \, f(y): \end{aligned} \quad (E-2)$$

References

- [1] S.Tasaki, Z.Suchanecki, I.Antoniu. *Physics Letters A*, 179 (1993) 103
- [2] S.Tasaki, T.Gilbert and J.R.Dorfman. *Chaos*, 2 (1998) 424
- [3] M.V.Jakobson. *Communications in Mathematical Physics*, 81 (1981) 39
- [4] T.Gilbert, C.D.Ferguson and J.R.Dorfman. *Phys. Rev. E* 59, 364 (1999)
- [5] K.Umeno. *Phys. Rev. E* 55 (1997) 5280
- [6] K.Umeno. *Jpn. J. Appl. Phys.* Vol. 39 (2000) Part 1, No. 3A, 1442-1456
- [7] G.Froyland. In *proceedings of the 1997 International Symposium on Nonlinear Theory and its Applications, Hawaii, Volume 2* (1997) 1129
- [8] G.Froyland. *Random Computational Dynamics*, 3 (4) (1995) 251.
- [9] M.Blank, G.Keller. *Nonlinearity*, 11 (1998) 1351-1364
- [10] S.Ulam, J.von Neumann, *Bull. Am. Math. Soc.* 53 (1947) 1120.
- [11] Z.X.Wang, D.R.Guo. *Special Functions*, World Scientific Publishing, (1989).
- [12] R.L.Devaney. *An Introduction to Chaotic Dynamical Systems*, Addison Wesley, (1982).
- [13] I.P.Cornfeld, S.V.Fomin and Ya.G.Sinai. *Ergodic Theory*. Springer-Verlag, Berlin, (1982).

- [14] J.R. Dorfman. An Introduction to Chaos in Nonequilibrium Statistical Mechanics, Cambridge, (1999)
- [15] G. Keller. Equilibrium States in a Ergodic Theory, Cambridge, (1998)
23–30

Figures Captions

Fig.1. Plot of $x_2(x; \cdot)$ map for $\alpha = 0:19$ (dashed curve) and plot of $x_3(x; \cdot)$ map for $\alpha = 0:19$, (solid curve). The location of maxima and minima and their values(0 or 1) are independent of parameter α , as it is shown in the figure.

Fig.4. Plot of $x_4(x; \cdot)$ map for $\alpha = 0:19$ (solid curve) and plot of $x_5(x; \cdot)$ map for $\alpha = 0:19$, (dashed curve). The location of maxima and minima and their values(0 or 1) are independent of parameter α , as it is shown in the figure.

Fig.3. Bifurcation diagram of $x_2(x; \cdot)$, for $0 < \alpha < 2$, it is ergodic and for $0 < \alpha < 1$, it has stable fixed point at $x = 1$.

Fig.4. Bifurcation diagram of $x_3(x; \cdot)$, where for $\frac{1}{3} < \alpha < 3$, it is ergodic and for $0 < \alpha < \frac{1}{3}$, it has stable fixed point at $x = 0$, while for $3 < \alpha < 1$, it has stable fixed point $x = 1$.

Fig.5 Bifurcation diagram of $x_{2,3}$, where for $0 < \alpha_1 < 2$ and $\frac{1}{3} < \alpha_2 < 3$ it is ergodic, while for $2 < \alpha_1 < 1$ and $0 < \alpha_2 < \frac{1}{3}$ ($3 < \alpha_2 < 1$) it has fixed point at $[0;0]$ ($[0;1]$).

Fig.6. Bifurcation diagram of $x_{3,2}$, where for $\frac{1}{3} < \alpha_1 < 3$ and $0 < \alpha_2 < 2$ it is ergodic, while for $0 < \alpha_1 < \frac{1}{3}$ ($3 < \alpha_1 < 1$) and $0 < \alpha_2 < 2$ ($2 < \alpha_2 < 1$) it has fixed point at $[0;0]$ ($[1;0]$).

Fig.7. Bifurcation diagram of $x_{3,3}$, where for $\frac{1}{3} < \alpha_1 < 3$ and $\frac{1}{3} < \alpha_2 < 3$ it is ergodic, while for $0 < \alpha_1 < \frac{1}{3}$ ($3 < \alpha_1 < 1$) and $0 < \alpha_2 < \frac{1}{3}$ ($3 < \alpha_2 < 1$) it has fixed point at $[0;0]$ ($[1;1]$).

Fig.8. Solid curve shows the variation of KS-entropy of $x_2(x; \cdot)$, in terms of the parameter α dots show the variation of Lyapunov characteristic exponent of $x_2(x; \cdot)$, in terms of the parameter α .

Fig.9. Solid curve shows the variation of K S-entropy of $\mathcal{S}_3(x; \cdot)$ in terms of the parameter α , dots show the variation of Lyapunov characteristic exponent of $\mathcal{S}_3(x; \cdot)$, in terms of the parameter α .

Fig.10. Solid surface shows the variation of K S-entropy of $\mathcal{S}_{2,2}$ in terms of the parameters α_1 and α_2 , while dotted surface shows the variation of Lyapunov characteristic exponent of $\mathcal{S}_{2,2}$ in terms of the parameters α_1 and α_2 .

Fig.11. Solid surface shows the variation of K S-entropy of $\mathcal{S}_{4,4}$ in terms of the parameters α_1 and α_2 , while dotted surface shows the variation of Lyapunov characteristic exponent of $\mathcal{S}_{4,4}$ in terms of the parameters α_1 and α_2 .

This figure "Fig1.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig2.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig3repair.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig4.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig5.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig6.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig7.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig8.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig9.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig10.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>

This figure "Fig11.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/nlin/0208011v1>