

HEUN EQUATION AND INOZEMTSEV MODELS

KOUCIHI TAKEMURA

ABSTRACT. The BC_N elliptic Inozemtsev model is a quantum integrable systems with N -particles whose potential is given by elliptic functions. Eigenstates and eigenvalues of this model are investigated.

1. INTRODUCTION

The BC_N Inozemtsev model [2] is a system of quantum mechanics with N -particles whose Hamiltonian is given by

$$(1) \quad H = -\sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \sum_{j=1}^N \sum_{i=0}^3 l_i(l_i + 1) \wp(x_j + \omega_i) + 2l(l+1) \sum_{1 \leq j < k \leq N} (\wp(x_j - x_k) + \wp(x_j + x_k)),$$

where $\wp(x)$ is the Weierstrass \wp -function with periods $(1, \tau)$, $\omega_0 = 0$, $\omega_1 = \frac{1}{2}$, $\omega_2 = -\frac{\tau+1}{2}$, $\omega_3 = \frac{\tau}{2}$ are half periods, and l and l_i ($i = 0, 1, 2, 3$) are coupling constants.

It is known that the BC_N Inozemtsev model is quantum completely integrable. More precisely, there exist operators of the form $H_k = \sum_{j=1}^N \left(\frac{\partial}{\partial x_j}\right)^{2k} + (\text{lower terms})$ ($k = 2, \dots, N$) such that $[H, H_k] = 0$ and $[H_{k_1}, H_{k_2}] = 0$ ($k, k_1, k_2 = 2, \dots, N$). Note that the BC_N Inozemtsev model is a universal completely integrable model of quantum mechanics with B_N symmetry, which follows from the classification due to Ochiai, Oshima and Sekiguchi [5]. For the case $N = 1$, finding eigenstates of the Hamiltonian is equivalent to solving the Heun equation [5, 7]. In this sense, the BC_N Inozemtsev model is a generalization of the Heun equation.

In this report, we are going to investigate eigenvalues and eigenstates of the BC_N Inozemtsev model.

A possible approach is to use the quasi-exact solvability. If the coupling constants l, l_0, l_1, l_2, l_3 satisfy some equation, the Hamiltonian H (see (1)) and the commuting operators of conserved quantities preserve some finite dimensional space of doubly periodic functions [1, 10]. On the finite dimensional space, eigenvalues are calculated by solving the

characteristic equation, and eigenfunctions are obtained by solving linear equations. In this sense, a part of eigenvalues and eigenfunctions is obtained exactly and this is the reason to use the phrase “quasi-exact solvability”.

Another approach is to use a method of perturbation. By the trigonometric limit $p = \exp(\pi\sqrt{-1}\tau) \rightarrow 0$, the Hamiltonian H of the BC_N elliptic Inozemtsev model tends to the Hamiltonian of the BC_N Calogero-Moser-Sutherland model H_T (see (2)), and eigenvalues and eigenstates of the BC_N Calogero-Moser-Sutherland model are known (see Proposition 2.1).

Based on the eigenstates for the case $p = 0$, we can obtain eigenvalues and eigenstates of the BC_N elliptic Inozemtsev model ($p \neq 0$) as formal power series in p . This procedure is sometimes called the algorithm of perturbation. Generally speaking, convergence of the formal power series obtained by perturbation is not guaranteed a priori, but for the case of the BC_N elliptic Inozemtsev model, the convergence radius of the formal power series in p is shown to be non-zero (see Corollary 3.4), and it is seen that this perturbation is holomorphic. As a result, real-holomorphy of the eigenvalues in p and the completeness of the eigenfunctions are shown.

In section 2, some results on the BC_N trigonometric Calogero-Moser-Sutherland model are reviewed. In section 3, we present some propositions about the perturbation of the BC_N elliptic Inozemtsev model from the trigonometric model. In section 4, we give some comments for future problems.

2. TRIGONOMETRIC LIMIT

In this section, we will consider the trigonometric limit ($\tau \rightarrow \sqrt{-1}\infty$) and review some results on the trigonometric model. Assume $l, l_0, l_1 \geq 0$ and set $p = \exp(\pi\sqrt{-1}\tau)$. Then $p \rightarrow 0$ as $\tau \rightarrow \sqrt{-1}\infty$.

If $p \rightarrow 0$, then $H \rightarrow H_T + C_T$, where

$$(2) \quad H_T = -\sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \sum_{1 \leq j < k \leq N} \left(\frac{2\pi^2 l(l+1)}{\sin^2 \pi(x_j - x_k)} + \frac{2\pi^2 l(l+1)}{\sin^2 \pi(x_j + x_k)} \right) \\ + \sum_{j=1}^N \left(\frac{\pi^2 l_0(l_0+1)}{\sin^2 \pi x_j} + \frac{\pi^2 l_1(l_1+1)}{\cos^2 \pi x_j} \right),$$

and $C_T = -\frac{N(N-1)\pi^2}{3}l(l+1) - \frac{N\pi^2}{3}\sum_{i=0}^3 l_i(l_i+1)$. The operator H_T is nothing but the Hamiltonian of the BC_N trigonometric Calogero-Moser-Sutherland model.

Now we solve the spectral problem for H_T by using hypergeometric functions. Set

$$(3) \quad \Phi_T(x) = \prod_{j=1}^N (\sin \pi x_j)^{l_0+1} (\cos \pi x_j)^{l_1+1} \\ \prod_{1 \leq j_1 < j_2 \leq N} (\sin \pi(x_{j_1} - x_{j_2}) \sin \pi(x_{j_1} + x_{j_2}))^{l+1}.$$

Let $D = \{(x_1, \dots, x_N) \in \mathbf{R}^N \mid 0 < x_1 \leq \dots \leq x_N < 1\}$. A function $f(x_1, \dots, x_N)$ is $W(B_N)$ -invariant iff $f(x_1, \dots, x_i, \dots, x_j, \dots, x_N) = f(x_1, \dots, x_j, \dots, x_i, \dots, x_N)$ for all $i < j$ and $f(x_1, \dots, x_i, \dots, x_N) = f(x_1, \dots, -x_i, \dots, x_N)$ for all i . The Hilbert space \mathbf{H} is defined by

$$(4) \quad \mathbf{H} = \left\{ f: \mathbf{R}^N \rightarrow \mathbf{C} \text{ measurable} \left| \begin{array}{l} \int_D |f(x)|^2 dx < +\infty, \\ \frac{f(x)}{\Phi_T(x)} \text{ is } W(B_N)\text{-invariant a.e. } x, \\ \frac{f(x+n)}{\Phi_T(x+n)} = \frac{f(x)}{\Phi_T(x)}, \forall n \in \mathbf{Z}^N \text{ a.e. } x \end{array} \right. \right\}.$$

Proposition 2.1. *Let $\mathcal{M}_N = \{(\lambda_1, \dots, \lambda_N) \in \mathbf{Z}^N \mid \lambda_1 \geq \dots \geq \lambda_N \geq 0\}$. There exists a complete orthonormal system $\{v_\lambda\}_{\lambda=(\lambda_1, \dots, \lambda_N) \in \mathcal{M}_N}$ on the Hilbert space \mathbf{H} such that*

$$H_T v_\lambda = E_\lambda v_\lambda,$$

where $E_\lambda = \pi^2 \sum_{j=1}^N (2\lambda_j + l_0 + l_1 + 2 + 2(l+1)(N-j))^2$.

Note that the function v_λ is expressed as the product of the ground-state $\Phi_T(x)$ and the BC_N Jacobi polynomial $\psi_\lambda(z_1, \dots, z_N)$ ($z_j = \exp(2\pi\sqrt{-1}x_j)$, $j = 1, \dots, N$). Essential selfadjointness of the operator H_T on the Hilbert space \mathbf{H} is obtained by applying Proposition 2.1.

3. PERTURBATION FROM THE TRIGONOMETRIC MODEL

We apply a method of perturbation and have an algorithm for obtaining eigenvalues and eigenfunctions as formal power series of p . The proofs of propositions in this report are obtained by imitating the proofs of the corresponding propositions in [4, 8]

Set $p = \exp(\pi\sqrt{-1}\tau)$. For the Hamiltonian of the BC_N Inozemtsev model, we adopt a notation $H(p)$ instead of H . The operator $H(p)$ admits the following expansion:

$$(5) \quad H(p)(= H) = H_T + C_T + \sum_{k=1}^{\infty} V_k(x)p^k,$$

where H_T is the Hamiltonian of trigonometric model defined in (2), $V_k(x)$ is some function in variables x_1, \dots, x_N , and C_T is a constant defined in section 2.

Based on the eigenvalues E_λ ($\lambda \in \mathcal{M}_N$) and the eigenfunctions v_λ of the operator H_T , we determine eigenvalues $E_\lambda(p) = E_\lambda + C_T + \sum_{k=1}^{\infty} E_\lambda^{\{k\}} p^k$ and normalized eigenfunctions $v_\lambda(p) = v_\lambda + \sum_{k=1}^{\infty} \sum_{\mu \in \mathcal{M}_N} c_{\lambda, \mu}^{\{k\}} v_\mu p^k$ of the operator $H(p)$ as formal power series in p . In other words, we will find $E_\lambda(p)$ and $v_\lambda(p)$ that satisfy equations

$$(6) \quad H(p)v_\lambda(p) = (H_T + C_T + \sum_{k=1}^{\infty} V_k(x)p^k)v_\lambda(p) = E_\lambda(p)v_\lambda(p),$$

$$\langle v_\lambda(p), v_\lambda(p) \rangle = 1,$$

as formal power series of p .

First we calculate coefficients $\sum_{\mu \in \mathcal{M}_N} d_{\lambda, \mu}^{\{k\}} v_\mu = V_k(x)v_\lambda$ ($k \in \mathbf{Z}_{>0}$, $\lambda \in \mathcal{M}_N$). Next we compute $E_\lambda^{\{k\}}$ and $c_{\lambda, \mu}^{\{k\}}$ for $k \geq 1$ and $\lambda, \mu \in \mathcal{M}_N$. By comparing coefficients of $v_\mu p^k$ in (6), we obtain recursive relations for $E_\lambda^{\{k\}}$ and $c_{\lambda, \mu}^{\{k\}}$.

Now we present results which are obtained by applying the Kato-Rellich theory. We use definitions written in Kato's book [3] freely. It is shown that the operator $H(p)$ ($-1 < p < 1$) is essentially self-adjoint on the Hilbert space \mathbf{H} , because the operator $\sum_{k=1}^{\infty} V_k(x)p^k$ is bounded. Let $\tilde{H}(p)$ ($-1 < p < 1$) be the unique extension of $H(p)$ to the selfadjoint operator.

Proposition 3.1. *The operators $\tilde{H}(p)$ form a holomorphic family of type (A) for $-1 < p < 1$.*

Proposition 3.2. *The spectrum $\sigma(\tilde{H}(p))$ contains only point spectra and it is discrete. The multiplicity of each eigenvalue is finite.*

Combining Theorem 3.9 in [3, VII-§3.5], propositions in this report and the selfadjointness of $\tilde{H}(p)$, the following theorem is proved:

Theorem 3.3. *All eigenvalues of $\tilde{H}(p)$ ($-1 < p < 1$) can be represented as $E_\lambda(p)$ ($\lambda \in \mathcal{M}_N$), which is real-holomorphic in $p \in (-1, 1)$ and $E_\lambda(0) = E_\lambda + C_T$. The eigenfunction $v_\lambda(p)$ of the eigenvalue $E_\lambda(p)$ is holomorphic in $p \in (-1, 1)$ as an element in L^2 -space, and the eigenvectors $v_\lambda(p)$ ($\lambda \in \mathcal{M}_N$) form a complete orthonormal family on \mathbf{H} .*

As an application of the theorem, the convergence of formal power series of eigenvalues in the variable p obtained by the algorithm of perturbation is shown.

Corollary 3.4. *Let $E_\lambda(p)$ ($\lambda \in \mathcal{M}_N$) (resp. $v_\lambda(p)$) be the formal eigenvalue (resp. eigenfunction) of the Hamiltonian $H(p)$ defined by (6). If $|p|$ is sufficiently small then the power series $E_\lambda(p)$ converges and as an element in L^2 space the power series $v_\lambda(p)$ converges.*

4. COMMENTS

In this report, holomorphy of perturbation for the Hamiltonian of the BC_N Inozemtsev model from the trigonometric one is established. Relationship between the perturbation and the complete integrability should be clarified. More precisely, holomorphy of perturbation for commuting operators of conserved quantities should be shown, although it is not succeed as of this writing. For the elliptic Calogero-Moser-Sutherland model of type A_N , holomorphy of perturbation for commuting operators is shown in [4, 6].

In [7, 9], Bethe Ansatz method for the BC_1 Inozemtsev model is proposed and some results on the finite gap integration for the BC_1 Inozemtsev model are established. We hope that some progress on the Bethe Ansatz method or the finite gap integration for the BC_N Inozemtsev model will be made.

REFERENCES

- [1] Finkel F, Gomez-Ullate D, Gonzalez-Lopez A, Rodriguez M A, and Zhdanov R 2001 New spin Calogero-Sutherland models related to B_N -type Dunkl operators *Nuclear Phys. B* **613** 472–496
- [2] Inozemtsev V I 1989 Lax representation with spectral parameter on a torus for integrable particle systems *Lett. Math. Phys.* **17** 11–17
- [3] Kato T 1980 Perturbation theory for linear operators, corrected printing of the second ed. (Berlin: Springer-Verlag).
- [4] Komori Y and Takemura K 2002 The perturbation of the quantum Calogero-Moser-Sutherland system and related results *Commun. Math. Phys.* **227** 93–118
- [5] Ochiai H, Oshima T, and Sekiguchi H 1994 Commuting families of symmetric differential operators *Proc. Japan. Acad.* **70** 62–66.
- [6] Takemura K 2000 On the eigenstates of the elliptic Calogero-Moser model *Lett. Math. Phys.* **53** 181-194
- [7] Takemura K 2003 The Heun equation and the Calogero-Moser-Sutherland system I: the Bethe Ansatz method, to appear in *Commun. Math. Phys.*, math.CA/0103077
- [8] Takemura K 2001 The Heun equation and the Calogero-Moser-Sutherland system II: the perturbation and the algebraic solution *Preprint* math.CA/0112179
- [9] Takemura K 2002 The Heun equation and the Calogero-Moser-Sutherland system III: the finite gap property and the monodromy *Preprint* math.CA/0201208
- [10] Takemura K 2002 Quasi-exact solvability of Inozemtsev models *J. Phys. A* **35** 8867-8881

DEPARTMENT OF MATHEMATICAL SCIENCES, YOKOHAMA CITY UNIVERSITY,
22-2 SETO, KANAZAWA-KU, YOKOHAMA 236-0027, JAPAN