

Dirac reduction revisited

Krzysztof M ARC IN IAK ^y and Maciej B LA SZAK ^z

^y Department of Science and Technology
Campus Norrköping, Linköping University
601-74 Norrköping, Sweden
E-mail: krzma@itn.liu.se

^z Institute of Physics, A . M ickiew icz University
Umultowska 85, 61-614 Poznań, Poland
E-mail: blaszakm@amu.edu.pl

March 7, 2003

Abstract

The procedure of Dirac reduction of Poisson operators on submanifolds is discussed within a particularly useful special realization of the general Marsden-Ratiu reduction procedure. The Dirac classification of constraints on 'first-class' constraints and 'second-class' constraints is reexamined.

AMS 2000 Subject Classification: 70H45, 53D17, 70G45

1 Introduction

Dirac bracket as well as Dirac's classification of constraints is nowadays a well recognized and very useful tool in the construction of Poisson dynamics on admissible submanifolds from a given Poisson dynamics on a given manifold. In this paper we consider the Dirac reduction procedure in a more general setting than is usually met in literature. In Section 2 we implement the Dirac reduction procedure into a particularly useful special realization of the general Marsden-Ratiu reduction scheme, based on the concept of transversal distributions. In Section 3 we reconsider the Dirac concept of first class constraints as it seems to be too restrictive.

Firstly we recall few basic notions from Poisson geometry. Given a manifold M , a Poisson operator on M is a mapping $\pi : T^*M \rightarrow TM$ that is bre-preserving (i.e. $\pi_{xM} : T_x^*M \rightarrow T_xM$ for any $x \in M$) and such that the induced bracket on the space $C^1(M)$ of all smooth real-valued functions on M

$$f;g : C^1(M) \times C^1(M) \rightarrow C^1(M), \quad f;g \stackrel{\text{def}}{=} \pi^*df; \pi^*dg \quad (1)$$

Partially supported by KBN grant No. 5P 03B 004 20

where h_{ij} is the dual map between T^*M and T^*M , is skew-symmetric and satisfies Jacobi identity (the bracket $\{ \}$ always satisfies the Leibniz rule $\{F, GH\} = G\{F, H\} + H\{F, G\}$). The symbol d denotes the operator of exterior differentiation. The operator $\{ \}$ can always be interpreted as a bivector, $\{ \} \in \wedge^2(T^*M)$ and in a given coordinate system (x^1, \dots, x^m) on M we have

$$\{ \} = \sum_{i < j}^{X^n} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j};$$

A function $C : M \rightarrow \mathbb{R}$ is called a Casimir function of the Poisson operator if for an arbitrary function $F : M \rightarrow \mathbb{R}$ we have $\{F, C\} = 0$ or, equivalently, if $dC = 0$.

2 Marsden-Ratiu reduction for transversal distributions

The Marsden-Ratiu reduction theorem [1] describes the procedure of reducing a Poisson operator on arbitrary submanifold S of our manifold M . This general procedure exists only if some conditions are satisfied. These conditions involve a distribution E (in the original notation of Marsden and Ratiu) that is a subbundle of T^*M . By a simple assumption, namely that this distribution is transversal, one can, however, satisfy all these conditions automatically. Below we reformulate the Marsden-Ratiu theorem in this more limited but useful setting.

Consider an m -dimensional manifold M equipped with a Poisson operator and an s -dimensional submanifold S of M . Fix a distribution Z of constant dimension $k = m - s$, that is a smooth collection of m -dimensional subspaces $Z_x \subset T_x M$ at every point x in M , which is transversal to S in the sense that no vector field $Z \in Z$ is at any point tangent to the submanifold S . Hence we have

$$T_x M = T_x S \oplus Z_x$$

for every $x \in S$ and, similarly,

$$T_x M = T_x S \oplus Z_x;$$

where $T_x S$ is the annihilator of Z_x and Z_x is the annihilator of $T_x S$. That means that if α is a one-form in $T_x S$ then $\alpha(Z) = 0$ for all vectors $Z \in Z_x$ and if β is a one-form in Z_x then β vanishes on all vectors in $T_x S$.

Definition 1 A function $F : M \rightarrow \mathbb{R}$ is invariant with respect to Z if $L_Z F = Z(F) = 0$ for any $Z \in Z$. Similarly, a function $F : M \rightarrow \mathbb{R}$ is invariant with respect to Z on S ($Z|_S$ -invariant in short) if $L_Z F|_S = Z(F)|_S = 0$ for any $Z \in Z$.

Here and in what follows the symbol L_Z means the Lie derivative along the vector field Z .

Definition 2 An operator $\{ \}$ is called invariant with respect to the distribution Z if the functions that are Z -invariant form a Poisson subalgebra, that is, if $F, G : M \rightarrow \mathbb{R}$ are two Z -invariant functions then $\{F, G\}$ is again a Z -invariant function. Similarly, an operator $\{ \}$ is called invariant with respect to the distribution Z on S if the functions that are $Z|_S$ -invariant form a Poisson subalgebra, that is, if $F, G : M \rightarrow \mathbb{R}$ are two $Z|_S$ -invariant functions then $\{F, G\}$ is again a $Z|_S$ -invariant function.

We denote these Poisson subalgebras by A and A_S respectively. Let us observe, that if \mathcal{Z} is \mathcal{Z}_j -invariant for any manifold S in a foliation of M then it is also \mathcal{Z} -invariant.

Theorem 3 (Marsden and Ratiu [1]): Let S be a submanifold of M equipped with a Poisson operator and let \mathcal{Z} be a distribution in M that is transversal to S . If the operator is invariant with respect to the distribution \mathcal{Z} on S , then the Poisson operator is reducible on S in the sense that on S there exists a (uniquely defined) Poisson operator P_R such that for any $f, g : S \rightarrow \mathbb{R}$ we have

$$fP_R g = P_R(f)g \quad (2)$$

for any \mathcal{Z}_j -invariant prolongations F and G of f and g respectively.

The above construction, however, is difficult to perform in practice since it is often hard to find explicit expressions for the prolongations F and G . We now show how this difficulty can be omitted.

Firstly, suppose that our submanifold S is given by k functionally independent equations $\phi_i(x) = 0, i = 1, \dots, k$ (constraints) and that our transversal distribution \mathcal{Z} is spanned by k vector fields Z_i chosen such that the following orthogonality relation holds

$$\langle d\phi_i, Z_j \rangle = \delta_{ij} \quad (3)$$

(this is no restriction since for any distribution \mathcal{Z} transversal to S we can choose its basis so that (3) is satisfied). We observe that in this case we have $[Z_i, Z_j] = 0$ for all i, j , where $[X, Y] = L_X Y - Y(X)$ is the Lie bracket (commutator) of the vector fields X, Y , so that $[Z_i, Z_j]$ is always tangent to S . Then, in case that the distribution \mathcal{Z} is involutive (integrable), this means that $[Z_i, Z_j] = 0$ for all i, j . Moreover, we define the vector fields X_i as

$$X_i = d\phi_i, \quad i = 1, \dots, k \quad (4)$$

There exists an important class of \mathcal{Z} -invariant Poisson operators

Lemma 4 [2] If

$$L_{Z_i} = \sum_{j=1}^k W_j^{(i)} \wedge Z_j, \quad i = 1, \dots, k \quad (5)$$

for some vector fields $W_j^{(i)}$, then the Poisson operator is invariant with respect to \mathcal{Z}

We sketch the proof here for the clarity of the text.

Proof. Assume, that $L_{Z_i} F = L_{Z_i} G = 0$ for all i . We have to show that $L_{Z_i} fP_R g = 0$ for all i , but, due to (5)

$$L_{Z_i} fP_R g = L_{Z_i} \langle dF, dG \rangle = \sum_{j=1}^k \langle dF, d(W_j^{(i)} \wedge Z_j) \rangle dG$$

since $L_{Z_i} (dF) = d(L_{Z_i} F) = 0$ (and similarly for G). On the other hand

$$\langle dF, d(W_j^{(i)} \wedge Z_j) \rangle = Z_j \langle F, W_j^{(i)} \rangle - Z_j \langle F, W_j^{(i)} \rangle \langle G \rangle = 0$$

since $Z_j(F) = L_{Z_j}F = 0$ (and similarly for G). ■

The condition (5) is sufficient but not necessary. For example, if

$$L_{Z_i} = \sum_{j=1}^k W_j \wedge [Z_i, Z_j] \quad i=1, \dots, k$$

for some vector fields W_i , then the operator is also Z -invariant (one shows it by computations similar to those in the above proof). In the case when (5) we apply the Lie derivative L_{Z_j} to both sides of the equation (4). Due to (5) we obtain

$$\begin{aligned} [Z_j, X_i] &= L_{Z_j}X_i = (L_{Z_j})d'_{i1} = \sum_1^X W_1^{(j)} \wedge Z_1 d'_{i1} = \\ &= \sum_1^X Z_1('_{i1})W_1^{(j)} - W_1^{(j)}('_{i1})Z_1 = W_i^{(j)} \sum_1^X W_1^{(j)}('_{i1})Z_1 \end{aligned} \quad (6)$$

We observe that, if F and G are two Z_j -invariant functions and V_j are arbitrary vector fields, then $dF; \sum_j V_j \wedge Z_j dG|_S = 0$ since $hdF; V_j \wedge Z_j dG|_S = Z_j(G)V_j(F) - Z_j(F)V_j(G) = 0$ on S . Thus the Poisson operator and its deformation of the form

$$D = \sum_j V_j \wedge Z_j \quad (7)$$

in spite of the fact that they act differently on A_S both generate the same bracket on S so that both can be used to define our restricted operator R on S through (2). Of course, the deformed operator D does not have to be Poisson, but nevertheless its restriction to S through (2) must be Poisson since it naturally coincides with similar restriction of P to S . If we now consider a whole foliation of M defined by the functions $'_i$ with leaves S (so that $S_0 = S$) then it turns out that we can choose our (undetermined so far) vector fields V_j in (7) so that

$$D(x) \in T_x S \quad \text{for any } x \in T_x M \text{ and any } x \in M; \quad (8)$$

which has a far reaching consequence.

Lemma 5 The deformation D given by (7) that also satisfies (8) is Poisson.

Proof. The condition that $D(x)$ is tangent to S for any $x \in T_x M$ is equivalent to the requirement that $hd'_{i1}; D(x)|_S = 0$ for all i . Due to the antisymmetry of D this requirement can be rewritten as $h_x; D('_{i1})|_S = 0$ for all i . Since x is arbitrary, the condition attains the form $D(d'_{i1}) = 0$ for $i = 1, \dots, k$. We now complete the set of functions $'_i$ with some functions x_j to a coordinate system $(x; ')$ on M . Then the matrix of the operator D has the last k rows and last k columns equal to zero while the $m-k$ dimensional upper left block coincides on every leaf S with the corresponding R which is Poisson by the Marsden-Ratiu construction. ■

Lem m a 6 The condition (8) can be written as

$$V_i - \sum_{j=1}^{X^k} V_j ('_i) Z_j = X_i: \quad (9)$$

P roof. We know that the condition (8) can be written as $_D (d'_i) = 0$ for $i = 1; \dots; k$. An easy calculation yields now that

$$\begin{aligned} 0 = _D (d'_i) &= (d'_i) - \sum_{j=1}^{X^k} (Z_j ('_i) V_j - V_j ('_i) Z_j) = \\ &= X_i - V_i + \sum_{j=1}^{X^k} V_j ('_i) Z_j \end{aligned}$$

due to the normalization condition (3). ■

We now restrict ourselves to only two limit cases, when all X_i are tangent to S and when X_i span Z .

2.1 The case when X_i are tangent to S

We firstly assume that all the vectors X_i are tangent to S . We have then naturally $X_i ('_j) = 0$. This in turn means that $f'_i; ' _j g = h d'_i; d'_j i = h d'_i; X_j i = 0$ so that all the vector fields X_i commute. In this case the simplest solution of (9) has the form $V_i = X_i$ and the corresponding deformation (7) attains the form

$$_D = \sum_{i=1}^{X^k} X_i \wedge Z_i: \quad (10)$$

This deformation has been recently widely used for projecting Poisson pencils on symplectic leaves of one of their operators [3]–[5].

Lem m a 7 [3] The vector fields $W_j^{(k)}$ in (5) can, in the case that all X_i are tangent to S , be chosen as tangent to S .

P roof. Consider the projections $\tilde{W}_j^{(i)}$ of the vector fields $W_j^{(i)}$ onto S :

$$\tilde{W}_j^{(i)} = W_j^{(i)} - \sum_{r=1}^{X^k} W_j^{(i)} ('_r) Z_r:$$

If $W_j^{(i)}$ are in Z , then $\tilde{W}_j^{(i)} = 0$. The vector field $\tilde{W}_j^{(i)}$ is indeed tangent to S since

$$\tilde{W}_j^{(i)} ('_1) = W_j^{(i)} ('_1) - \sum_{r=1}^{X^k} W_j^{(i)} ('_r) 1_r = 0:$$

Now

$$\sum_{j=1}^{X^k} \tilde{W}_j^{(i)} \wedge Z_j = \sum_{j=1}^{X^k} W_j^{(i)} \wedge Z_j - \sum_{j;r=1}^{X^k} W_j^{(i)} ('_r) Z_r \wedge Z_j$$

the last term being equal to zero since $L_{Z_k} f'_i; ' _j g = 0$ implies $W_j^{(i)} ('_r) = W_r^{(i)} ('_j)$. Thus $\sum_{j=1}^{X^k} W_j^{(i)} \wedge Z_j = \sum_{j=1}^{X^k} \tilde{W}_j^{(i)} \wedge Z_j$. ■

Due to this gauge freedom, if we choose $W_j^{(i)}$ as tangent to S (which means that $W_j^{(i)} ('_r) = 0$) then the formula (6) yields that $W_j^{(i)} = [Z_i; X_j]$: Thus, due to the fact that we assumed (5),

$$L_{Z_i} = \sum_{j=1}^{X^k} [Z_i; X_j] \wedge Z_j: \quad (11)$$

Remark 8 In the case that the functions $'_i$ are Casimir functions of ω we have $X_i = d'_i = 0$ so that the formula (11) yields $L_{Z_i} = 0$ for all i ; i.e. the vector fields Z_i are symmetries of ω . In this case our reduction procedure (2) coincides with the standard projection onto a level set of Casimir functions (symplectic leaf in case there are no other Casimirs apart from $'_i$) [6].

From what we have said above it becomes clear that the above reduction scheme can be interpreted as a two-step procedure: firstly we deform the original Poisson tensor to a Poisson tensor ω_D and then we obtain ω_R as standard projection of ω_D onto the level set S of its Casimirs $'_i$ (thus we need not calculate the prolongations F and G in order to define ω_R).

Now we check what can be said about our vector fields Z_i .

According to Remark 8 $L_{Z_i} \omega_D = 0$. On the other hand, due to (10),

$$0 = L_{Z_i} \omega_D = \sum_{j=1}^{X^k} [Z_i; X_j] \wedge Z_j - \sum_{j=1}^{X^k} L_{Z_i} X_j \wedge Z_j - \sum_{j=1}^{X^k} X_j \wedge L_{Z_i} Z_j$$

so that $\sum_{j=1}^{X^k} X_j \wedge [Z_i; Z_j] = 0$. Of course one of the possible realizations of this condition is the case that the distribution Z be integrable since then $[Z_i; Z_j] = 0$. There are, however, other possibilities here. For example, if $[Z_i; Z_j] = \sum_{s=1}^{X^k} c_{ij}^s X_s$ with $c_{ij}^s = c_{sj}^i$, $\sum_{j=1}^{X^k} X_j \wedge [Z_i; Z_j] = 0$ as well.

2.2 The case when X_i span Z

This time we assume that $X_i = \sum_{k=1}^P '_{ki} Z_k$ for some real valued functions $'_{ij}$, which due to (3) yields

$$'_{ij} = \sum_{k=1}^P '_{kj} Z_k ('_i) = X_j ('_i) = f'_i; ' _j g: \quad (12)$$

The functions $'_{ij}$ define a k -dimensional skew-symmetric matrix $' = ('_{ij}); i, j = 1; \dots; k$. The only condition imposed on $'$ is related to the determinant and that X_i span Z , i.e. $\det' \neq 0$: We thus do not have to assume (5) this time since now the distribution Z is spanned by

the Hamiltonian vector fields X_i and thus is automatically invariant with respect to Z as $L_{X_i} = 0$ for all i . It can be easily shown that

$$[X_j, X_i] = X_{f'_{ij}} = df'_{ij} = d'_{ij}.$$

Now we look for solutions of (9) in the simple form $V_i = X_i$. Inserting this into (9) and using the fact that $f'_{ij} = -f'_{ji}$ we obtain

$$0 = X_i \sum_{j=1}^{X^k} X_j (f'_{ij}) Z_j - X_i = X_i + \sum_{j=1}^{X^k} f'_{ji} Z_j - X_i = (2 - 1) X_i$$

so that $a = 1=2$ and $V_i = \frac{1}{2} X_i$. In this case the deformation (7) attains the form :

$$D = \frac{1}{2} \sum_{i=1}^{X^k} X_i \wedge Z_i \quad (13)$$

and is, as mentioned above, Poisson. It is easy to check that our operator D defines the following bracket on M

$$fF;Gg_D = fF;Gg + \sum_{i,j=1}^{X^k} fF;f'_{ij}g (-f'_{ij})_{ij} f'_{ij}Gg; \quad (14)$$

where $f, G : M \rightarrow \mathbb{R}$ are now two arbitrary functions on M , which is just the well known Dirac deformation [7] of the bracket $f;g$ associated with D .

Remark 9 If $C : M \rightarrow \mathbb{R}$ is a Casimir function of D , then it is also a Casimir function of D since in this case (14) yields

$$fF;Cg_D = fF;Cg + \sum_{i,j=1}^{X^k} fF;f'_{ij}g (-f'_{ij})_{ij} f'_{ij}Cg = 0 \quad 0 = 0: \quad (15)$$

We also know that the constraints f'_{ij} are Casimirs of the deformed operator D . Thus we can state that Dirac deformation preserves all the old Casimir functions and introduces new Casimirs f'_{ij} .

It is now possible to restrict our Poisson operator D (or our Poisson bracket $f;g_D$) to a Poisson operator D_R (bracket $f;g_R$) on the submanifold S , i.e. the level set $f'_{ij} = \dots = f'_{mn} = 0$ of Casimirs of D , in a standard way.

3 Existence of Dirac reduction

We now present some realizations of the above Dirac case and discuss the classical concept of the Dirac classification of constraints. We will show that the classification of constraints as being either of first-class or of second-class, proposed by Dirac, should be reexamined when one looks at the problem from a more general point of view.

We recall that a constraint ϕ_k is of first class if its Poisson bracket with all the remaining constraints ϕ_i vanishes on S , that is if

$$\{\phi_k, \phi_i\} = 0; \quad i = 1, \dots, m; \quad (16)$$

Otherwise ϕ_k is of second-class. In the case that at least one of the constraints is of the first class, the matrix γ_{ij} in (12) is singular on S so that the formula (14) cannot be used in order to define R . However, it may still be possible to define R via the above general scheme. This indicates that the concept of first class constraint is too narrow. Below we demonstrate the examples of Dirac reduction in case when constraints are of first class.

We start with a simple example. Consider a $2n$ -dimensional manifold M parametrized by coordinates $(q_1, \dots, q_n; p_1, \dots, p_n)$ and equipped with a Poisson operator of the form

$$= \begin{pmatrix} 0 & Q_n \\ Q_n & 0 \end{pmatrix};$$

where Q is a diagonal matrix of the form $Q_n = \text{diag}(q_1, \dots, q_n)$. Consider a submanifold S given by a pair of constraints $\phi_1(q; p) = q_n = 0$ and $\phi_2(q; p) = p_n = 0$. Then the matrix γ has the form

$$\gamma = \begin{pmatrix} 0 & q_n \\ q_n & 0 \end{pmatrix}$$

so that it is clearly singular on S ($\det(\gamma) = 0$ on S) and

$$\gamma^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \frac{1}{q_n}$$

so that the Dirac formula (14) cannot be applied. However, the vector fields $Z_1 = q_n^{-1} X_2$ and $Z_2 = -q_n^{-1} X_1$ that span our distribution Z are not singular on S since $X_1 = q_n \frac{\partial}{\partial p_n}$ and $X_2 = q_n \frac{\partial}{\partial q_n}$ so that the deformation (13) becomes

$$D = \frac{1}{q_n} X_1 \wedge X_2 = q_n \frac{\partial}{\partial q_n} \wedge \frac{\partial}{\partial p_n} = \sum_{i=1}^{n-1} q_i \frac{\partial}{\partial q_i} \wedge \frac{\partial}{\partial p_i}$$

and is clearly reducible on S . The operator R obtained on S parametrized by coordinates $(q_1, \dots, q_{n-1}; p_1, \dots, p_{n-1})$ is

$$R = \begin{pmatrix} 0 & Q_{n-1} \\ Q_{n-1} & 0 \end{pmatrix}$$

This simple example clearly illustrates that Dirac's classification is too strong. As a second example we consider a particle moving in a Riemannian manifold Q of dimension three with a contravariant metric tensor

$$G = \begin{pmatrix} 2 & 0 & 3 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

given in some coordinates $(q^1; q^2; q^3)$. Suppose that this particle is subordinated to a holonomic constraint on Q given by

$$\phi_1(q) = q^1 q^2 + q^3 = 0; \quad (17)$$

This defines a submanifold of Q : The velocity $v = \sum_{i=1}^3 v^i \partial/\partial q^i$ of this particle must then remain tangent to this submanifold so that

$$0 = \text{hd}'_k; v^i = \sum_{i=1}^3 \frac{\partial'_{\text{k}}}{\partial q^i} v^i;$$

and thus in our coordinates $v^i = \sum_j^P G^{ij} p_j$ the motion of the particle in the phase space $M = T^*Q$ is constrained not only by (17) but also by the relation

$$\sum_{i,j=1}^3 G^{ij} \frac{\partial'_{\text{k}}}{\partial q^i} p_j - p_1 + p_2 q^1 + p_3 q^2 = 0 \quad (18)$$

that is nothing else than the lift of (17) to M . The constraints (17)–(18) define a four-dimensional submanifold S of M . We now introduce the following Poisson structure on M :

$$= \begin{pmatrix} 2 & 0 & 0 & 0 & q^1 & 1 & 0 \\ 6 & 0 & 0 & 0 & q^2 & 0 & 1 \\ 6 & 0 & 0 & 0 & 2q^3 & q^2 & q^1 \\ 6 & q^1 & q^2 & 2q^3 & 0 & p_2 & p_3 \\ 4 & 1 & 0 & q^2 & p_2 & 0 & 0 \\ 5 & 0 & 1 & q^1 & p_3 & 0 & 0 \end{pmatrix}:$$

Again the matrix $'$ is singular, since $'_{12} = 2(q^1 q^2 + q^3) = 2'_{11}$ which obviously vanishes on S . One can, however, perform the deformation (13). A quite lengthy but straightforward computation shows that in this case

$$D = \begin{pmatrix} 2 & 0 & 0 & 0 & q^1 & 1 & 0 \\ 6 & 0 & 0 & 0 & q^2 & 0 & 1 \\ 6 & 0 & 0 & 0 & 2q^1 q^2 & q^2 & q^1 \\ 6 & q^1 & q^2 & 2q^1 q^2 & 0 & p_2 & p_3 \\ 4 & 1 & 0 & q^2 & p_2 & 0 & 0 \\ 5 & 0 & 1 & q^1 & p_3 & 0 & 0 \end{pmatrix}$$

and this operator can be projected onto S . To do this, one can first pass to the Casimir variables

$$(q^1; q^2; '_{11}(q); '_{22}(q; p); p_2; p_3)$$

since, due to the fact that it is easiest to eliminate q^3 and p_1 from the system of equations $'_{11} = '_{11}(q) = 0; '_{22} = '_{22}(q; p) = 0$, we parametrize our submanifold by the coordinates $(q^1; q^2; p_2; p_3)$. In these variables the operator R attains the canonical form

$$R = \begin{pmatrix} 2 & 0 & 0 & 1 & 0 \\ 6 & 0 & 0 & 0 & 1 \\ 4 & 1 & 0 & 0 & 0 \\ 5 & 0 & 1 & 0 & 0 \end{pmatrix}:$$

Our two examples show that the condition (16) is only a necessary condition for nonexistence of R on S , but is not a sufficient one. Hence the definition of first class constraints

has to be made weaker. Even in the case when we deal with a real first class constraint we can obtain π_R on S coming from the Dirac reduction of π : We demonstrate this below.

Firstly we assume that we have a pair of constraints f'_1, f'_2 that define our submanifold $S = \{f'_1 = 0, f'_2 = 0\}$ and such that they are of second class, i.e. that $\{f'_1, f'_2\} \neq 0$. It is clear that our submanifold S can be parametrized in infinitely many different ways by constraints $e_1 = 0, e_2 = 0$; where

$$e_1 = f'_1 + \alpha_2 f'_2; \quad e_2 = \alpha_3 f'_1 + \alpha_4 f'_2 \quad (19)$$

and where α_i are some functions on M such that $\alpha_i \neq 0$ and such that

$$D \frac{D(e_1, e_2)}{D(f'_1, f'_2)} = \alpha_1 \alpha_4 - \alpha_2 \alpha_3 \neq 0 \quad (20)$$

One can prove the following

Lemma 10 The deformations (13) given by the pair f'_1, f'_2 of constraints and by the pair e_1, e_2 of constraints define the same reduced Poisson operator π_R on S .

Proof. For the moment we denote the deformation (13) defined through f'_1, f'_2 by π_D and the corresponding deformation defined through e_1, e_2 by π_{e_D} . Applying (13) we easily get that for any two functions $A, B : M \rightarrow \mathbb{R}$

$$fA; B g_{\pi_D} = fA; B g + \frac{fA; f'_2 g \quad fB; f'_1 g \quad fA; f'_1 g \quad fB; f'_2 g}{f'_1; f'_2 g};$$

where we have assumed that $f'_1; f'_2 g$ does not vanish on S . Similarly

$$fA; B g_{\pi_{e_D}} = fA; B g + \frac{fA; e_2 g \quad fB; e_1 g \quad fA; e_1 g \quad fB; e_2 g}{f e_1; e_2 g}; \quad (21)$$

where $f e_1; e_2 g$ does not vanish on S due to (20). Using the relations (19) between the deformed constraints e_i and the original constraints f'_i , the Leibniz property of Poisson brackets and the fact that the functions f'_i vanish on S we obtain

$$f e_1; e_2 g = D f'_1; f'_2 g$$

and

$$\begin{aligned} & (fA; e_2 g \quad fB; e_1 g \quad fA; e_1 g \quad fB; e_2 g) \\ &= D (fA; f'_2 g \quad fB; f'_1 g \quad fA; f'_1 g \quad fB; f'_2 g) \end{aligned}$$

so that the nonzero term sD in the numerator and denominator of (21) cancel and we obtain $fA; B g_{\pi_D|_S} = fA; B g_{\pi_{e_D}|_S}$ which implies that the projections of π_D and π_{e_D} onto S coincide. ■

In this nonsingular case the distribution Z along which we project a Poisson tensor usually changes after reparametrization, but $Z|_S$ remains the same as can be easily demonstrated. Thus in case of the second class constraints one has a "canonical" way of projecting onto S .

We now suppose that the constraints f'_i are of first class, that is $f'_1; f'_2 g_{ij} = 0$ and that the singularity in D is not removable. We may still attempt to define the projection π_R by reparametrizing S as in (19) above. It turns out that among an infinite set of admissible reparametrizations there are some exceptional which, although they fulfill the condition (16), nevertheless eliminate the singularity in D . In this case, however, by choosing a new parametrization e_1, e_2 of S we change the distribution Z even on S so that we cannot expect that the projection π_R will be independent of the choice of the parametrization. We lose a natural, "canonical" choice of projection, but we still can perform the projection, although in infinitely many nonequivalent ways. We illustrate this below in a sequence of examples.

Consider a six-dimensional manifold M parametrized with coordinates $(q_1, q_2, q_3, p_1, p_2, p_3)$ with the following Poisson operator:

$$D = \begin{pmatrix} 2 & 0 & 0 & 0 & 1 & q_1 & 0 & 3 \\ 6 & 0 & 0 & 0 & q_1 & 2q_2 + 1 & q_3 & 7 \\ 6 & 0 & 0 & 0 & 0 & q_3 & 0 & 7 \\ 6 & 1 & q_1 & 0 & 0 & p_1 & 0 & 7 \\ 4 & q_1 & 2q_2 & 1 & q_3 & p_1 & 0 & 5 \\ & 0 & q_3 & 0 & 0 & p_3 & 0 & \end{pmatrix} :$$

Consider now a four-dimensional submanifold S in M given by the relations

$$f'_1(q;p) = q_3 = 0, \quad f'_2(q;p) = p_3 = 0. \quad (22)$$

It is clear that $f'_1; f'_2 g$ vanishes on the whole manifold M (and thus on S) so that these constraints do not define any Dirac deformation at all. We now deform (22) as

$$e_1 = f'_1 + f'_2; \quad e_2 = (p_2 - q_1 p_1) f'_1 + f'_2 \quad (23)$$

Calculation shows $f e_1; e_2 g = (p_3 - q_3) q_3$ so that $f e_1; e_2 g_{ij} = 0$. One can show that after introducing the Casimir variables $(q_1, q_2, e_1, p_1, p_2, e_2)$ the deformed operator D attains the form

$$D = \begin{pmatrix} 2 & 0 & 2 \frac{q_1 q_3}{q_3 p_3} & 0 & 1 & q_1 & 0 & 3 \\ 6 & 2 \frac{q_1 q_3}{q_3 p_3} & 0 & 0 & q_1 + 2 \frac{p_1 q_3}{q_3 p_3} & q_1^2 + & 0 & 7 \\ 6 & 0 & 0 & 0 & 0 & 0 & 0 & 7 \\ 6 & 0 & 0 & 0 & 0 & 0 & 0 & 7 \\ 6 & 1 & q_1 & 2 \frac{p_1 q_3}{q_3 p_3} & 0 & 0 & p_1 & 0 \\ 4 & q_1 & q_1^2 & 0 & p_1 & 0 & 0 & 5 \\ & 0 & 0 & 0 & 0 & 0 & 0 & \end{pmatrix} ;$$

where now $q_3 = q_3(q;p;e)$ and $p_3 = p_3(q;p;e)$ and $= (q_3 + p_2 q_3 + q_1 q_3 p_1) = (q_3 - p_3)$, and as such is clearly singular on S and thus unreducible. This situation seems to be the most common, i.e. a spontaneous choice of parametrization almost always leads to a singularity. However, if we perform a slightly different deformation of (22):

$$e_1 = f'_1; \quad e_2 = (p_2 - q_1 p_1) f'_1 + f'_2 \quad (24)$$

so that $f\mathbf{e}_1; \mathbf{e}_2 g = q_3^2$ is again zero on S , then the operator D becomes nonsingular and its projection on S has the following form

$$R = \begin{array}{ccccc} & 2 & & & 3 \\ & 0 & 0 & 1 & q_1 \\ \begin{array}{c} 6 \\ 6 \\ 4 \end{array} & 0 & 0 & q_1 & 1 & q_1^2 & \begin{array}{c} 7 \\ 7 \\ 5 \end{array} \\ & 1 & q_1 & 0 & p_1 & \\ & q_1 & q_1^2 & 1 & p_1 & 0 \end{array}$$

in the variables $(q_1; q_2; p_1; p_2)$. Yet another deformation (even this time of the form (19)):

$$\mathbf{e}_1 = q_2' \mathbf{e}_1, \quad \mathbf{e}_2 = (p_2 + q_2') \mathbf{e}_1 \quad (25)$$

yields a quite complicated expression on $f\mathbf{e}_1; \mathbf{e}_2 g$:

$$f\mathbf{e}_1; \mathbf{e}_2 g = (3q_2 + 1)q_3^2 + q_3^3;$$

so that it again vanishes on S , but D is again nonsingular and in the same variables $(q_1; q_2; p_1; p_2)$ its projection becomes

$$R = \begin{array}{ccccc} & 2 & & & 3 \\ & 0 & 0 & 1 & \frac{q_1^2}{3q_2 + 1} & 0 \\ \begin{array}{c} 6 \\ 6 \\ 4 \end{array} & 0 & 0 & \frac{q_1 q_2}{3q_2 + 1} & 0 & \begin{array}{c} 7 \\ 7 \\ 7 \end{array} \\ & \frac{q_1^2}{3q_2 + 1} & 1 & \frac{q_1 q_2}{3q_2 + 1} & 0 & \frac{q_1 p_2}{3q_2 + 1} \\ & 0 & 0 & \frac{q_1 p_2}{3q_2 + 1} & 0 & \end{array}$$

which concludes our series of examples.

4 Conclusions

In this article we have focused on two issues involving Dirac reductions of Poisson operators on submanifolds. In the first part of the article we have shown how the Dirac reduction procedure fits in a natural way, i.e. as a result of two natural assumptions about the deformation D of ω , in the general Marsden-Ratiu reduction scheme. In the second part of our considerations we have demonstrated that the Dirac reduction procedure is often possible even in cases when the constraints that define our submanifold are of first class (in Dirac terminology), possibly after some suitably chosen reparametrization of the submanifold S .

Acknowledgement One of the authors (M.B.) would like to thank Department of Science and Technology, Linköping University, for its kind hospitality during his stay in Sweden in the fall of 2002.

References

- [1] J. Marsden, T. Ratiu, Reduction of Poisson manifolds, Lett. Math. Phys. 11 (1986) 161{169.
- [2] I. Vaisman, Lectures on the Geometry of Poisson Manifolds. Progress in Math., vol. 118, Birkhäuser Basel, Boston, Berlin 1994.

- [3] G. Falqui, M. Pedroni, Separation of variables for bi-Hamiltonian systems, preprint nlin.SI/0204029 v1 15 April 2002.
- [4] M. Błaszak, From bi-Hamiltonian geometry to separation of variables: stationary Harry-Dym and the KdV dressing chain, J. Nonlin. Math. Phys. 9 Supplement 1 (2002) 1{13.
- [5] G. Falqui, F. Magri, G. Tondo, Reduction of bihamiltonian systems and separation of variables: an example from the Boussinesq hierarchy, Theor. Math. Phys. 122 (2000) 176{192.
- [6] P. J. Olver, Application of Lie groups to differential equations, Springer-Verlag New York, 1986.
- [7] P. A. M. Dirac, Generalized Hamiltonian Dynamics, Can. J. Math. 2 (1950) 129{148.