

Non-Vanishing Profiles for the Kuramoto-Sivashinsky Equation on the Infinite Line

April 16, 2024

G. van Baalen¹, J.-P. Eckmann^{1;2}

¹ Département de Physique Théorique, Université de Genève

² Section de Mathématiques, Université de Genève

Abstract

We study the Kuramoto-Sivashinsky equation on the infinite line with initial conditions having arbitrarily large limits $\lim_{x \rightarrow \pm\infty} u(x,0) = \pm 1$. We show that the solutions have the same limits for all positive times. This implies that an attractor for this equation cannot be defined in L^1 . To prove this, we consider profiles with limits at $x = \pm 1$, and show that initial conditions L^2 -close to such profiles lead to solutions which remain L^2 -close to the profile for all times. Furthermore, the difference between these solutions and the initial profile tends to 0 as $x \rightarrow \pm 1$, for any fixed time $t > 0$. Analogous results hold for L^2 -neighborhoods of periodic stationary solutions. This implies that profiles and periodic stationary solutions partition the phase space into mutually unattainable regions.

1 Introduction

The Kuramoto-Sivashinsky equation

$$u_t(x;t) = -u u_x(x;t) - \frac{1}{2} (u_x(x;t))^2; \quad u(x;0) = u_0(x); \quad (1.1)$$

is an interesting model for stabilization mechanisms of very indirect type. It can be considered on a finite interval of length L with periodic boundary conditions (KS_L) or on the infinite line (KS_1). In both cases, one can formally multiply the equation with u and integrate, leading to (after integration by parts)

$$\frac{1}{2} \partial_t \int_{-\infty}^{\infty} u^2 dx = - \int_{-\infty}^{\infty} u^3 dx - \int_{-\infty}^{\infty} u (u_x)^2 dx - \frac{1}{2} \int_{-\infty}^{\infty} (u_x)^3 dx.$$

Note that the last term vanishes identically since it equals $\int_{-\infty}^{\infty} (u_x)^3 dx = 0$ and thus, surprisingly, the non-linearity does not contribute directly to the decay of initial conditions with large L^2 norm, in contrast to equations like the Ginzburg-Landau equation [12] which derive from a potential.¹

¹On the other hand, as we shall see, it is precisely this feature which allows for L^2 bounds which grow only exponentially in time.

Clearly, on the other hand, without the non-linear term, the equation is unstable. It has been shown [14, 7, 3, 5] that for finite L , the equation KS_L has an attractor in L^2 as well as in L^1 , whose radius is finite (known to be bounded by $L^{8=5}$, resp. $L^{48=25}$, see *e.g.*, [3, 4]). Numerical experiments seem to indicate that these bounds should in fact be extensive (with at worst $O(\log L)$ corrections), *i.e.*, $L^{1=2}$ for the L^2 radius and L^0 for L^1 . Here, we will show that this conjecture is wrong for initial conditions in $L^1(\mathbf{R})$, since we shall construct initial data φ_Y satisfying $\lim_{x \rightarrow \pm 1} \varphi_Y(x) = \pm Y$ and for which the corresponding solution² satisfies

$$\lim_{t \rightarrow 1} \varphi(x; t) = \pm Y \quad (1.2)$$

for all $t > 0$. The behavior of (1.2) is similar to what happens for the diffusion equation, where initial conditions with different limits at ± 1 also maintain this property as time increases, see *e.g.*, [2]. Note that due to the Galilean invariance of (1.1), the extensivity conjecture $O(L^0)$ for the attractor in L^1 is trivially wrong without a requirement ruling out constants – which are global solutions of (1.1) – as admissible initial data. The usual restriction to break the Galilean symmetry is to require the initial datum to be an odd function (see *e.g.* [3]). For our purpose however, it is enough to assume that the limits at $x = \pm 1$ are of opposite sign and equal magnitude.

What do we learn from (1.2)? Basically, it shows, that if there is ever to be a definition of attractor for KS_1 it must contain constraints on the initial condition which are much stronger than just being in L^1 (plus asymmetry and arbitrary regularity). Rather, if there is any hope to define a bounded attractor, it would have to come from a condition which says that the initial condition looks everywhere “like” those well-known [9] patterns one encounters in numerical simulations. In the absence of a technique replacing localization as in [1, 13] this seems currently impossible to achieve. Note however that it is known [10] that periodic stationary solutions are universally bounded. The present paper obtains more information on the structure of the phase space, if not on an eventual attractor, by showing that periodic stationary solutions and profiles divide naturally the phase space into mutually unattainable regions.

Our proof of (1.2) is based on the following simple idea. First of all, constants are clearly stationary solution of KS_1 . Furthermore, (1.1) has a one parameter family of explicit (albeit unbounded) solutions of the form

$$\varphi(x; t) = \frac{bx}{1 + bt} ;$$

with $b > 0$, showing that positive constant slopes are rotated clockwise. Our starting point consists in combining these two special solutions by taking as an initial condition the function

$$\varphi_a(x) = a \arctan(x) ;$$

where $a = 2Y = \frac{1}{2}$. The ‘middle’ of this function is like the constant slope example (with $b = a$) while for large x it reaches very quickly $\pm Y$. It is therefore natural to assume that

$$\varphi_a(x; t) = a \arctan \frac{x}{1 + at}$$

²the well known Bunsen flame fronts (see [11]).

is a good approximate profile for $t > 0$. In fact, while the definition of $\varphi_a(x; t)$ is suggestive, as long as we only have bounds, and not small bounds in L^2 , we may, and will, work with the *fixed* function $\varphi_a = \varphi_a(x)$ that is, with a profile which is not changing in time. Our main result is that *if the initial condition φ_0 is L^2 -close to $\varphi_a(x)$ and the difference decays at infinity then the solution remains L^2 -close to $\varphi_a(x)$, and the difference still decays at infinity for all $t > 0$* . This result also holds if φ_a is replaced by φ_{per} , with φ_{per} a periodic and analytic stationary solution of (1.1) as constructed in [10]. Therefore, there exist solutions of KS_1 which stay near

φ_a at infinity for all times, and thus we have found a family of large initial conditions whose evolution does not get smaller in L^1 as time goes to ∞ .³ On the other hand, initial conditions which behave asymptotically like periodic stationary solutions apart from L^2 corrections remain so for all times. Since the difference of two periodic functions with different periods is *not* in L^2 , this shows that the phase space naturally splits into disconnected components. This last result is an extension of [6].

The discussion above suggests to consider the equation for $\varphi(x; t) = \varphi_a(x; t) + \varphi_a(x)$, which reads

$$\varphi_t = -\varphi \varphi_x - \frac{1}{2}(\varphi^2)_x + \varphi_a \varphi_{ax}; \quad \varphi(x; 0) = \varphi_0(x) \quad (1.3)$$

where

$$\varphi_a = \varphi_{\text{per}} + \varphi_0;$$

and $\lim_{x \rightarrow \pm \infty} \varphi_0(x) = 0$. We will consider (1.3) either with $\varphi_a = \varphi_a$ and corresponding φ_{ax} , or with $\varphi_a = \varphi_{\text{per}}$ a periodic analytic stationary solution for which $\varphi_{ax} = \varphi_{\text{per},x} = 0$. Instead of φ_a , we could have used the stationary profiles (*i.e.*, stationary solutions of (1.1)) constructed in [10], or even the explicit one

$$\varphi(x; t) = \frac{15}{361} \frac{p}{209} - 9 \tanh \frac{p}{38} x + 11 \tanh \frac{p}{38} x^3$$

found by Kuramoto [8]. Note that these profiles are uniformly bounded. The advantage of these choices would have been that $\varphi_a = 0$, the disadvantage is the lack of explicit formulas, in particular for the Fourier transform of the profiles. While adding an inhomogeneous term to the equation, the choice of φ_a retains the main properties of these stationary profiles, *e.g.* in terms of analyticity. As is easily seen, high frequency modes are strongly damped by (1.1) at the linear level. It is known (see *e.g.*, [4]) that solutions corresponding to periodic antisymmetric initial condition in $L^2([-\frac{1}{2}, \frac{1}{2}])$ become analytic in finite time in a strip of finite width around the real axis. The error term φ_a of the equation (see (1.3)) and φ_a are analytic in the strip $|\text{Im } z| < 1$ and uniformly bounded in any smaller strip—these two facts are better seen in Fourier space, since the Fourier transform $\tilde{\varphi}_a$ of φ_a exists as a distribution and is given by

$$\tilde{\varphi}_a(k) = \frac{e^{-k/2}}{k} : \quad (1.4)$$

³Note that we do not claim (and it quite probably is not true) that $\varphi(x; t) - \varphi_a(x; t)$ stays bounded in L^2 . We will rather see that it grows (quickly) in L^2 . But the only thing which matters is that it remains in L^2 and decays at infinity.

Definition 1.1 Throughout, we denote by A the operator $A = \mathcal{P} \overline{\partial_x^2}$.

Remark 1.2 We fix $\Upsilon > 0$, and we tacitly admit that all constants occurring in the sequel may depend on Υ .

Theorem 1.3 There are constants C and $\delta > 0$ such that the following holds. For any initial condition $(\cdot; 0)$ with $\|(\cdot; 0)\|_2 < 1$, the solution of (1.3) exists for all $t > 0$ and

$$\sup_{t \geq 0} e^{-\delta t} \|(\cdot; t)\|_2 \leq \|(\cdot; 0)\|_2 + C :$$

Furthermore, the flow is regularizing in the sense that there exist constants $\delta > 0$, $C < 1$ such that

$$\sup_{t \geq 0} e^{-\delta t} k e^{m \ln(\cdot; t)A} (\cdot; t) \|_2 \leq \|(\cdot; 0)\|_2 + C :$$

Corollary 1.4 For every $m = 0, 1, \dots$ there exists a constant C_m such that

$$\| \mathfrak{P}_x^m(x; t) \| \leq \frac{C_m}{m \ln(\cdot; t)^m} e^{-\delta t} \text{ and } \lim_{x \rightarrow 1} \mathfrak{P}_x^m(x; t) = 0$$

for all $t > 0$.

Proof. By Theorem 1.3 there exists a C^0 such that $k e^{m \ln(\cdot; t)A} (\cdot; t) \|_2 \leq C^0 e^{-\delta t}$. By the Schwarz inequality,

$$\|k A^m \sim(\cdot; t)\|_2 \leq \sum_{j=1}^m \|k \mathfrak{J}_j^m\|_2 e^{-\delta t} \leq \sum_{j=1}^m \|k \mathfrak{J}_j^m\|_2 e^{-\delta t} \leq C^0 e^{-\delta t} \sum_{j=1}^m \|k \mathfrak{J}_j^m\|_2 :$$

This immediately implies the first assertion since $\sup_x \|\mathfrak{P}_x^m(x; t)\|_2 \leq \|k A^m \sim(\cdot; t)\|_2$. This bound also implies (by the Riemann-Lebesgue theorem) that $\lim_{x \rightarrow 1} \mathfrak{P}_x^m(x; t) = 0$. ■

2 Functional Spaces, Estimates

In this section, we collect a few straightforward bounds on the function \mathfrak{a} and the operator $A = \mathcal{P} \overline{\partial_x^2}$. We denote by W_{δ}^p the (Banach) space obtained by completing $C_0([0, +\infty]; \mathcal{C}^1(\mathbf{R}))$ in the norm $\sup_{t \geq 0} \|\cdot\|_p e^{(\delta t)A}$, where, throughout, $\|\cdot\|_p$ is the L^p norm. We also denote by B_d the open ball of radius d , centered on 0 in W_{δ}^p . The bounds of this section serve to control the non-linear and mixed terms in Eq.(1.3).

Lemma 2.1 There is a $\delta > 0$ such that for all $t \geq [0, +\infty]$ one has

$$k e^{(\delta t)A} \mathfrak{a}(\cdot) f(\cdot; t) \|_2 \leq \| \mathfrak{a} \|_p \overline{k 1 + A^2 }^p e^{(\delta t)A} f(\cdot; t) \|_2 :$$

Remark 2.2 One can choose $\delta = 1$ as will be seen from the proof. (This value is related to the domain of analyticity of φ_a .)

Proof. Define $F(x; t) = e^{(t-A)x} \varphi_a(x) f(x; t)$. We write the Fourier transform of F as

$$\tilde{F}(k; t) = e^{(t-A)k} \int_{-\infty}^{\infty} e^{-ikx} \varphi_a(x) f(x; t) dx :$$

Using (1.4), we find (using principal values)

$$\tilde{F}(k; t) = a e^{(t-A)k} \int_{-\infty}^{\infty} \frac{e^{-ikx}}{k} f(x; t) dx :$$

Denote $g(x; t) = e^{(t-A)x} f(x; t)$, so that $f(x; t) = e^{-(t-A)x} g(x; t)$. Rearranging exponentials, we get

$$\begin{aligned} \tilde{F}(k; t) &= a \int_{-\infty}^{\infty} \frac{e^{-ikx}}{k} e^{(t-A)k} e^{-(t-A)x} g(x; t) dx \\ &= a \int_{-\infty}^{\infty} \frac{e^{-ikx} e^{(t-A)k}}{k} e^{-(t-A)x} g(x; t) dx : \end{aligned}$$

We decompose this as

$$\tilde{F}(k; t) = a \int_{-\infty}^{\infty} \frac{e^{-ikx} e^{(t-A)k}}{k} g(x; t) dx + a \int_{-\infty}^{\infty} G(k; x; t; (1 - e^{-(t-A)x})) g(x; t) dx ;$$

where

$$G(k; x; t; \delta) = \frac{e^{-ikx}}{k} e^{(t-A)k} e^{-(t-A)x} (1 - e^{-\delta x}) :$$

One checks easily, using the triangle inequality, that for $\delta > 0$,

$$|G(k; x; t; \delta)| \leq |G(k; x; t; 0)| e^{\frac{\delta}{2} |k|} \leq \frac{1}{1 + (k\delta)^2} :$$

Using $\int_{-\infty}^{\infty} dk (1 + k^2)^{-1} = \pi$, we get

$$\begin{aligned} \|F(\cdot; t)\|_2 &\leq \|a e^{(t-A)\cdot} f(\cdot; t)\|_2 + a \sup_{x \in \mathbb{R}} \|G(\cdot; x; t; (1 + \delta)^{-1})\|_2 \|g(\cdot; t)\|_2 \\ &\leq \frac{a}{2} \|e^{(t-A)\cdot} f(\cdot; t)\|_2 + a \sup_{x \in \mathbb{R}} \|G(\cdot; x; t; 0)\|_2 e^{\frac{\delta}{2} \|x\|_1} \|g(\cdot; t)\|_2 \\ &\leq \frac{a}{2} \left(\frac{1}{2} + \frac{\delta}{2} \right) \|e^{(t-A)\cdot} f(\cdot; t)\|_2 + \frac{a}{2} \frac{\delta}{1 + \delta^2} \|e^{(t-A)\cdot} f(\cdot; t)\|_2 \\ &\leq \frac{a}{2} \frac{\delta}{1 + \delta^2} \|e^{(t-A)\cdot} f(\cdot; t)\|_2 ; \end{aligned}$$

provided $\delta > 0$. The proof of Lemma 2.1 is complete. ■

Lemma 2.3 Let p_{per} be periodic of period L , let $q = \frac{2}{L}$ and assume that there exist constants C_{per} ; such that $\sum_{m \in \mathbb{Z}} |p_{\text{per},m}| e^{qjm} < C_{\text{per}}$, where $p_{\text{per},m}$ denotes the m -th Fourier coefficient of p_{per} , then

$$\|e^{(t-A)p_{\text{per}}} f(\cdot; t)\|_2 \leq C_{\text{per}} \|e^{(t-A)} f(\cdot; t)\|_2$$

for all $t \in [0; +\infty]$.

Proof. As in Lemma 2.1, we define $F(x; t) = e^{(t-A)p_{\text{per}}}(x)f(x; t)$. The Fourier transform of F satisfies

$$\tilde{F}(k; t) = e^{(t-A)jk} \sum_{m \in \mathbb{Z}} p_{\text{per},m} \tilde{f}(k - qm);$$

so that

$$\begin{aligned} \|F(\cdot; t)\|_2^2 &= \sum_{m \in \mathbb{Z}} \int_{\mathbb{Z}} |dk| e^{2(t-A)jk + qjm} |\tilde{f}(k)|^2 \\ &= \|e^{(t-A)p_{\text{per}}} f(\cdot; t)\|_2^2 = \sum_{m \in \mathbb{Z}} \int_{\mathbb{Z}} |dk| e^{(t-A)qjm} |\tilde{f}(k)|^2; \end{aligned}$$

which completes the proof of the lemma since $0 \leq t < \infty$. ■

Lemma 2.4 Let $\gamma > 0$ and $k(1 + A^2)e^{-A} \|f\|_2 + ke^{-A} \|g\|_2 + ke^{-A} \|h\|_2 < 1$. Then

$$\|dx e^{-A} f - e^{-A} (gh)^0\|_2^p \leq k(1 + A^2)e^{-A} \|f\|_2^p + ke^{-A} \|g\|_2^p + ke^{-A} \|h\|_2^p;$$

Proof. Set $F = e^{-A} f$, $G = e^{-A} g$ and $H = e^{-A} h$. Since $\|j\|_2 \leq \|j\|_1 + \|j\|_2 = 0$ by the triangle inequality, we have

$$\begin{aligned} \|dx e^{-A} f - e^{-A} (gh)^0\|_2^p &= \left\| \int_{\mathbb{Z}} dk i k \tilde{F}(k) - \int_{\mathbb{Z}} dk e^{(jk - k - j)j} \tilde{G}(k - j) \tilde{H}(j) \right\|_2^p \\ &\leq \frac{p}{1 + A^2} \|A \tilde{F}\|_2^p + k \|G\|_2^p + k \|H\|_2^p \\ &\leq \frac{p}{1 + A^2} \|F\|_2^p + k \|G\|_2^p + k \|H\|_2^p; \end{aligned}$$

where we used again $\int_{\mathbb{Z}} dk (1 + k^2)^{-1} = 1$. ■

The following proposition estimates how close ϕ_a is to a solution of KS_1 .

Proposition 2.5 Define $\phi_a(x) = a \arctan(x)$ and let $\phi_a = \sum_{a=0}^{\infty} \phi_a^{(0)}$. Then, for $0 < \frac{1}{2}$, one has

$$\sup_{t \in [0; 1]} \|e^{tA} \phi_a\|_2 \leq B;$$

for some B depending only on a .

Proof. The Fourier transform of $e^{tA} \sum_{a=0}^{\infty} \phi_a^{(0)}$ is of the form $i^3 a e^{tjk} k^3 e^{-jk}$ so that for $t \in [0; 1]$ we get

$$\|e^{tA} \sum_{a=0}^{\infty} \phi_a^{(0)}\|_2^2 \leq a^2 \int_{\mathbb{Z}} dk e^{2(1-t)jk} k^6 = O((1-t)^{-7});$$

A similar bound holds for $\sum_{a=0}^{\infty} \phi_a^{(0)}$. The term $e^{tA} \phi_a^{(0)}$ is bounded using Lemma 2.1 with $\gamma = \frac{1}{2}$ instead of 1 and $f = \phi_a^{(0)}$. This yields a bound on the square of the L^2 norm which is of the form $a^2 O(1) \int_{\mathbb{Z}} dk (1 + k^2) e^{2jk} e^{-2jk}$, and combining the bounds completes the proof. ■

3 The Local Cauchy Problem in L^2

In this section, we consider the local (in time) Cauchy problem

$$u_t = \sum_{|\alpha| \leq m} a_\alpha(x) \partial^\alpha u + b(x); \quad (x; t) = (x_0(x); t) \quad (3.1)$$

for (1.3) with $u_0 \in L^2$. We will show, using a contraction argument, that it is well posed on any time interval $t \in [0; t_0]$ with $t_0 = \min(C \|u_0\|_2^{-2})$.⁴ To this end, we construct the map $\mathcal{F}(u)$ defined by $\mathcal{F}(u) = v$, where v is the solution of

$$v_t = \sum_{|\alpha| \leq m} a_\alpha(x) \partial^\alpha v + b(x); \quad (x; t) = (x_0(x); t) \quad (3.2)$$

and show that if $\|u_0\|_2$ is sufficiently small ($\|u_0\|_2 \leq k_0$) then \mathcal{F} is a contraction in a ball of radius $> \|u_0\|_2$ in W . Namely, let $f = e^{(t-t_0)A} u_0$ and $g = e^{(t-t_0)A} u_0$. Multiplying (3.1) with $f e^{(t-t_0)A}$, integrating over the space variable and using the results of the preceding section, we have

$$\begin{aligned} \frac{1}{2} \partial_t \|f\|_2^2 &= \|A^{1/2} f\|_2^2 - \|A^{3/2} f\|_2^2 + \|A f\|_2^2 + \frac{p-1}{2} \|k(1+A^2) f\|_2 \|f\|_2 \|g\|_2 \\ &\quad + 2 \|a\|_2 \|A f\|_2 \|k\|_2 \frac{p}{1+A^2} \|f\|_2^2 + B \|f\|_2^2; \end{aligned} \quad (3.3)$$

The first term on the r.h.s. comes from the time derivative of the exponential $e^{(t-t_0)A}$, the second and third from the space derivatives of f . The next term uses Lemma 2.4, while the last two use Lemma 2.1 and Proposition 2.5, respectively. Then, we use the inequalities

$$\|A f\|_2 \leq \|k\|_2 \frac{p}{1+A^2} \|f\|_2; \quad \|A^{3/2} f\|_2^2 \leq \|k(1+A^2) f\|_2 \|f\|_2$$

and get

$$\begin{aligned} \frac{1}{2} \partial_t \|f\|_2^2 &= \|A^{1/2} f\|_2^2 - \|A^{3/2} f\|_2^2 + \|A f\|_2^2 + B \|f\|_2^2 \\ &\quad + \frac{p-1}{2} \|k\|_2 \|g\|_2 + 2 \|a\|_2 \|k(1+A^2) f\|_2 \|f\|_2; \end{aligned}$$

We also have

$$\begin{aligned} \|A^{1/2} f\|_2^2 &= \|f\|_2^{3/2} \|A^{1/2} f\|_2^{1/2} \leq \frac{3}{4} \|f\|_2^2 + \frac{1}{4} \|A^{1/2} f\|_2^2; \\ \|A f\|_2^2 &= \|f\|_2 \|A^2 f\|_2 \leq \frac{1}{2} \|f\|_2^2 + \frac{1}{2} \|A^2 f\|_2^2; \\ \|k(1+A^2) f\|_2 &\leq \frac{1}{2} \|X\|_2^2 + \frac{3}{2} \|A^2 f\|_2^2 + 2 \|A f\|_2^2 + \|f\|_2^2; \end{aligned}$$

for all $X_i, \mu_i > 0; i=1,2,3$. Using these inequalities with sufficiently small μ_i shows that there is a positive constant c such that

$$\partial_t \|f\|_2^2 \leq (c + \|k\|_2^2) \|f\|_2^2 + B^2;$$

⁴Note that by our choice of t_0 , we have $t \leq \frac{1}{2}$.

from which we get that for all \mathbf{x} in the ball of radius d in W , $F(\cdot)$ satisfies

$$\sup_{t \in [0; +\infty[} \|e^{(t-\tau)A} F(\cdot) - F(\cdot)\|_{k_2} \leq e^{(c+d^2)=2} \frac{Q}{k_0 k_2^2 + B^2} :$$

For all $d > \frac{P}{k_0 k_2^2 + 2B^2}$, there exists a $\alpha = O(k_0 k_2^2)$ such that F maps B_d into itself. On the other hand $F' = F(\cdot_1) - F(\cdot_2)$ satisfies

$$F' = \sum_{i=1}^m \sum_{j=1}^m \frac{1}{2} (F')^0_{ij} \frac{1}{2} ((\cdot_1 - \cdot_2))^0_{ij} (F')^0_a; \quad F'(\mathbf{x}; \cdot) = 0;$$

where $\cdot = (\cdot_1 + \cdot_2)/2$ and $\cdot = (F(\cdot_1) + F(\cdot_2))/2$. Since \cdot_1 and \cdot_2 are in B_d in W , similar arguments also show that for the same d and α as above

$$\sup_{t \in [0; +\infty[} \|e^{(t-\tau)A} (F(\cdot_1) - F(\cdot_2))\|_{k_2} < \sup_{t \in [0; +\infty[} \|e^{(t-\tau)A} (\cdot_1 - \cdot_2)\|_{k_2};$$

so that F is a contraction in B_d in W . Thus the sequence of approximating solutions $\mathbf{x}_{n+1} = F(\mathbf{x}_n)$ converges to a unique solution of (3.1) in B_d in W .

Note that the results of this section also hold for the equation

$$\mathbf{x}' = \sum_{i=1}^m \sum_{j=1}^m \frac{1}{2} (\mathbf{x}')^0_{ij} (\mathbf{x}_{\text{per}})^0_{ij}; \quad (\mathbf{x}; \cdot) = \mathbf{x}_0(\mathbf{x}) : \quad (3.4)$$

It follows easily from [10] (see also [6]) that periodic stationary solutions \mathbf{x}_{per} of (1.1) satisfy the hypotheses of Lemma 2.3. The procedure is then exactly the same, *i.e.*, to show that the analog of the map F is a contraction in B_d in W , for some d ; and \cdot . Using obvious notations, we find that (3.3) is replaced by

$$\frac{1}{2} \partial_t \|f\|_{k_2}^2 = \|A^{1/2} f\|_{k_2}^2 - \|A^{1/2} f\|_{k_2}^2 + \|A f\|_{k_2}^2 + \frac{P}{2} k(1 + A^2) \|f\|_{k_2} \|f\|_{k_2} \|g\|_{k_2} + c_{\text{per}} \|A f\|_{k_2} \|f\|_{k_2};$$

from which it follows that

$$\partial_t \|f\|_{k_2}^2 \leq (c + k \|g\|_{k_2}^2) \|f\|_{k_2}^2;$$

for some positive c . The remainder of the proof is straightforward.

4 Proof of Theorem 1.3

Let now $\mathbf{x}_0 = 0$ and $(\mathbf{x}; \cdot_0) = \mathbf{x}_0(\mathbf{x})$ with $\mathbf{x}_0 \in L^2$, and define $\mathbf{x}_0 = k_0 k_2$. From the results of the preceding section, we know that there exists a unique solution of (3.1) in B_{d_0} in $W_{0;0}$ for $d_0 = D_0$ and $\mathbf{x}_0 = C_0^{-2}$, with C_0 so small that $\mathbf{x}_0 < \cdot$. Let $\mathbf{x}_1 = \mathbf{x}_0 + \cdot_0$. By the definition of W , (see also Corollary 1.4), \mathbf{x}_1 and all its derivatives tend to 0 as $\mathbf{x}_R \rightarrow 1$ for all $t \in [0; 1]$. In particular, the trilinear form $(\cdot_2)^0$ satisfies $(\cdot_2)^0 = \frac{1}{3} (\cdot_3)^0 = 0$. Hence, multiplying (1.3) with \cdot_1 and integrating over the space variable, we get:

$$\begin{aligned} \frac{1}{2} \partial_t \|\cdot_1\|^2 &= \sum_{i=1}^m \sum_{j=1}^m \frac{1}{2} (\cdot_1)^0_{ij} (\cdot_1)^0_{ij} - \sum_{i=1}^m \sum_{j=1}^m \frac{1}{2} (\cdot_1)^0_{ij} (\cdot_1)^0_{ij} \\ &= (\cdot_1)^0_{11} + (\cdot_1)^0_{22} - \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^m (\cdot_1)^0_{ij} (\cdot_1)^0_{ij} : \end{aligned}$$

Note that because the trilinear form vanishes, we get only quadratic (and linear) terms in \mathbf{z} , and therefore it is natural to find an exponential bound in time for the evolution of the L^2 norm; this is the main explanation for the bounds which follow below. Using $\|\mathbf{k}\|^4 + \|\mathbf{k}\|^2 \leq \frac{1}{4}$, we have the inequalities

$$\begin{aligned} \|\mathbf{z}\| &\leq \|\mathbf{z}\|^2 + \|\mathbf{z}\| \leq \frac{1}{4} \|\mathbf{z}\|^2 + \|\mathbf{z}\|; \\ \|\mathbf{z}\| &\leq \|\mathbf{z}\| \leq \frac{1}{2} \|\mathbf{z}\|^2 + \frac{1}{2} \|\mathbf{z}\|^2 : \end{aligned}$$

We find for, $t \in [0; 1]$, with $2(\frac{3}{4} + a)$,

$$\frac{1}{2} \|\mathbf{z}\|_t^2 \leq \frac{1}{2} \|\mathbf{z}\|^2 + \frac{1}{2} \|\mathbf{z}\|_a^2 \leq \frac{1}{2} \|\mathbf{z}\|^2 + \frac{B^2}{2} : \quad (4.1)$$

This differential inequality is valid for all $t \in [0; 1]$, and implies that

$$\|\mathbf{k}(\mathbf{i}_1)\mathbf{k}_2\| \leq e^{\frac{1}{2}} \sqrt{\frac{1}{4} + B^2} : \quad 1 :$$

Again, from the results of the preceding section, we now see that there exists a unique solution of (3.1) in $B_{d_1} \cap W_{1,1}$ for $d_1 = D_{-1}$ and $1 = C_{-1}^2$. Thus (4.1) is valid for all $t \in [0; 2]$ with $2 = 1 + 1$, and we get

$$\|\mathbf{k}(\mathbf{i}_2)\mathbf{k}_2\| \leq e^{\frac{1}{2}} \sqrt{\frac{1}{4} + B^2} : \quad 2 :$$

Continuing by induction, we find

$$\|\mathbf{k}_n\| = C_{-n}^2 = \frac{e^{-n}}{\frac{1}{4} + B^2} ;$$

so that

$$\|\mathbf{k}_{n+1}\| = \|\mathbf{k}_n\| + E e^{-n} ;$$

with $E = (\frac{1}{4} + B^2)^{-1}$. This implies that $\lim_{n \rightarrow \infty} \|\mathbf{k}_n\| = 1$, and therefore (4.1) is valid for all $t > 0$. This completes the proof of Theorem 1.3 for the profile case. The periodic case follows along the same lines.

Acknowledgements

This research was partially supported by the Fonds National Suisse.

References

- [1] P. Collet and J.-P. Eckmann. The time dependent amplitude equation for the Swift-Hohenberg problem. *Comm. Math. Phys.* **132** (1990), 139–153.

- [2] P. Collet and J.-P. Eckmann. Space-time behaviour in problems of hydrodynamic type: a case study. *Nonlinearity* **5** (1992), 1265–1302.
- [3] P. Collet, J.-P. Eckmann, H. Epstein, and J. Stubbe. A global attracting set for the Kuramoto-Sivashinsky equation. *Comm. Math. Phys.* **152** (1993), 203–214.
- [4] P. Collet, J.-P. Eckmann, H. Epstein, and J. Stubbe. Analyticity for the Kuramoto-Sivashinsky equation. *Phys. D* **67** (1993), 321–326.
- [5] J. Goodman. Stability of the Kuramoto-Sivashinsky and related systems. *Comm. Pure Appl. Math.* **47** (1994), 293–306.
- [6] Z. Grujić. Spatial analyticity on the global attractor for the Kuramoto-Sivashinsky equation. *J. Dynam. Differential Equations* **12** (2000), 217–228.
- [7] J. S. Il'yashenko. Global analysis of the phase portrait for the Kuramoto-Sivashinsky equation. *J. Dynam. Differential Equations* **4** (1992), 585–615.
- [8] Y. Kuramoto and T. Tsuzuki. Persistent propagation of concentration waves in dissipative media far from thermal equilibrium. *Progress of Theor. Phys.* **55** (1976), 356–368.
- [9] P. Manneville. The Kuramoto-Sivashinsky equation: a progress report. In: *Propagation in systems far from equilibrium (Les Houches, 1987)*, volume 41 of *Springer Ser. Synergetics* (Berlin: Springer, 1988), pp. 265–280.
- [10] D. Michelson. Steady solutions of the Kuramoto-Sivashinsky equation. *Phys. D* **19** (1986), 89–111.
- [11] D. Michelson. Stability of the Bunsen flame profiles in the Kuramoto-Sivashinsky equation. *SIAM J. Math. Anal.* **27** (1996), 765–781.
- [12] A. Mielke. The Ginzburg-Landau equation in its role as a modulation equation. In: *Handbook of dynamical systems, Vol. 2* (Amsterdam: North-Holland, 2002), pp. 759–834.
- [13] A. Mielke and G. Schneider. Attractors for modulation equations on unbounded domains—existence and comparison. *Nonlinearity* **8** (1995), 743–768.
- [14] B. Nicolaenko, B. Scheurer, and R. Temam. Some global dynamical properties of the Kuramoto-Sivashinsky equations: nonlinear stability and attractors. *Phys. D* **16** (1985), 155–183.