

ALGEBRO-GEOMETRIC SOLUTION OF THE DISCRETE KP EQUATION OVER A FINITE FIELD OUT OF A HYPERELLIPTIC CURVE

MARIUSZ BIAŁECKI AND ADAM DOLIWA

ABSTRACT. We transfer the algebro-geometric method of construction of solutions of the discrete KP equation to the finite field case. We emphasize role of the Jacobian of the underlying algebraic curve in construction of the solutions. We illustrate in detail the procedure on example of a hyperelliptic curve.

1. INTRODUCTION

Cellular automata are dynamical systems on a lattice with values being discrete (usually finite) as well. They are one of the more popular and distinctive classes of models of complex systems. Introduced in various contexts [28, 23] around 1950 they have found wide applications in different areas, from physics through chemistry and biology to social sciences [29].

One of the most interesting properties of cellular automata is that complex patterns can emerge from very simple update rules. However, usually one cannot easily predict how a given cellular automaton will behave without going through a number of time steps on a computer. Due to their completely discrete nature, cellular automata are naturally suitable for computer simulations, but also here it would be instructive to have examples of rules with large classes of analytical solutions, integrals of motions and other "integrable features".

The problem of construction of integrable cellular automata is not new and was undertaken in a number of papers (see, for example, [8, 4, 5, 26]). In particular in [27, 18] it was given a systematic method, called ultra-discretization, of obtaining cellular automaton version of a given discrete integrable system.

Recently a new approach to integrable cellular automata was proposed in [7]. Its main idea is to keep the form of a given integrable discrete system but to transfer the algebro-geometric method of construction of its solutions [14, 1] from the complex field \mathbb{C} to a finite field. This method, which in principle can be applied to any integrable discrete system with known algebro-geometric method of solution, has been applied to the fully discrete 2D Toda system (the Hirota equation) and in [3] to discrete KP and KdV equations (in Hirota form). In particular, the finite field valued multisoliton solutions of these equations have been constructed. We remark that the algebraic geometry over finite fields, although conceptually similar to that over the field of

2000 *Mathematics Subject Classification.* 14H70, 37K10, 37B15.

Key words and phrases. Integrable systems; cellular automata; finite fields; algebraic curves; discrete KP equation.

complex numbers [10], has its own tools and peculiarities [6, 21]. It is also nowadays very important in practical use in modern approaches to public key cryptography [13] and in the theory of error correcting codes [25].

The aim of this paper is to study in the finite field context the very distinguished example of integrable system — the discrete KP equation. We present the algebro-geometric scheme of construction of its solutions in a finite field and we demonstrate its linearization on the level of the abstract Jacobi variety of the corresponding algebraic curve. We illustrate details of the construction on example of a hyperelliptic curve.

We remark that in [4] it was observed that the Lax representation of the discrete sine-Gordon equation of Hirota [11] has a meaning also when the field of complex numbers is replaced by a finite field of the form F_{p^2} , where p is a prime number. The authors of [4] showed also that, in principle, the corresponding integrals of motion can be calculated. Finally, we note that possibility of considering the soliton theory in positive characteristic has been anticipated in [22].

The organization of the paper is as follows. In Section 2 we first summarize the finite field version of Krichever's construction of solutions of the discrete KP equation, then we present its abstract Jacobian picture. In Section 3 we apply the method starting from an algebraic curve of genus two.

2. THE FINITE FIELD SOLUTION OF THE DISCRETE KP EQUATION OUT OF NONSINGULAR ALGEBRAIC CURVES

We first briefly recall algebro-geometric construction of solutions of the discrete KP equation over finite fields [7, 3]. We discuss in addition a possible degeneracy of the linear problem and its consequences. Then we present the Jacobian picture of the construction, in which the integrable nature of the equation is evident. We point out some aspects of the representation which will help us to construct effectively solutions of the equation.

2.1. General construction. Consider an algebraic projective curve $C=\mathbb{K}$ (or simply C), absolutely irreducible, nonsingular, of genus g , defined over the finite field $\mathbb{K} = F_q$ with q elements, where q is a power of a prime integer p (see, for example [25, 10]). By $C(\mathbb{K})$ we denote the set of \mathbb{K} -rational points of the curve. By $\overline{\mathbb{K}}$ denote the algebraic closure of \mathbb{K} , i.e., $\overline{\mathbb{K}} = \bigcup_{i=1}^{\infty} F_{q^i}$, and by $C(\overline{\mathbb{K}})$ denote the corresponding infinite set (often identified with C) of $\overline{\mathbb{K}}$ -rational points of the curve. The action of the Galois group $G(\overline{\mathbb{K}}=\mathbb{K})$ (of automorphisms of $\overline{\mathbb{K}}$ which are identity on \mathbb{K} , see [17]) extends naturally to action on $C(\overline{\mathbb{K}})$.

Let us choose:

1. four points $A_i \in C(\mathbb{K})$, $i=0;1;2;3$,
2. effective \mathbb{K} -rational divisor of order g , i.e., g points $B_j \in C(\overline{\mathbb{K}})$, $j=1; \dots; g$, which satisfy the following \mathbb{K} -rationality condition

$$\sum_{j=1}^g \sigma_j(B_j) = \sum_{i=0}^3 A_i \quad (\sigma_j \in G(\overline{\mathbb{K}}=\mathbb{K})); \quad (B_j) = B_j \circ:$$

As a rule we assume here that all the points used in the construction are distinct and in general position. In particular, the divisor $\sum_{j=1}^g B_j$ is non-special.

Definition 1. Fix K -rational local parameter t_0 at A_0 . For any integers $n_1; n_2; n_3 \in \mathbb{Z}$ define the function $\psi(n_1; n_2; n_3)$ as a rational function on the curve C with the following properties:

1. it has pole of the order at most $n_1 + n_2 + n_3$ at A_0 ,
2. the first nontrivial coefficient of its expansion in t_0 at A_0 is normalized to one,
3. it has zeros of order at least n_i at A_i for $i = 1; 2; 3$,
4. it has at most simple poles at points $B_j, j = 1; \dots; g$.

As usual, zero (pole) of a negative order means pole (zero) of the corresponding positive order. Correspondingly one should exchange the expressions "at most" and "at least" in front of the orders of poles and zeros. By the standard (see e.g., [1]) application of the Riemann–Roch theorem (and the general position assumption) we conclude that the wave function $\psi(n_1; n_2; n_3)$ exists and is unique. The function $\psi(n_1; n_2; n_3)$ is K -rational, which follows from K -rationality conditions of sets of points in their definition.

Remark. In what follows we will often normalize functions in a sense of point 2 of Definition 1.

Fix K -rational local parameters t_i at $A_i, i = 1; 2; 3$. In the generic case, which we assume in the sequel, when the order of the pole of $\psi(n_1; n_2; n_3)$ at A_0 is $n_1 + n_2 + n_3$ denote by $\psi_k^{(i)}(n_1; n_2; n_3), i = 0; 1; 2; 3$, the K -rational coefficients of expansion of $\psi(n_1; n_2; n_3)$ at A_i , respectively, i.e.,

$$\psi(n_1; n_2; n_3) = \frac{1}{t_0^{(n_1+n_2+n_3)}} \sum_{k=1}^{\infty} \psi_k^{(0)}(n_1; n_2; n_3) t_0^k ;$$

$$\psi(n_1; n_2; n_3) = \sum_{k=0}^{\infty} t_1^{n_i} \psi_k^{(i)}(n_1; n_2; n_3) t_1^k ; \quad i = 1; 2; 3;$$

Denote by T_i the operator of translation in the variable $n_i, i = 1; 2; 3$, for example $T_1 \psi(n_1; n_2; n_3) = \psi(n_1 + 1; n_2; n_3)$. Uniqueness of the wave function implies the following statement.

Proposition 1. *Generically, the function ψ satisfies equations*

$$(1) \quad T_i \psi - T_j \psi + \frac{T_j \psi_0^{(i)}}{\psi_0^{(i)}} = 0; \quad i \neq j; \quad i; j = 1; 2; 3;$$

Remark. When the genericity assumption fails then the linear problem (1) degenerates, i.e., some of its terms are absent.

Notice that equation (1) gives

$$(2) \quad \frac{T_j \psi_0^{(i)}}{\psi_0^{(i)}} = \frac{T_i \psi_0^{(j)}}{\psi_0^{(j)}}; \quad i \neq j; \quad i; j = 1; 2; 3;$$

Define

$$(3) \quad \psi_i = \sum_{j < i}^P \psi_0^{(j)}; \quad i = 1; 2; 3;$$

then equation (2) implies existence of a K -valued potential (the τ -function) defined (up to a multiplicative constant) by formulas

$$(4) \quad \frac{T_i}{T_{i-1}} = \tau_i; \quad i = 1; 2; 3:$$

Finally, equations (1) give rise to condition

$$(5) \quad \frac{T_{2-1}}{T_1} - \frac{T_{3-1}}{T_1} + \frac{T_{3-2}}{T_2} = 0;$$

which written in terms of the τ -function gives the discrete KP equation [12] called also the Hirota equation

$$(6) \quad (\tau_1 - \tau_2)(\tau_2 \tau_3 - \tau_1 \tau_3) - (\tau_2 - \tau_3)(\tau_3 \tau_1 - \tau_1 \tau_2) + (\tau_3 - \tau_1)(\tau_1 \tau_2 - \tau_2 \tau_3) = 0:$$

Corollary 2. Equation (5) can be obtained also from expansion of equation (1) at A_k , where $k = 1; 2; 3$, $k \notin i; j$.

Remark. Absence of a term in the linear problem (1) (see the remark after Proposition 1) reflects, due to Corollary 2, in absence of the corresponding term in equation (6). This implies that in the non-generic case, when we have not defined the τ -function yet, we are forced to put it to zero. Let us notice in advance (see the next section) that it is in complete analogy with the well known (complex field) interpretation of the algebro-geometric solution of the discrete KP equation as, essentially, the Riemann theta function.

2.2. The Jacobian interpretation. Denote by $D \text{div}(C)$ the abelian group of the divisors on the curve C and by $P \text{ic}^0(C)$ the group of equivalence classes of degree zero divisors $D \text{div}^0(C)$ modulo the principal divisors. There exists [16, 20] an abelian variety $J(C)$ of dimension g (the Jacobian of the curve) and an injective map $\alpha: C \rightarrow J(C)$ (the Abel map) such that the extension of α to $D \text{div}(C)$ establishes an isomorphism between $P \text{ic}^0(C)$ and $J(C)$. Moreover, if there exists a K -rational point $A \in C(K)$ of the curve, then α can be defined by

$$\alpha(P) = \mathbb{P}^1 - A \in J(C);$$

where $\mathbb{P}^1 - A$ designates the class of the degree zero divisor $P - A$ in $P \text{ic}^0(C)$.

Denote by $D_r(C)$ the effective divisors of degree r of the curve C and by $\mathbb{P}^1 - A$ the extension of α to $D_r(C)$

$$D_r(C) \ni D \in \mathbb{P}^1 - A \in J(C):$$

The direct image of $\mathbb{P}^1 - A$ is a subvariety W_r of dimension r if $0 \leq r \leq g$, and of dimension g if $r > g$. In particular, W_{g-1} defines a divisor in $J(C)$.

The group $P \text{ic}^0(C; K)$ of equivalence classes of K -rational degree zero divisors $D \text{div}^0(C; K)$ modulo the principal K -rational divisors can be identified with the abelian group $J(C; K)$ of K -rational points of the Jacobian variety. For finite field K the group $P \text{ic}^0(C; K)$ is finite as well and its order can be found using properties of the zeta function of the curve (see, for example [25]).

Let us present in this picture the description of the wave function and of the function. We choose the point A_0 as the reference point A and consider the following divisor $D(n_1; n_2; n_3) \in \text{Div}^0(C; K)$ of degree zero

$$D(n_1; n_2; n_3) = n_1(A_0 - A_1) + n_2(A_0 - A_2) + n_3(A_0 - A_3) + \sum_{i=1}^{g-1} (B_i - A_i);$$

with linear dependence on n_1, n_2 and n_3 . Its equivalence class in $\text{Pic}^0(C; K)$ has the unique K -rational representant of the form

$$X(n_1; n_2; n_3) = \sum_{i=1}^{g-1} X_i(n_1; n_2; n_3) - A_i;$$

This equivalence is given by a function whose divisor reads

$$(7) \quad n_1(A_1 - A_0) + n_2(A_2 - A_0) + n_3(A_3 - A_0) + \sum_{i=1}^{g-1} X_i(n_1; n_2; n_3) - \sum_{i=1}^{g-1} B_i = 0;$$

If we normalize such a function at A_0 according to Definition 1 it becomes the wave function ψ . Notice that some of X_i could be A_0 which would mean that $D(n_1; n_2; n_3) \in W_{g-1}$. This correspondence gives rise to a set of important facts.

Corollary 3. *Evolutions in variables n_1, n_2 and n_3 define linear flows in the Jacobian.*

Corollary 4. *Points $X_i, i = 1, \dots, g$ indicate zeros of the wave function which are not specified in the previous construction.*

Corollary 5. *If $D(n_1; n_2; n_3) \in W_{g-1}$ then the pole of the wave function at A_0 has the order less than $(n_1 + n_2 + n_3)$, i.e., we are in the non-generic case, thus $\psi(n_1; n_2; n_3) = 0$.*

Remark. Notice that because $D(0; 0; 0) \notin W_{g-1}$ then $\psi(0; 0; 0) \neq 0$.

Remark. If $K = \mathbb{C}$ then the algebraic curve C is the compact Riemann surface, theorems of Abel and Jacobi identify the Jacobian with quotient of C^g by the period lattice, and a theorem of Riemann identifies W_{g-1} with a certain translate of the zero locus of the Riemann theta function (see [9]). Then, as we mentioned in remark after Corollary 2, the algebro-geometric solution of the discrete KP equation becomes, with appropriate understanding of its argument via the divisor $D(n_1; n_2; n_3)$ and up to a not essential and non-vanishing multiplier, the Riemann theta function (see, e.g. [15]). In particular, in such an interpretation the zeros of ψ are located in points of the translate W_{g-1} of the Theta divisor.

Let us discuss periodicity of solutions of the finite field version of the KP equation obtained using the above method. Denote by $\nu_i, i = 1; 2; 3$, the ranks of cyclic subgroups of $J(C; K)$ generated by divisors $A_i - A_0$, then for arbitrary $k_i \in \mathbb{Z}, i = 1; 2; 3$,

$$D(n_1 + k_1; n_2 + k_2; n_3 + k_3) - D(n_1; n_2; n_3) = 0;$$

In particular, $\psi(n_1; n_2; n_3) = 0$ implies $\psi(n_1 + k_1; n_2 + k_2; n_3 + k_3) = 0$.

There exist unique (normalized at A_0) functions h_i , $i = 1; 2; 3$, with zeros of order ν_i at A_i , poles of order μ_i at A_0 and no other singularities and zeros such that

$$(8) \quad (n_1 + k_1 - \mu_1; n_2 + k_2 - \mu_2; n_3 + k_3 - \mu_3) = h_1^{k_1} h_2^{k_2} h_3^{k_3} (n_1; n_2; n_3):$$

Remark. Generalizing above considerations, if for $\nu_i \in \mathbb{Z}$, $i = 1; 2; 3$,

$$X^3 \prod_{i=1}^3 (A_i - A_0)^{\nu_i} = 0;$$

then $(\nu_1; \nu_2; \nu_3)$ is the period vector of zeros of the X -function and vector of quasi-periodicity (in the above sense) of the wave function.

Equation (8) implies quasi-periodicity of the functions $\psi_0^{(i)}$, $i = 1; 2; 3$,

$$\psi_0^{(i)}(n_1 + k_1 - \mu_1; n_2 + k_2 - \mu_2; n_3 + k_3 - \mu_3) = c_{(i)1}^{k_1} c_{(i)2}^{k_2} c_{(i)3}^{k_3} \psi_0^{(i)}(n_1; n_2; n_3);$$

with the (non-zero) factors $c_{(i)j} \in \mathbb{K}$ equal to

$$c_{(i)j} = \frac{h_j}{t_j^{\nu_j}} \Big|_{P=A_i} :$$

The multiplicative group \mathbb{K} is a cyclic group of order $q - 1$, therefore the functions $\psi_0^{(i)}$ are periodic. Their periods in variable n_j are equal to ν_j times the order of the subgroup of \mathbb{K} generated by $c_{(i)j}$ (a divisor of $q - 1$). Due to possible change of sign (see equation (3)) the periods of $\psi_0^{(i)}$ can be eventually doubled with respect to the corresponding periods of $\psi_0^{(i)}$. Again, periodicity of $\psi_0^{(i)}$ implies quasi-periodicity of ψ with a factor from \mathbb{K} , thus the period of ψ in variable n_i can be maximally $q - 1$ times the period of $\psi_0^{(i)}$ in that variable.

3. A "HYPERELLIPTIC" SOLUTION OF THE DISCRETE KP EQUATION

Our goal here is to demonstrate how does the method described above work. We perform all steps of the construction (see also [2] for details) starting from a given algebraic curve, which we have chosen to be a hyperelliptic curve, due to relatively simple description of Jacobians of such curves [19]. We consider a hyperelliptic curve of genus $g = 2$ but the technical tools used here can be applied directly to hyperelliptic curves of arbitrary genus.

3.1. A hyperelliptic curve and its Jacobian. Consider a hyperelliptic curve C of genus $g = 2$ defined over the field \mathbb{F}_7 and given by the equation

$$(9) \quad C : v^2 + uv = u^5 + 5u^4 + 6u^2 + u + 3:$$

The $(u; v)$ coordinates of its \mathbb{F}_7 -rational points are presented in Table 1. The curve has one point at infinity, denoted by 1_∞ , whose preimage on the nonsingular model of C consists of one point only [24], and where the local uniformizing parameter can be chosen as $u^2 = v$ (u is a polynomial function of order 2, and v is a polynomial function of order 5). The point opposite (with respect to the hyperelliptic automorphism) to P is denoted by \bar{P} . The only two special point of the curve are $(6; 4)$ and the infinity point 1_∞ .

i	P_i	\tilde{P}_i
0	1	P_0
1	(1;1)	(1;5)
2	(2;2)	(2;3)
3	(5;3)	(5;6)
4	(6;4)	P_4

TABLE 1. F_7 -rational points of the curve C .

i	P_i	\tilde{P}_i	P_i	\tilde{P}_i
5	(0;21)	(0;28)	\tilde{P}_5	P_5
6	(3;9)	(3;44)	\tilde{P}_6	P_6
7	(4;26)	(4;33)	\tilde{P}_7	P_7
8	(7;5)	(7;44)	(42;5)	(42;9)
9	(8;22)	(8;26)	(43;29)	(43;33)
10	(11;5)	(11;47)	(46;5)	(46;12)
11	(12;6)	(12;45)	(47;6)	(47;10)
12	(13;14)	(13;29)	(48;35)	(48;22)
13	(14;8)	(14;34)	(35;43)	(35;27)
14	(15;13)	(15;28)	(36;48)	(36;21)
15	(16;17)	(16;23)	(37;38)	(37;30)
16	(17;0)	(17;39)	(38;0)	(38;18)
17	(18;4)	(18;41)	(39;4)	(39;20)
18	(19;9)	(19;28)	(40;44)	(40;21)
19	(20;12)	(20;31)	(41;47)	(41;24)
20	(22;4)	(22;30)	(29;4)	(29;23)
21	(25;6)	(25;32)	(32;6)	(32;25)
22	(27;7)	(27;22)	(34;42)	(34;29)

TABLE 2. F_{49} -rational points of the curve C (which are not F_7 -rational); here \tilde{P} is the opposite of P , and \tilde{P} denotes its conjugate with respect to the lift of the Frobenius automorphism.

We identify the field F_{49} as the extension of F_7 by the polynomial $x^2 + 2$, i.e., $F_{49} = F_7[x]/(x^2 + 2)$. Let us introduce the following notation: the element $k \in F_{49}$ represented by the polynomial $x + \dots$ is denoted by the *natural* number $7 + \dots$. The Galois group $G(F_{49}/F_7) = \langle \text{id}, \sigma \rangle$, where σ is the Frobenius automorphism, acts on elements of $F_{49} \setminus F_7$ in the following way:

$$\sigma(k) = 7(7 + \dots) + \dots$$

The coordinates of F_{49} -rational points of the curve (which are not F_7 -rational) are presented in Table 2.

In the next step we find the group of the F_7 -rational points $J(\mathbb{C}; F_7)$ of the Jacobian of the curve. The number of its points can be found from the number of F_7 -rational and F_{49} -rational points of the curve by application of properties of the zeta function of the curve C (see for instance [25, 13]). In our case the curve has 8 F_7 -rational points and 74 F_{49} -rational points which implies the following form of the zeta function $(\mathbb{C}; T)$

$$(\mathbb{C}; T) = \frac{P(T)}{(1-T)(1-7T)}; \quad P(T) = 1 + 12T^2 + 49T^4;$$

The number $\# J(\mathbb{C}; F_7)$ of the F_7 -rational points of the Jacobian is equal to $P(1) = 62$, and therefore $J(\mathbb{C}; F_7)$ is the direct sum of cyclic groups of orders 31 and 2.

Let us choose the infinity point 1 as the basepoint. The group law in the Jacobian of a hyperelliptic curve can be intuitively described in a way which is a higher-genus analog of the well known addition operation for points of elliptic curves. We present here only its sketch for genus $g = 2$ and in the generic case of addition of two points of $J(\mathbb{C})$ with representants of the form

$$E_i = Q_i + R_i - 21; \quad i = 1; 2;$$

with all points distinct. If $E_3 = Q_3 + R_3 - 21$ is the representant of $[E_1 + E_2]$, i.e.,

$$E_1 + E_2 = (g) + E_3;$$

then

$$(10) \quad E_1 + E_2 + \tilde{E}_3 = 0;$$

where we have used the fact that for any point $P \in C(\overline{\mathbb{K}})$ of a hyperelliptic curve the divisor $P + \tilde{P} - 21$ is principal. Therefore, there exists a normalized polynomial function f of the order six, thus necessarily of the form

$$f = a + bu + cu^2 + du^3 + ev;$$

with divisor given by the left hand side of equation (10). Its zeros at Q_i and R_i , $i = 1; 2$, and the normalization condition allow to fix the coefficients and then to find two other zeros Q_3 and R_3 .

Geometrically, we are looking for two other intersection points of the cubic interpolating known four points with the hyperelliptic curve. In cases when some points of $E_1 + E_2$ are repeated, the interpolation step must be adjusted to ensure tangency to the curve with sufficient multiplicity. When divisors have less points then we consider the interpolating curve of lower degree (some intersection points are at infinity). Finally, the transition function g is the unique normalized function with the nominator equal to f and the denominator being the normalized polynomial function with the divisor $E_3 + \tilde{E}_3$.

The full description of the group $J(\mathbb{C}; F_7)$ is given in Table 3. The divisor $D_1 = P_1 - 1$ generates the subgroup of order 31 and the divisor $D_4 = P_4 - 1$ generates the subgroup of order 2. We present the reduced representants $[nD_1 + mD_4]_K$ of elements $[nD_1 + mD_4]$ of $J(\mathbb{C}; F_7)$, where $n \in \{0; 1; \dots; 30\}g$ and $m \in \{0; 1\}g$. Moreover we write down the transition functions $g_m(n)$ defined by the following divisor equation

$$(11) \quad [nD_1 + mD_4]_K + D_1 = (g_m(n)) + [(n+1)D_1 + mD_4]_K;$$

Also the transition functions for the sums $[nD_1 + mD_4]_k + D_4$ can be read off from Table 3. In particular, to find such a transition function (we call it $W(0;1)$) with $n = 29$ and $m = 0$, i.e.,

$$(12) \quad (1;5) + (1;5) + D_4 \quad 21 = (W(0;1)) + (12;6) + (47;6) \quad 21 ;$$

we make use of the fact that the analogous transition function for $n = 30$ and $m = 0$ is 1. Then

$$(13) \quad W(0;1) = g_0(29) \cdot [g_0(29)]^{-1} = \frac{2 + 3u + 4u^2 + v}{6 + 4u + u^2};$$

where in the last equality we have used equation (9) of the curve to get rid of v from the denominator.

3.2. Construction of the wave and functions. In order to find a solution of the discrete KP equation let us fix the following points of the curve C ,

$$A_0 = 1 ; \quad A_1 = (1;1); \quad A_2 = (2;2); \quad A_3 = (5;3);$$

with the uniformizing parameters

$$t_0 = u^2=v; \quad t_1 = u - 1; \quad t_2 = u - 2; \quad t_3 = u - 5;$$

and

$$B_1 = (12;6); \quad B_2 = (47;6):$$

Because

$$\begin{aligned} A_0 - A_1 &= 30D_1; & A_0 - A_2 &= 21D_1 + D_4; & A_0 - A_3 &= 17D_1 + D_4; \\ B_1 + B_2 &= 2A_0 - 29D_1 + D_4; \end{aligned}$$

then the points $X_1(n_1;n_2;n_3)$ and $X_2(n_1;n_2;n_3)$ where the wave function $(n_1;n_2;n_3)$ has additional zeros can be found from Table 3 and from equation

$$(14) \quad X_1(n_1;n_2;n_3) + X_2(n_1;n_2;n_3) \quad 21 = [nD_1 + mD_4]_k;$$

where $n \in \{0,1,\dots,30\}$ and $m \in \{0,1\}$ are given by

$$(15) \quad n = 29 + 30n_1 + 21n_2 + 17n_3 \pmod{31};$$

$$(16) \quad m = 1 + n_2 + n_3 \pmod{2};$$

Remark. Notice that the infinity point 1 is the Weierstrass point of the curve C , which violates the assumption of general position of points used in the construction of solutions of the discrete KP equation. This will not change the Jacobian picture of the construction but in some situations, which we will point out, will affect uniqueness of the wave function. We remark that such a choice is indispensable in reduction of the method from the discrete KP equation to the discrete KdV equation (see, for example [15, 3]).

To find effectively the wave function we will constraint the range of parameters from Z^3 to the parameters of the group of F_7 -rational points of the Jacobian. Let us introduce functions h_1 and h_4 corresponding to generators of the two cyclic subgroups

n	$\lfloor nD_1 \rfloor_r$	$g_0(n)$	$\lfloor nD_1 + D_4 \rfloor_r$	$g_1(n)$
0	0	1	(6;4) 1	1
1	(1;1) 1	1	(1;1) + (6;4) 21	$\frac{5+5u+3u^2+v}{6+4u+u^2}$
2	(1;1) + (1;1) 21	$\frac{u+5u^2+v}{(2+u)^2}$	(12;45) + (47;10) 21	$\frac{1+5u^2+v}{2+5u+u^2}$
3	(5;6) + (5;6) 21	$\frac{1+u+4u^2+v}{(2+u)(5+u)}$	(15;28) + (36;21) 21	$\frac{6u^2+v}{2+u^2}$
4	(2;3) + (5;3) 21	$\frac{2+4u^2+v}{5+4u+u^2}$	(7;44) + (42;9) 21	$\frac{5+u+v}{4+6u+u^2}$
5	(19;9) + (40;44) 21	$\frac{4u+2u^2+v}{5+5u+u^2}$	(11;5) + (46;5) 21	$\frac{6+6u+u^2+v}{3+6u+u^2}$
6	(22;4) + (29;4) 21	$\frac{5+2u+6u^2+v}{(2+u)(5+u)}$	(18;41) + (39;20) 21	$\frac{5+3u+5u^2+v}{5+3u+u^2}$
7	(2;3) + (5;6) 21	$\frac{5+6u+2u^2+v}{5+2u+u^2}$	(16;17) + (37;38) 21	$\frac{5+4u+4u^2+v}{3+u+u^2}$
8	(27;22) + (34;29) 21	$\frac{1+3u+2u^2+v}{1+u^2}$	(17;39) + (38;18) 21	$\frac{3+2u+u^2+v}{(5+u)(6+u)}$
9	(14;34) + (35;27) 21	$\frac{1+5u+v}{(1+u)(5+u)}$	(1;5) + (2;2) 21	6 + u
10	(2;2) + (6;4) 21	$\frac{3+5u+5u^2+v}{(5+u)^2}$	(2;2) 1	1
11	(2;3) + (2;3) 21	$\frac{6+u+6u^2+v}{3+2u+u^2}$	(1;1) + (2;2) 21	$\frac{4+2u^2+v}{3+5u+u^2}$
12	(13;14) + (48;35) 21	$\frac{3+6u+4u^2+v}{2+2u+u^2}$	(8;22) + (43;29) 21	$\frac{2+4u+v}{(2+u)(6+u)}$
13	(20;12) + (41;47) 21	$\frac{5u+u^2+v}{(1+u)(2+u)}$	(1;5) + (5;3) 21	6 + u
14	(5;3) + (6;4) 21	$\frac{6+5u+2u^2+v}{6+6u+u^2}$	(5;3) 1	1
15	(25;32) + (32;25) 21	$\frac{5+u^2+v}{6+6u+u^2}$	(1;1) + (5;3) 21	$\frac{u+5u^2+v}{(2+u)(6+u)}$
16	(25;6) + (32;6) 21	$\frac{6+5u+2u^2+v}{(1+u)(2+u)}$	(1;5) + (5;6) 21	6 + u
17	(5;6) + (6;4) 21	$\frac{5u+u^2+v}{2+2u+u^2}$	(5;6) 1	1
18	(20;31) + (41;24) 21	$\frac{3+6u+4u^2+v}{3+2u+u^2}$	(1;1) + (5;6) 21	$\frac{2+4u+v}{3+5u+u^2}$
19	(13;29) + (48;22) 21	$\frac{6+u+6u^2+v}{(5+u)^2}$	(8;26) + (43;33) 21	$\frac{4+2u^2+v}{(5+u)(6+u)}$
20	(2;2) + (2;2) 21	$\frac{3+5u+5u^2+v}{(1+u)(5+u)}$	(1;5) + (2;3) 21	6 + u
21	(2;3) + (6;4) 21	$\frac{1+5u+v}{1+u^2}$	(2;3) 1	1
22	(14;8) + (35;43) 21	$\frac{1+3u+2u^2+v}{5+2u+u^2}$	(1;1) + (2;3) 21	$\frac{3+2u+u^2+v}{3+u+u^2}$
23	(27;7) + (34;42) 21	$\frac{5+6u+2u^2+v}{(2+u)(5+u)}$	(17;0) + (38;0) 21	$\frac{5+4u+4u^2+v}{5+3u+u^2}$
24	(2;2) + (5;3) 21	$\frac{5+2u+6u^2+v}{5+5u+u^2}$	(16;23) + (37;30) 21	$\frac{5+3u+5u^2+v}{3+6u+u^2}$
25	(22;30) + (29;23) 21	$\frac{4u+2u^2+v}{5+4u+u^2}$	(18;4) + (39;4) 21	$\frac{6+6u+u^2+v}{4+6u+u^2}$
26	(19;28) + (40;21) 21	$\frac{2+4u^2+v}{(2+u)(5+u)}$	(11;47) + (46;12) 21	$\frac{5+u+v}{2+u^2}$
27	(2;2) + (5;6) 21	$\frac{1+u+4u^2+v}{(2+u)^2}$	(7;5) + (42;5) 21	$\frac{6u^2+v}{2+5u+u^2}$
28	(5;3) + (5;3) 21	$\frac{u+5u^2+v}{(6+u)^2}$	(15;13) + (36;48) 21	$\frac{1+5u^2+v}{6+4u+u^2}$
29	(1;5) + (1;5) 21	(6 + u)	(12;6) + (47;6) 21	$\frac{5+5u+3u^2+v}{(1+u)(6+u)}$
30	(1;5) 1	(6 + u)	(1;5) + (6;4) 21	6 + u

TABLE 3. The group $\mathcal{J}(C;F_7)$ of F_7 -rational points of the Jacobian as the simple sum of its cyclic subgroups.

of $J(\mathbb{C}; \mathbb{F}_7)$. The function h_1 with the divisor $31D_1$ and normalized at the infinity point is equal to the product $\prod_{i=0}^{30} g_0(i)$ and reads

$$h_1 = 1 + 2u + u^2 + 4u^3 + 3u^5 + u^6 + 3u^7 + u^8 + 4u^9 + 4u^{10} + 2u^{11} + 5u^{12} + 2u^{13} + 4u^{14} + 3u^{15} + 5u + 2u^2 + 5u^3 + 4u^5 + 6u^6 + 4u^7 + 3u^9 + 5u^{10} + 5u^{11} + 4u^{12} + u^{13} v;$$

where we also used equation of the curve (9) to reduce higher order terms in v . The normalized function h_4 with the divisor $2D_4$ is

$$h_4 = u - 6;$$

Let us introduce other auxiliary functions f_2 and f_3 to factorize the zeros of the wave function at A_2 and A_3 . Notice that

$$A_2 + 21D_1 + D_4 - 1 = 0;$$

which implies that there exists a polynomial function on \mathbb{C} with simple zero at A_2 and other zeros in the distinguished (by our choice of description of $J(\mathbb{C}; \mathbb{F}_7)$) points $(1;1)$ and $(6;4)$. Define f_2 as the unique such function normalized at the infinity point 1, then

$$f_2 = 1 + 5u + u^2 + 4u^4 + 6u^5 + 4u^6 + 4u^7 + 3u^8 + 4u^9 + 6u^{11} + 6 + 4u + 2u^2 + 5u^3 + 6u^4 + 6u^6 + u^7 + u^8 + u^9 v;$$

Similarly we define the normalized function

$$f_3 = 1 + 6u + 2u^2 + 6u^5 + u^6 + 5u^7 + 5u^8 + 4u^9 + 4 + 3u + 5u^2 + 4u^5 + 2u^6 + u^7 v;$$

with the divisor $A_3 + 17D_1 + D_4 - 1$.

Uniqueness of the wave function implies that it can be decomposed as follows

$$(17) \quad (n_1; n_2; n_3) = \frac{f_2^{n_2} f_3^{n_3}}{h_1^{p_1} h_4^q} W(m_1; m_2);$$

where $W(m_1; m_2)$ is the unique normalized function with the divisor

$$(18) \quad m_1 D_1 + m_2 D_4 + Y_1(m_1; m_2) + Y_2(m_1; m_2) - (12; 6) - (47; 6);$$

where

$$(19) \quad Y_1(m_1; m_2) + Y_2(m_1; m_2) = X_1(n_1; n_2; n_3) + X_2(n_1; n_2; n_3);$$

and the new variables m_1 i m_2 are given by

$$(20) \quad 21n_2 + 17n_3 - n_1 = 31p - m_1; \quad m_1 \in \mathbb{F}_0; 1; \dots; 30g;$$

$$(21) \quad n_2 + n_3 = 2q - m_2; \quad m_2 \in \mathbb{F}_0; 1g;$$

To find the functions $W(m_1; m_2)$ for all $m_1 \in \mathbb{F}_0; 1; \dots; 30g$ and $m_2 \in \mathbb{F}_0; 1g$ let us first notice that $W(0; 0) = 1$ and $W(0; 1)$ is indeed the function found in equation (13). For $m_1 \in \mathbb{F}_1; \dots; 30g$ and $m_2 \in \mathbb{F}_0; 1g$ define the functions $w_{m_2}(m_1)$ as follows

$$W(m_1; m_2) = w_{m_2}(m_1) W(m_1 - 1; m_2);$$

Equations (11), (14)-(16) and (18)-(21) imply that for such range of m_1 and m_2 we have

$$g_m(n) = w_{m_2}(m_1);$$

where

$$m_2 = 1 \quad m \pmod{2}; \quad m_1 = 29 \quad n \pmod{31};$$

Finally, under identification $w_{m_2}(0) = W(0; m_2)$ we obtain

$$W(m_1; m_2) = \sum_{i=0}^{\mathbb{F}-1} w_{m_2}(i);$$

which, together with factorization (17), gives the wave function for all values of $(n_1; n_2; n_3) \in \mathbb{Z}^3$.

Remark. For $(m_1; m_2) = (29; 1)$ we have $X_1 = X_2 = 1$. Because the infinity point 1 is the Weierstrass point of order two, there exist functions with divisor of poles equal to 21. This means that is not uniquely determined in this case. However it is natural to keep the divisor of , and therefore itself, exactly like it is given from the flow on Jacobian. Notice that because for $X_1 = X_2 = 1$ we stay in the divisor W_{g-1} then this ambiguity does not affect construction of the -function.

The coefficients $c_0^{(k)}(n_1; n_2; n_3)$, $k = 1; 2; 3$, of expansion of the wave function can be obtained from factorization (17) and are given by

$$(22) \quad c_0^{(1)}(n_1; n_2; n_3) = 6^{n_3+p} 4^q \frac{W(m_1; m_2)}{t_1^{m_1}} \Big|_{t_1=0};$$

$$(23) \quad c_0^{(2)}(n_1; n_2; n_3) = 6^{n_2} 5^{n_3+q} 4^p W(m_1; m_2) \Big|_{t_2=0};$$

$$(24) \quad c_0^{(3)}(n_1; n_2; n_3) = 5^{n_2+n_3} 3^p 6^q W(m_1; m_2) \Big|_{t_3=0};$$

Using definition of the function for nonzero t_i , i.e. equation (4), and putting $t_i = 0$ for points of the divisor W_{g-1} we obtain the corresponding \mathbb{F}_7 -valued solution of the discrete KP equation (6). This -function is presented in Figure 1. Notice that due to quasi-periodicity of the -function, we have to calculate the solution in this way only for a finite range of values of the variables.

The periods c_i , $i = 1; 2; 3$, of zeros of the -function are, respectively, 31, 62 and 62. Equivalently, the "period vectors" of zeros can be chosen as

$$v_1 = \begin{pmatrix} 0 & 1 \\ 4 & 1 \\ 1 & 1 \end{pmatrix} A \quad v_2 = \begin{pmatrix} 0 & 1 \\ 11 & 1 \\ 2 & 0 \end{pmatrix} A \quad v_3 = \begin{pmatrix} 0 & 1 \\ 6 & 1 \\ 6 & 0 \end{pmatrix} A :$$

Because $c_{(1)1} = 6$, $c_{(2)1} = 3$ and $c_{(3)1} = 5$ then periods of the functions c_i , $i = 1; 2; 3$, in that variable are, respectively, 2 31, 3 31 2 and 6 31. Moreover, we have $(31; 0; 0) = 3$ which gives $(62; 0; 0) = 5 = 3 \pmod{7}$, and therefore the period of in n_1 is 6 2 31.

ACKNOWLEDGMENTS

The paper was partially supported by the University of Warmia and Mazury in Olsztyn under the grant 522-1307-0201 and by KBN grant 2 P03B 12622.




FIGURE 1. F_7 -valued solutions of discrete KP equation out of genus two hyperelliptic curve C . Variables n_1 (directed to the right) and n_2 (directed up) take values from 0 to 34. Subsequent figures are for values of $n_3 = 1; 0; 1; 2$.

REFERENCES

1. E. D. Belokolos, A. I. Bobenko, V. Z. Enol'skii, A. R. Its, and V. B. Matveev, *Algebro-geometric approach to nonlinear integrable equations*, Springer-Verlag, Berlin, 1994.
2. M. Białeckki, *Methods of algebraic geometry over finite fields in construction of integrable cellular automata*, PhD dissertation, Warsaw University, Institute of Theoretical Physics, 2003.
3. M. Białeckki and A. Doliwa, *The discrete KP and KdV equations over finite fields*, Theor. Math. Phys. **137** (2003), 1410–1416.
4. A. Bobenko, M. Bordemann, Ch. Gunn, and U. Pinkall, *On two integrable cellular automata*, Comm. Math. Phys. **158** (1993), 127–134.
5. M. Bruschi and P. M. Santini, *Cellular automata in 1+1, 2+1 and 3+1 dimensions, constants of motion and coherent structures*, Physica D **70** (1994), 185–209.
6. G. Cornell and J. H. Silverman (eds.), *Arithmetic geometry*, Springer-Verlag, New York, 1986.
7. A. Doliwa, M. Białeckki, and P. Klimczewski, *The Hirota equation over finite fields: algebro-geometric approach and multisoliton solutions*, J. Phys. A **36** (2003), 4827–4839.

8. A.S. Fokas, E.P. Papadopoulou, and Y.G. Saridakis, *Soliton cellular automata*, Physica D **41** (1990), 297–321.
9. P. Griffiths and J. Harris, *Principles of algebraic geometry*, John Wiley and Sons, New York, 1978.
10. R. Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977.
11. R. Hirota, *Nonlinear partial difference equations. III. Discrete sine-Gordon equation*, J. Phys. Soc. Jpn. **43** (1977), 2079–2086.
12. ———, *Discrete analogue of a generalized Toda equation*, J. Phys. Soc. Jpn. **50** (1981), 3785–3791.
13. N. Koblitz, *Algebraic aspects of cryptography*, Springer-Verlag, Berlin, 1998.
14. I. M. Krichever, *Algebraic curves and non-linear difference equations*, Uspekhi Mat. Nauk **33** (1978), 215–216.
15. I. M. Krichever, P. Wiegmann, and A. Zabrodin, *Elliptic solutions to difference non-linear equations and related many body problems*, Commun. Math. Phys. **193** (1998), 373–396.
16. S. Lang, *Abelian varieties*, Interscience Publishers, Inc., New York, 1958.
17. ———, *Algebra*, Addison-Wesley, Reading, Mass., 1970.
18. J. Matsukidaira, J. Satsuma, D. Takahashi, T. Tokihiro, and M. Torii, *Toda-type cellular automaton and its N -soliton solution*, Phys. Lett. A **225** (1997), 287–295.
19. A. J. Menezes, Y. H. Wu, and R. J. Zuccherato, *An elementary introduction to hyperelliptic curves*, appendix in [13], pp. 151–178.
20. J. S. Milne, *Jacobian varieties*, Chapter VII in [6], pp. 167–212.
21. C. Moreno, *Algebraic curves over finite fields*, University Press, Cambridge, 1991.
22. D. Mumford, *An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg–de Vries equation, and related nonlinear equations*, Proceedings of the International Symposium on Algebraic Geometry (M. Nagata, ed.), Kinokuniya, Tokyo, 1978, pp. 115–153.
23. J. von Neumann, *The general and logical theory of automata*, The collected works of John von Neumann (A. W. Taub, ed.), vol. 5, Pergamon Press, 1963.
24. I. Shafarevich, *Basic algebraic geometry*, Springer-Verlag, Heidelberg, 1974.
25. H. Stichtenoth, *Algebraic function fields and codes*, Springer-Verlag, Berlin, 1993.
26. D. Takahashi and J. Satsuma, *A soliton cellular automaton*, J. Phys. Soc. Jpn. **59** (1990), 3514–3519.
27. T. Tokihiro, D. Takahashi, J. Matsukidaira, and J. Satsuma, *From soliton equations to integrable cellular automata through a limiting procedure*, Phys. Rev. Lett. **76** (1996), 3247–3250.
28. S. Ulam, *Random processes and transformations*, Proceedings of the International Congress of Mathematicians, Cambridge, MA, 30 August–6 September 1950 (P. A. Smith and O. Zariski, eds.), AMS, Providence, 1952, pp. 264–275.
29. S. Wolfram, *Theory and application of cellular automata*, World Scientific, Singapore, 1986.

M. BIAŁECKI, INSTYTUT GEOFIZYKI PAN, UL. KSIĘCIA JANUSZA 64, 01-452 WARSZAWA, POLAND

E-mail address: bialecki@igf.edu.pl

A. DOLIWA, WYDZIAŁ MATEMATYKI I INFORMATYKI, UNIwersytet WARMIŃSKO–MAZURSKI, UL. ŻOŁNIERSKA 14A, 10-561 OLSZTYN, POLAND

E-mail address: doliwa@matman.uwm.edu.pl