

Symmetry constraints for dispersionless integrable equations and systems of hydrodynamic type

L.V. Bogdanov and B.G. Konopelchenko^y

Abstract

Symmetry constraints for $(2+1)$ -dimensional dispersionless integrable equations are considered. It is demonstrated that they naturally lead to systems of hydrodynamic type which arise within the reduction method. One also easily obtains an associated complex curve (Sato function) and corresponding generating equations. Dispersionless KP and 2D TL hierarchy are considered as illustrative examples.

1 Introduction

Dispersionless integrable equations arise in various contexts and have variety of applications from complex analysis to topological field theory (see [1]–[22]). The reduction method seems to be the most developed method of construction of exact solutions of dispersionless equations [5, 6, 8, 17], [23]–[34]. It reveals a close connection between such equations and certain systems of hydrodynamic type, emphasising a role of complex curves (Sato functions). The reduction method has been successfully applied to several dispersionless equations. However, an intrinsic background of its validity and efficiency seems to remain not very clear.

In the present paper it is shown that systems of hydrodynamic type arising within the reduction method are nothing but the symmetry constrained $(2+1)$ -dimensional integrable equations. We will consider here the dispersionless KP and 2D TL hierarchies.

First, we find a general infinitesimal nonispectral¹ symmetry transformation of the dispersionless τ -function (free energy F). Then we impose symmetry constraint of the type 'ispectral symmetry = nonispectral symmetry', similar to dispersionfull case [35]–[40]. It is demonstrated that

¹L.D. Landau ITP, Kosygin str. 2, Moscow 119334, Russia

^yDipartimento di Fisica dell'Università di Lecce and Sezione INFN, 73100 Lecce, Italy

such a constraint reduces $(2+1)$ -dimensional dispersionless equation to a set of $(1+1)$ -dimensional systems of hydrodynamic type with a finite number of dependent variables. We present also a simple way to find an associated complex curve and to construct generating equations.

Our approach starts with the Θ -method applied to dispersionless equations. It provides us with a simple and natural way to consider symmetries and symmetry constraints. At the same time, an analysis of such constraints indicates a direction of a possible extension of the quasiclassical Θ -method.

2 Quasiclassical Θ -method

Dispersionless integrable hierarchies can be described in different forms within different approaches. In the papers [41, 42, 44] it was shown that such hierarchies can be introduced starting with the nonlinear Beltrami equation (quasi-classical Θ -problem)

$$S_z = W(\bar{z}; z; S_z); \quad (1)$$

where $z \in \mathbb{C}$, \bar{z} means complex conjugation, $S_z = \frac{\partial S(z; \bar{z})}{\partial z}$; $S_{\bar{z}} = \frac{\partial S(z; \bar{z})}{\partial \bar{z}}$; and W (quasi-classical Θ -data) is an analytic function of S_z ,

$$W(\bar{z}; z; S_z) = \sum_{p=0}^{\infty} w_p(\bar{z}; z) (S_z)^p; \quad (2)$$

Applying the quasi-classical Θ -dressing method based on equation (1), one can get dispersionless integrable hierarchies and the corresponding addition formulae in a very regular and simple way. Such an approach reveals also the connection of dispersionless hierarchies with the quasi-conformal mappings on the plane.

The specific choice of the hierarchy depends on the choice of domain G which is the support of the Θ -data. Once the domain is fixed, we introduce the class of functions $S_0(z)$, having the derivative $\partial_z S_0(z)$ analytic in G , and formulate the boundary problem in G in the following way. Let the function $S_0(z)$ be given. The problem is to find the function $S = S_0 + \mathcal{S}$, satisfying (1) in G , with \mathcal{S} analytic outside G and decreasing at infinity.

Introducing parameterization of the class of functions $S_0(z)$ in terms of finite number of variables (times), and using the technique of quasiclassical Θ -dressing method, it is possible to demonstrate that $S(z; \bar{z}; t)$ is a solution of Hamilton-Jacobi equations of corresponding dispersionless hierarchy.

For dispersionless KP (dKP) hierarchy we take unit disc D as G , and parameterization of $S_0(z)$ is given by

$$S_0(z;t) = \sum_{n=1}^{\infty} t_n z^n; \quad (3)$$

$S(z;t) = S_0 + \mathfrak{S}$, \mathfrak{S} is analytic outside the unit disc, and at infinity it has an expansion $\mathfrak{S} = \sum_{i=1}^{\infty} S_i(t) z^{-i}$. The quantity $p = \frac{\partial S}{\partial t_1}$ is a basic homomorphism [42]. Important role in the theory of dKP hierarchy is played by the equation

$$p(z) - p(z_1) + z_1 \exp(-D(z_1)S(z)) = 0; \quad z \in \mathbb{C}; \quad z_1 \in \mathbb{C} \cap D; \quad (4)$$

(where $D(z)$ is the quasiclassical vertex operator, $D(z) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{z^n} \frac{\partial}{\partial t_n}$, $|z| > 1$) which generates Hamilton-Jacobi equations of the hierarchy by expansion into the powers of z^{-1} at infinity (see, e.g., [47]). This equation also implies existence of the \mathfrak{F} -function, characterized by the relation

$$\mathfrak{S}(z;t) = -D(z)F(t); \quad (5)$$

and provides the dispersionless addition formula.

For the dispersionless 2D TL (d2D TL) hierarchy the \mathfrak{G} -data are localized on the domain G which is an annulus $a < |z| < b$, where a, b ($a, b \in \mathbb{R}$, $a, b > 0$; $b > a$) are arbitrary, S_0 can be represented as [8, 12, 47]

$$S_0(t;x;y) = t \log z + \sum_{n=1}^{\infty} z^n x_n + \sum_{n=1}^{\infty} z^{-n} y_n; \quad (6)$$

We assume that $\mathfrak{S}(z) = \sum_{n=1}^{\infty} \frac{S_n}{z^n}$ as $z \rightarrow \infty$ and denote $\mathfrak{S}(0) = \mathfrak{S}$, $G_+ = \{z; |z| > b\}$, $G_- = \{z; |z| < a\}$. The functions $p_+ = \frac{\partial S}{\partial x_1}$ and $p_- = \frac{\partial S}{\partial y_1}$ have pole singularities while $p = \frac{\partial S}{\partial t}$ has a logarithmic singularity. Relations, characterizing the \mathfrak{F} -function F , are $\mathfrak{F} = D F$, $\mathfrak{S}(z_1) = \sum_{n=1}^{\infty} D_+(z_1) F(z_1 \in G_+)$, $\mathfrak{S}(z_2) = \sum_{n=1}^{\infty} D_-(z_2) F(z_2 \in G_-)$, where $D_+(z) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{z^n} \frac{\partial}{\partial x_n}$, $D_-(z) = \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{z^n} \frac{\partial}{\partial y_n}$, $D = \frac{\partial}{\partial t}$. Hamilton-Jacobi equations of the hierarchy are generated by the relations [8, 12, 47]

$$e^{p(z)} - e^{p(z_1)} + z_1 e^{D_+(z_1)S(z)} = 0; \quad z \in \mathbb{C}; \quad z_1 \in G_+; \quad (7)$$

$$e^{p(z)} - 1 - e^{D_-(z_2)S(z)} = z_2 e^{(D - D_-(z_2))} ; \quad z \in \mathbb{C}; \quad z_2 \in G_-; \quad (8)$$

these relations also provide dispersionless addition formulae for F .

3 τ -function and symmetries

It was noted in [44] that equation (1) is a Lagrangian one. It can be obtained by variation of the action (for the boundary problem in G)

$$f = \frac{1}{2} \int_{\partial G} \bar{z} z \frac{1}{2} \mathbb{S}_z \mathbb{S}_{\bar{z}} W_{-1}(z; z; S_z) dz \wedge d\bar{z}; \quad (9)$$

where

$$W_{-1}(z; z; S_z) = \sum_{p=0}^{\infty} w_p(z; z) \frac{(S_z)^{p+1}}{p+1}; \quad @ W_{-1}(z; z; \cdot) = W(z; z; \cdot):$$

One should consider independent variations of \mathbb{S} , possessing required analytic properties (analytic outside G , decreasing at infinity), keeping S_0 fixed.

In [48] it was proved for dispersionless KP and 2D TL hierarchies that the action (9) evaluated on the solution of nonlinear Beltrami equation (1) gives a τ -function of the corresponding hierarchy. Here we formulate first an auxiliary statement, valid for an hierarchy with arbitrary G , and then recall the results of [48].

Lemma 1 Let us evaluate the action (9) on the solution of the boundary problem for (1) and consider it as a functional of S_0 ,

$$F(S_0) = \frac{1}{2} \int_{\partial G} \bar{z} z \frac{1}{2} \mathbb{S}_z \mathbb{S}_{\bar{z}} W_{-1}(z; z; S_z) dz \wedge d\bar{z}; \quad (10)$$

Then

$$F = \frac{1}{2} \int_{\partial G} \bar{z} \mathbb{S} d S_0; \quad (11)$$

Using the formula (11), it is easy to reproduce the statements of the work [48].

Proposition 1 The function

$$F(t) = \frac{1}{2} \int_{\partial D} \bar{z} z \frac{1}{2} \mathbb{S}_z(t) \mathbb{S}_{\bar{z}}(t) W_{-1}(z; z; S_z(t)) dz \wedge d\bar{z}; \quad (12)$$

i.e., the action (9) evaluated on the solution of the problem (1), with S_0 given by (3), is a τ -function of dKP hierarchy.

Proposition 2 The function

$$F(t; x; y) = \frac{1}{2} \oint_G \frac{1}{z} \mathbb{S}_z(t; x; y) \mathbb{S}_z(t; x; y) W_1(z; z; S_z(t; x; y)) dz \wedge dz;$$

with S_0 given by (6), evaluated on the solution of the problem (1), is a $\bar{\partial}$ -function of dispersionless 2D TL hierarchy (G is an annulus defined above).

The function W is the $\bar{\partial}$ -data for the quasiclassical $\bar{\partial}$ -problem. Its variations provide us with a wide class of variations of the function F (first we will use formula (10) defined for arbitrary G).

For the functions W of the form (2), varying $w_n(z; z)$, one has

$$W = \sum_{n=1}^{\infty} w_n(z; z) (S_z)^n; \quad W_1 = \sum_{n=1}^{\infty} \frac{w_n}{n+1} (S_z)^{n+1};$$

and

$$F = \frac{1}{2} \oint_G U(z; z; S_z) dz \wedge dz; \quad (13)$$

where $U(z; z; \cdot)$ is an arbitrary function with the support in z -plane belonging to G , analytic in \cdot . The formula (13) gives a general variation (in infinitesimal symmetry transformation) of the $\bar{\partial}$ -function. Let us consider some special symmetry transformations.

Considering elementary variation $w_{n_0} = \delta_{n_0}(z - z_0)$, $w_n = 0; n \neq n_0$, one gets

$$F = \frac{1}{2} \oint \frac{1}{z} \frac{1}{(n_0 + 1)} (S_z)^{n_0+1} \Big|_{z=z_0}; \quad (14)$$

and, respectively,

$$\mathbb{S} = \frac{1}{2} \oint \frac{1}{z} \frac{1}{(n_0 + 1)} D(z) (S_z(z_0))^{n_0+1}; \quad (15)$$

Taking superposition of elementary variations (14), we obtain a variation of the form

$$F = f(\mathbb{S}(z_0)); \quad (16)$$

where f is an arbitrary analytic function (summation over different points and integration over z_0 are also possible).

Another simple symmetry transformation is given by

$$F = c(S(z_2) - S(z_1)); \quad (17)$$

where z_1, z_2 are arbitrary points belonging to the domain G , and c is some constant. Indeed,

$$\begin{aligned} S(z_2) - S(z_1) &= \int_{z_1}^{z_2} d(S) = \int_{z_1}^{z_2} S_z dz + \int_{z_1}^{z_2} S_z dz \\ &= \int_{z_1}^{z_2} S_z dz + \int_{z_1}^{z_2} W(z; z; S_z) dz; \end{aligned}$$

thus the function $S(z_2) - S(z_1)$ corresponds to the special case of general variation (13). In the limit $z_2 \rightarrow z_1$ the transformation (17) is reduced to elementary symmetries (14).

4 Symmetry constraints

Given infinitesimal symmetries of the form

$$F = \dots;$$

(cf., (13), (16), (17)), it is possible to introduce symmetry constraints

$$\frac{\partial}{\partial t_1} F = 0; \quad (18)$$

These constraints are preserved by the flows of the hierarchy and represent a reduction of the hierarchy. There is a lot of similarity between constraints (18) and symmetry constraints in the $(2+1)$ -dimensional dispersionless case, which are rather well studied [35]–[40]. Below we will demonstrate that constraints (18) lead to hydrodynamic reductions, and give a regular way to introduce basic objects connected with these reductions.

Technically, in the dispersionless case one starts with linear equations of the hierarchy (which give the hierarchy itself in the form of compatibility conditions). Symmetry constraints connect the potentials in these equations with some special wave functions, and in the end one obtains $(1+1)$ -dimensional integrable system for the set of wave functions. The simplest symmetry constraint for KP hierarchy leads to nonlinear Schrödinger equation.

In the dispersionless case we will follow the same strategy. The role of linear equations of the hierarchy is played by Hamilton–Jacobi equations of the dispersionless hierarchy.

5 Zakharov reduction for the dKP hierarchy

We begin with a simple example of symmetry constraint for the dKP hierarchy. The first Hamilton-Jacobi equation of dKP hierarchy (dispersionless analogue of Lax equation) is

$$S_y = p^2 + 2u; \quad p = S_x; \quad u = \partial_x S_1; \quad (19)$$

where $x = t_1, y = t_2$. Let us consider the symmetry constraint

$$F_x = f(S_z(z_0)); \quad (20)$$

which is equivalent to the condition (cf., (5))

$$u = \partial_x f(S_z(z_0)):$$

Differentiating (19) with respect to z and evaluating the result at $z = z_0$, we get

$$\frac{\partial_y S_z(z_0)}{\partial_x S_z(z_0)} = 2p_0; \quad p_0 = p(z_0):$$

Using this equation and evaluating (19) at $z = z_0$, we obtain a system of hydrodynamic type

$$\begin{aligned} \partial_y p_0 &= \partial_x (p_0^2 + 2u); \\ \partial_y u &= 2\partial_x (p_0 u); \end{aligned} \quad (21)$$

This system is well known [2], it represents a dispersionless limit of nonlinear Schrödinger equation. Thus a constraint (20) represents a proper dispersionless analogue of the simplest symmetry constraint of KP hierarchy.

It is possible to move further and demonstrate that the symmetry constraint (20) leads to explicit expression for the function $z(p)$, which plays a fundamental role in the picture of constrained hierarchy and allows to find Riemann invariants and hodograph equations. To do this, we will use generating Hamilton-Jacobi equation (4) rewritten as

$$D(z)S(z^0) = \log \frac{p(z) - p(z^0)}{z}; \quad |j| > 1; \quad (22)$$

which produces infinite set of Hamilton-Jacobi equations by expansion into the powers of z^{-1} at infinity (see, e.g., [47]). Differentiating this equation by z^0 and evaluating the result at $z^0 = z_0$, we get

$$\frac{D(z)S_z(z_0)}{\partial_x S_z(z_0)} = \frac{1}{p - p_0}:$$

Then, using the constraint (20) and simple formula

$$D(z)F_x = z - p;$$

we obtain the relation

$$z = p + \frac{u}{p - p_0}; \quad (23)$$

The function $z(p)$ (Sato function) plays a crucial role in the picture of dispersionless hierarchy. Dispersionless KP hierarchy (for general $z(p)$) can be written in the form $\partial_n z(p) = f z_+^n; z_0$: Relation (23) defines the function $z(p)$ for the constrained hierarchy, it is well known [2] and is usually called Zakharov reduction. Thus we have demonstrated that symmetry constraint (20) leads to Zakharov reduction (23).

Riemann invariants for the constrained hierarchy of equations of hydrodynamic type (the first of which is (21)), are given by the values of $z(p)$ (23) at p_i where $\frac{\partial z}{\partial p}$ vanishes, $z_i = z(p_i)$, $\frac{\partial z}{\partial p}(p_i) = 0$. It is rather straightforward to introduce hodograph transform for the solution of constrained hierarchy [15].

6 Generic symmetry constraint for the dKP hierarchy

Now we will use the symmetry (17) and consider the constraint

$$F_x = \sum_{i=1}^N c_i (S_i - \bar{S}_i); \quad (24)$$

where $S_i = S(z_i)$, $\bar{S}_i = S(\bar{z}_i)$, z_i, \bar{z}_i are some sets of points, and c_i are arbitrary constants. In equivalent form,

$$u = \partial_x \sum_{i=1}^N c_i (S_i - \bar{S}_i) = \sum_{i=1}^N c_i (p_i - \bar{p}_i):$$

Evaluating Hamilton-Jacobi equation (19) and its higher counterpart

$$S_t = p^3 + 3up + \frac{3}{2}S_1;$$

where $t = t_3$, at z equal to z_i, \bar{z}_i , we immediately get two systems of hydrodynamic type

$$\begin{aligned} \sum_{k=1}^8 \partial_y p_k &= \partial_x (p_k^2) + 2 \sum_{i=1}^N c_i (p_i - \bar{p}_i) \\ \vdots \\ \sum_{k=1}^8 \partial_y p_k &= \partial_x (p_k^2) + 2 \sum_{i=1}^{X^i} c_i (p_i - \bar{p}_i) \quad k = 1, \dots, N \end{aligned} \quad (25)$$

and

$$\begin{aligned} \sum_{k=1}^N \partial_t p_k &= \partial_x (p_k^3) + 3p_k \sum_{i=1}^N c_i(p_i - p_i) + \frac{3}{2} \sum_{i=1}^N c_i(p_i^2 - p_i^2) \\ \sum_{k=1}^N \partial_t p_k &= \partial_x (p_k^3) + 3p_k \sum_{i=1}^N c_i(p_i - p_i) + \frac{3}{2} \sum_{i=1}^N c_i(p_i^2 - p_i^2) \end{aligned} \quad (26)$$

These systems can be written in Hamiltonian form,

$$\begin{aligned} \partial_t p &= J \frac{H_{n+1}}{p}; \quad J = \begin{pmatrix} C^{-1} & 0 \\ 0 & C^{-1} \end{pmatrix} \partial_x; \\ H_3 &= \frac{1}{3} \sum_{i=1}^N dx \sum_{i=1}^N c_i(p_i^3 - p_i^3) + 3 \sum_{i=1}^N c_i(p_i - p_i)^2; \\ H_4 &= \frac{1}{4} \sum_{i=1}^N dx \sum_{i=1}^N c_i(p_i^4 - p_i^4) + 6 \sum_{i=1}^N c_i(p_i - p_i) \sum_{i=1}^N c_i(p_i^2 - p_i^2); \end{aligned}$$

where $p = (p_1, \dots, p_N, p_1, \dots, p_N)^t$, C is diagonal $N \times N$ matrix with the entries c_i , $C_{ik} = c_i \delta_{ik}$, $i, k = 1, \dots, N$.

Common solution of the systems (25), (26) generates a solution $u = \sum_{i=1}^N c_i(p_i - p_i)$ of the dKP equation.

The Sato function $z(p)$, which allows to construct Riemann invariants and higher Hamiltonians, is the central object for the constrained hierarchy. To find it, we, similar to the first example, use the generating Hamilton-Jacobi equation (22). Evaluating this equation at z^0 equal to z_i , z_i and combining the results, we get

$$D(z) \sum_{i=1}^N c_i(S_i - S_i) = \sum_{i=1}^N c_i \log \frac{p - p_i}{p - p_i}; \quad (27)$$

Then, due to the constraint (24),

$$D(z)F_x = \sum_{i=1}^N c_i \log \frac{p - p_i}{p - p_i};$$

and finally we obtain the function $z(p)$,

$$z = p \sum_{i=1}^N c_i \log \frac{p - p_i}{p - p_i}; \quad (28)$$

Equation $\frac{\partial z}{\partial p} = 0$ is an algebraic one, having $2N$ roots p , and Riemann invariants for the constrained hierarchy are $\lambda = z(p)$.

Expansion of $z(p)$ at $p \rightarrow 1$ looks like

$$z = p + \sum_{n=1}^N \frac{v_n}{p^n}; \quad v_n = \sum_{i=1}^N \frac{c_i (p_i^n - \bar{p}_i^n)}{n}; \quad (29)$$

Equation (22) allows us to obtain the generating equation for the whole hierarchy of systems of hydrodynamic type associated with the constraint (24). Indeed, differentiating (22) with respect to x and evaluating it at the points z_i, \bar{z}_i , one gets

$$\begin{aligned} D(z)p_k &= \partial_x \log(p - p_k) \\ D(z)\bar{p}_k &= \partial_x \log(p - \bar{p}_k) \end{aligned} \quad k = 1; \dots; N; \quad (30)$$

where p is a function of $z, (p_1; \dots; p_N; \bar{p}_1; \dots; \bar{p}_N)$, defined by the relation (28). Expanding both sides of this system into the powers of z^{-1} , one gets the systems (25), (26) and their higher counterparts. Introducing to $S_0(z; t)$ dependence on extra time t_{\log} , $S_0(z^0) \rightarrow S_0(z^0) - t_{\log} \log(1 - \frac{z^0}{z})$, $\frac{\partial}{\partial t_{\log}} = D(z)$, we may consider (30) as a system of hydrodynamic type with the time t_{\log} . Hamiltonian for this system is given by the expression

$$H_{\log} = \int dx \left[\frac{1}{2} p^2 \sum_{i=1}^N c_i p_i - p_i + p_i \log(p - p_i) - \bar{p}_i \log(p - \bar{p}_i) \right];$$

Differentiating (30) by z , we will get another form of generating system,

$$\begin{aligned} \partial_z D(z)p_k &= \partial_x \frac{p_z}{p - p_k} \\ \partial_z D(z)\bar{p}_k &= \partial_x \frac{p_z}{p - \bar{p}_k} \end{aligned} \quad k = 1; \dots; N; \quad (31)$$

where p, p_z are functions of $z, (p_1; \dots; p_N; \bar{p}_1; \dots; \bar{p}_N)$, defined by the relation (28). This system can be considered as a system of hydrodynamic type with the time t_{pole} , $S_0(z^0) \rightarrow S_0(z^0) + t_{\text{pole}} (z - z^0)^{-1}$, $\frac{\partial}{\partial t_{\text{pole}}} = \partial_z D(z)$. Hamiltonian for the generating system (31) is remarkably simple,

$$H_{\text{pole}} = \int p dx;$$

and its expansion into the powers of z^{-1} gives a general formula for the Hamiltonians of constrained hierarchy,

$$H_n = \frac{1}{n} \int \text{res}_1 z(p)^n dx = \partial_x \mathcal{E}_n; \quad (32)$$

which is in agreement with general results [1].

In a similar way one can treat the symmetry constraints

$$\frac{\partial F}{\partial t_n} = \sum_{i=1}^N c_i (S_i - \tilde{S}_i): \quad (33)$$

In particular, since

$$\frac{\partial S}{\partial t_n} = p^n + u_{n-2} p^{n-2} + \dots + u_0$$

one obtains a corresponding Sato function

$$E(p) = z^n = p^n + u_{n-2} p^{n-2} + \dots + u_0 + \sum_{i=1}^N c_i \log \frac{p - p_i}{p - \tilde{p}_i}: \quad (34)$$

Remark. It is easy to check that in the limit $z_i = z_i + \epsilon_i \epsilon_i \rightarrow 0$, $c_i \epsilon_i = \text{const}$, the Sato function (34) reproduces the curve

$$z^n = p^n + u_{n-2} p^{n-2} + \dots + u_0 + \sum_{i=1}^N \frac{a_i}{p - p_i}; \quad (35)$$

which has been discussed in [15, 17]. At $n = 1$ one gets a general Zakharov reduction.

7 Symmetry constraint for the d2D TL hierarchy

Now we will consider symmetry reduction of d2D TL hierarchy defined by the constraint

$$F_t = \sum_{i=1}^N c_i (S_i - \tilde{S}_i): \quad (36)$$

Technically this case is very similar to that considered in the previous section, and we will omit some details. The first Hamilton-Jacobi equations of d2D TL hierarchy are

$$\partial_x S(z) = e^p - \partial_t \tilde{S}_1; \quad (37)$$

$$\partial_y S = e^{-t-p}; \quad (38)$$

where $x = x_1$, $y = y_1$. Using the constraint (36), from these equations we get two systems of hydrodynamic type, namely

$$\begin{cases} \partial_x p_i = \partial_t e^{p_i} + \sum_k c_k (e^{p_k} - e^{\tilde{p}_k}) \\ \partial_x \tilde{p}_i = \partial_t e^{\tilde{p}_i} + \sum_k c_k (e^{p_k} - e^{\tilde{p}_k}) \end{cases} \quad i = 1, \dots, N \quad (39)$$

and

$$\begin{cases} \partial_y p_i = \partial_t \exp \left(p_i + \sum_{k=1}^N c_k (p_k - p_k) \right) \\ \partial_y p_i = \partial_t \exp \left(p_i + \sum_{k=1}^N c_k (p_k - p_k) \right) \end{cases} \quad i = 1, \dots, N \quad (40)$$

Both systems can be written in Hamiltonian form,

$$\partial_x p = J \frac{H_1^+}{p}; \quad \partial_y p = J \frac{H_1}{p}; \quad J = \begin{pmatrix} E & C^{-1} \\ 0 & C^{-1} \end{pmatrix} \partial_t;$$

where E is $2N \times 2N$ matrix with all entries equal to 1, $E_{ik} = 1$, $i, k = 1, \dots, 2N$, C is diagonal $N \times N$ matrix with the entries c_i , $C_{ik} = c_i \delta_{ik}$, $i, k = 1, \dots, N$, and the Hamiltonians are, respectively,

$$H_1^+ = \sum_{k=1}^N \frac{d}{dt} c_k (e^{p_k} - e^{p_k}); \quad (41)$$

$$H_1 = \sum_{k=1}^N \frac{d}{dt} \exp \left(\sum_{k=1}^N c_k (p_k - p_k) \right) \sum_{k=1}^N c_k (e^{p_k} - e^{p_k}); \quad (42)$$

Using generating equation (7), we obtain the function $z(p)$ for the constrained hierarchy,

$$z = e^p \prod_{k=1}^N \frac{e^p - e^{p_k}}{e^p - e^{p_k}}^{c_k}; \quad (43)$$

Starting with second generating equation (8), we get equivalent form of this relation,

$$z = \exp p \prod_{k=1}^N c_k (p_k - p_k) \prod_{k=1}^N \frac{e^p - e^{p_k}}{e^p - e^{p_k}}^{c_k}; \quad (44)$$

Equation $\frac{\partial z}{\partial p} = 0$ is algebraic (with respect to the variable $p = e^p$), having $2N$ roots, and Riemann invariants for the constrained hierarchy are $z(p)$.

Generating systems for the constrained hierarchy read

$$\begin{cases} D_+(z)p_k = \partial_t \log(e^p - e^{p_k}) \\ D_+(z)p_k = \partial_t \log(e^p - e^{p_k}) \end{cases} \quad k = 1, \dots, N; \quad (45)$$

$$\begin{cases} D_-(z)p_k = \partial_t \log(1 - e^{p - p_k}) \\ D_-(z)p_k = \partial_t \log(1 - e^{p - p_k}) \end{cases} \quad k = 1, \dots, N; \quad (46)$$

(logarithmic form) and

$$\begin{aligned} & \leq D_z(z) p_k = \partial_t \frac{p_z}{e^{p_k} p - 1} \\ & : D_z(z) p_k = \partial_t \frac{p_z}{e^{p_k} p - 1} \quad k = 1, \dots, N; \end{aligned} \quad (47)$$

(pole form). Hamiltonian for the pole form of generating equations is

$$H(z) = \frac{1}{z} \int^Z p dt;$$

Expansion of this Hamiltonian at $z = 1$ and $z = 0$ gives general expressions for the Hamiltonians of constrained hierarchy,

$$H_n^+ = \frac{1}{n} \int^Z \text{res}_1 z(t)^{n-1} dt; \quad (48)$$

$$H_n^- = \frac{1}{n} \int^Z \text{res}_0 z(t)^{n-1} dt; \quad (49)$$

where $\partial_t = e^p$.

In more detail, the Hamiltonian and bi-Hamiltonian structure of equation considered above will be considered elsewhere.

Symmetry constraints of the type $F_{x_n} = \sum_{i=1}^N C_i (S_i - \bar{S}_i)$ and $F_{y_n} = \sum_{i=1}^N C_i (S_i - \bar{S}_i)$ are treated analogously. The corresponding Sato function is given by (34) with substitutions $p \rightarrow e^p, p \rightarrow e^{-p}$, respectively.

8 Conclusion

As we have seen, the systems of hydrodynamic type and associated complex curves (Sato functions) are direct consequences of the generating Hamilton-Jacobi equations (4), (7,8) under the symmetry constraint.

We would like to note here that complex curves discussed above are connected with the ∂ -equation in a very natural way. Let us consider the dKP hierarchy as an example. The function $p = \partial_x S$ solves a linear Beltrami equation $p_z = W \bar{p}_z$, and under certain conditions is a basic homeomorphism. The inverse function $z(p; p)$ is analytic in some neighborhood of infinity. The Cauchy-Green formula implies that

$$z = p + \frac{1}{2\pi i} \int^Z \frac{dp^0 \wedge dp^0 \partial z}{p^0 p \partial p^0}; \quad (50)$$

In the neighborhood of infinity one has

$$z = p + \sum_{n=1}^{\infty} \frac{v_n}{p^n}; \quad (51)$$

where

$$v_n = \frac{1}{2} \sum_{i=1}^N \int_{\gamma_i} p^{n-1} \frac{\partial z}{\partial p^0} dp^0 \wedge dp^1; \quad (52)$$

Let us choose a special $\frac{\partial z}{\partial p}$ with a support along the curve $\gamma = \bigcup_{i=1}^N \gamma_i$ composed of N disconnected pieces γ_i with the ends at the points z_i and \bar{z}_i and such that $\frac{\partial z}{\partial p} = \sum_{i=1}^N c_i \gamma_i$. For such 'constrained' ∂ -data formula (50) gives

$$z = p \sum_{i=1}^N c_i \log \frac{p - p_i}{p - \bar{p}_i};$$

while from (52) we obtain

$$v_n = \sum_{i=1}^N \frac{c_i (p_i^n - \bar{p}_i^n)}{n};$$

that coincides with (28,29). For the ∂ -data given by $\frac{\partial z}{\partial p} = \sum_{i=1}^N a_i (p - p_i)$, equation (50) immediately leads to the Sato function associated with Zakharov reduction (see (35)). Thus, the Cauchy-Green formula is a generator of complex curves associated with equations of hydrodynamic type.

Equation (50) is the two-dimensional extension of the integral equation discussed in [2], [3].

Acknowledgments

Authors would like to thank Luis Martinez Alonso for useful discussions. LVB is also grateful to E.V. Ferapontov for attracting his interest to the problem of hydrodynamic reductions. LVB was supported in part by RFBR grant 01-01-00929 and President of Russia grant 1716-2003; BGK was supported in part by the grant COFIN 2002 'Sintesi'.

References

- [1] B.A. Kuperschmidt and Yu. I. Manin, Funk. Anal. Appl. 11 (3), 31{42 (1977); 17 (1), 25{37 (1978).
- [2] V.E. Zakharov, Func. Anal. Pribl. 14, 89{98 (1980); Physica 3D, 193{202 (1981).
- [3] J. Gibbons, Phys. D 3 (1981), no. 3, 503{511.
- [4] P.D. Lax and C.D. Levermore, Commun. Pure Appl. Math., 36, 253{290, 571{593, 809{830 (1983).
- [5] I.M. Krichever, Func. Anal. Pribl., 22, 37{52 (1988).
- [6] Y. Kodama, Phys. Lett. 129A, 223{226 (1988); 147A, 477{482 (1990).
- [7] B.A. Dubrovin and S.P. Novikov, Russian Math. Surveys, 44, 35{124 (1989).

- [8] K. Takasaki and T. Takebe, *Int. J. Mod. Phys. A*, Suppl. 1B, 889{922 (1992).
- [9] I. M. Krichever, *Commun. Pure Appl. Math.*, 47, 437{475 (1994).
- [10] N. M. Ercolani et al, eds., *Singular limits of dispersive waves*, Nato Adv. Sci. Inst. Ser. B Phys. 320, Plenum, New York (1994).
- [11] V. E. Zakharov, *Dispersionless limit of integrable systems in (2+1)-dimensions*, in [10], pp. 165{174 (1994)
- [12] K. Takasaki and T. Takebe, *Rev. Math. Phys.*, 7, 743{808 (1995).
- [13] R. Carroll and Y. Kodama, *J. Phys. A Math. Gen.* 28, 6373{6387 (1995).
- [14] S. Jin, C. D. Levermore and D. W. McLaughlin, *Comm. Pure and Appl. Math.*, 52, 613{654 (1999).
- [15] I. M. Krichever, *Comm. Math. Phys.*, 143, 415{429 (1992).
- [16] B. A. Dubrovin, *Comm. Math. Phys.* 145, 195{207 (1992); B. A. Dubrovin, *Nucl. Phys. B* 379, 627{289 (1992); B. A. Dubrovin and Y. Zhang, *Comm. Math. Phys.* 198, 311{361 (1998).
- [17] S. Aoyama and Y. Kodama, *Commun. Math. Phys.* 182, 185{219 (1996).
- [18] Y. Kodama, *SIAM J. Appl. Math.* 59, 2162{2192 (1999)
- [19] Y. Gibbons and S. P. Tsarev, *Phys. Lett.* 258A, 263{271 (1999).
- [20] M. M. Inceev-W einstein P. B. W iegmann and A. Zabrodin, *Phys. Rev. Lett.* 84, 5106{5109 (2000).
- [21] P. B. W iegmann and A. Zabrodin, *Commun. Math. Phys.* 213, 523{538 (2000).
- [22] M. Dunajski, L. J. Mason and P. Tod, *J. Geom. Phys.* 37, 63{93 (2001).
- [23] E. V. Ferapontov, D. A. Kirotkin and V. A. Shramchenko, *Class. Quantum Grav.* 19, L205{L210 (2002).
- [24] E. V. Ferapontov and M. V. Pavlov, *Classical Quantum Gravity* 20 (2003), no. 11, 2429{2441
- [25] M. V. Pavlov, *J. Math. Phys.* 44 (2003), no. 9, 4134{4156
- [26] E. V. Ferapontov and K. R. Khusnutdinova, *On integrability of (2+1)-dimensional quasilinear systems*, arXiv:nlin.SI/0305044
- [27] E. V. Ferapontov and K. R. Khusnutdinova, *The characterization of two-component (2+1)-dimensional integrable systems of hydrodynamic type*, arXiv:nlin.SI/0310021
- [28] M. Maras, L. Mart nez Alonso and E. Medina, *J. Phys. A* 35 (2), 401{417 (2002).
- [29] L. Mart nez Alonso and M. Maras, *J. Math. Phys.* 44 (8), (2003).
- [30] L. Mart nez Alonso and M. Maras, *A hodograph transformation which applies to the heavenly equation*, arXiv:nlin.SI/0209050
- [31] F. Guil, M. Maras and L. Mart nez Alonso, *J. Phys. A* 36 (2003), no. 14, 4047{4062
- [32] J.-H. Chang and M.-H. Tu, *J. Math. Phys.* 41 (2000), no. 12, 8117{8131
- [33] J.-H. Chang and M.-H. Tu, *J. Phys. A* 34 (2001), no. 2, 251{272
- [34] Y.-T. Chen and M.-H. Tu, *J. Phys. A* 36 (2003), 9875{9883
- [35] B. Konopelchenko and W. Strampp, *Inverse Problems* 7 (1991), no. 2, L17{L24

- [36] B. Konopelchenko, J. Sidorenko and W. Strampp, *Phys. Lett. A* 157 (1991), no. 1, 17{21
- [37] B. Konopelchenko and W. Strampp, *J. Math. Phys.* 33 (1992), no. 11, 3676{3686
- [38] Y. Cheng and Y. S. Li, *Phys. Lett. A* 157 (1991), no. 1, 22{26
- [39] Z. Yunbo, *J. Phys. A* 24 (1991), no. 18, L1065{L1072
- [40] A. Yu. Orlov, *Symmetries for Unifying Different Soliton Systems into a Single Integrable Hierarchy*, preprint IINS/Oe04/03, March 1991; the report at the NEEDS workshop, Gallipoli 1991.
- [41] B. G. Konopelchenko, L. Mart nez Alonso and O. Ragnisco, *J. Phys. A : Math. Gen.* 34, 10209{10217 (2001).
- [42] B. Konopelchenko and L. Mart nez Alonso, *Phys. Lett. A* 286, 161{166 (2001).
- [43] B. Konopelchenko and L. Mart nez Alonso, *J. Math. Phys.* 43 (7), 3807{3823 (2002)
- [44] B. Konopelchenko and L. Mart nez Alonso, *Stud. Appl. Math.* 109 (2002), no. 4, 313{336
- [45] I. N. Vekua, *Generalized analytic functions*, Pergamon Press, Oxford (1962).
- [46] I. K. Kostov et al, ψ -function for analytic curves, in: *Random matrices and their applications*, MSRI Publications, v. 40, 1{15 (2001).
- [47] L. V. Bogdanov, B. G. Konopelchenko and L. Mart nez Alonso, *Teor. Mat. Fiz.* 134 (1), 46{54 (2003).
- [48] L. V. Bogdanov and B. G. Konopelchenko, *Nonlinear Beltrami equation and ψ -function for dispersionless hierarchies*, arXiv:nlin.SI/0310040