M ultifractal Behavior of the Korean Stock-market Index KOSPI

Jae W oo Lee^{a;b;} K young Eun Lee^a Per Ame R ikvold^{b;c}

^aD epartm ent of Physics, Inha University, Incheon 402-751, Korea ^bSchool of Computational Science, Florida State University, Tallahassee FL 32306-4120, USA

^cD epartm ent of Physics, Center for Materials Research and Technology, and National High Magnetic Field Laboratory, Florida State University, Tallahassee, FL 32306-4350, USA

A bstract

We investigate multifractality in the K orean stock-market index KOSPI. The generalized qth order height-height correlation function shows multiscaling properties. There are two scaling regimes with a crossover time around $t_c=40\,\mathrm{m}$ in. We consider the original data sets and the modiled data sets obtained by removing the daily jumps, which occur due to the dilerence between the closing index and the opening index. To clarify the origin of the multifractality, we also smooth the data through convolution with a Gaussian function. After convolution we observe that the multifractality disappears in the short-time scaling regimet < t_c , but remains in the long-time scaling regimet > t_c , regardless of whether or not the daily jumps are removed. We suggest that multifractality in the short-time scaling regime is caused by the local uctuations of the stock index. But the multifractality in the long-time scaling regime appears to be due to the intrinsic trading properties, such as herding behavior, information outside the market, the long memory of the volatility, and the nonlinear dynamics of the stock market.

Keywords: Econophysics, multi-scaling, multifractal, stock market PACS: 89.65 Gh, 89.75 Da, 05.40 Fb, 05.45 Tp,

Corresponding author. Tel.: 82+ 32+ 8607660; fax: 82+ 32+ 8727562 Em ailaddress: jaewlee@inha.ac.kr (Jae W oo Lee).

1 Introduction

In recent years, concepts and techniques from statistical physics have been widely applied to economics [1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19], and the complex behaviors of economic systems have been found to be very similar to those of complex systems customarily studied in statistical physics. Stock-market indexes around the world have been accurately recorded for many years and therefore represent a rich source of data for quantitative analysis, and the statistical behaviors of stock markets have been studied by various methods, such as distribution functions [10,11,12,19], correlation functions [12,13,14], multifractal analysis [19,20,21], and network analysis of the market structure [15,16].

Multiscaling properties have been reported form any economic time series [22,23,24,25,26,27,28,29,30,31 Multifractality has been observed in stock markets [20,29,32,33,34], the price of crude oil [25], the price of commodities [34], and foreign exchange rates [24,35]. In daily stock indexes and foreign exchange rates, the generalized Hurst exponent H_q decreases monotonically with q [24,29,32,33,34,35] (see formal denitions of H_q and q in Eqs. (1) and (2) below). In the U.S.NASDAQ index, two scaling regimes have been reported: one quasi-Brownian and the other multifractal [33]. Two scaling domains have also been reported in the uctuations of the price of crude oil: H_q increases with q in the short-time domain, but decreases with q in the long-time domain [25]. The existence of multiple scaling regimes seems to depend on the resolution of the data sets in the economic time series.

Although many economic time series display multifractality (in the theory of surface scaling analysis referred to as multia nity), the origins of multifractality in the stock market are not well understood. It has been suggested that herding behavior and nonlinear complex dynamics of the stock market induce multiscaling [10]. However, it is very dicult to quantify herding behavior and complex dynamics in the stock market. Buend a et al. observed multianity in a frustrated spring-network model simulating the surface structure of cross-linked polymer gels [36]. Removing vertical discontinuities from the rough surface by convolution with a Gaussian, they observed that the multiane surface changed into a self-ane one [37]. They concluded that vertical discontinuities can be one cause of multianity. Mitchell conversely introduced artical vertical discontinuities into a self-ane surface and observed that the surface became multiane [38].

In the present paper we investigate multifractality in the Korean stock-market index KOSPI (Korean Composite Stock Price Index). We observe that local uctuations, including jumps, are responsible for multifractality in the short-time scaling regime. However, multifractality in the long-time scaling regime

is not rem oved by smoothing of the time series.

2 M ethod and R esults

We consider a set of data recorded every m inute of trading from M arch 30, 1992, through N ovember 30, 1999. We count the time during trading hours and remove closing hours, weekends, and holidays from the data. Denoting the stock-market index as x (t), the generalized qth order height-height correlation function (GHCF) F_q (t) is defined by

$$F_{q}(t) = h_{x}(t^{0} + t) \times (t^{0}) f^{1} i^{1-q};$$
 (1)

where the angular brackets denote a time average over the time series. The GHCF F_q (t) characterizes the correlation properties of the time series x (t), and for a multia ne series a power-law behavior like

$$F_{\alpha}(t) \quad t^{H_{\alpha}}$$
 (2)

is expected, where H $_{\rm q}$ is the generalized qth order H urst exponent [39]. If H $_{\rm q}$ is independent of q, the time series is monofractal. If H $_{\rm q}$ depends on q, the time series is multifractal. Multifractality is a distinctive property of the stock market index observed.

In Fig. 1(a) we present the GHCF as a function of the time interval t. We observe clear multifractal behavior in the time series of the stock index. There are two dierent scaling regimes separated by a crossover time of about $t_c=40\,\mathrm{m}$ in. The slopes of the log-log plot depend on q in each scaling regime. As shown in Fig. 1(b), H $_q=1=q$ for large q in both scaling regimes, while it saturates for small q. The 1=q behavior is consistent with numerical observations in Refs. [37,38] and analytical results for functions that include discontinuities in Ref. [38]. These behaviors are dierent from the multifractality in the price of crude oil [25]. A lvarez-R am irezet al. reported the existence of two scaling regimes for crude-oil prices [25], but H $_q$ increased with q in the long-time scaling regime. K in et al. did not observe a short-time scaling regime for the daily Yen-D ollar exchange rate [22].

2.1 Removal of daily jumps

We analyzed the KOSPI time series to not the origin of multifractality in the Korean stock market. Inspired by Refs. [36,37,38], we removed the daily jumps




Fig. 1. (Color online.) (a) Log-log plot of the generalized height-height correlation function $F_{\rm q}$ (t) versus timet for the prices of the Korean stock-market index, KOSPI with ${\rm q}=0.5;1;2;$; 10 from bottom to top. (b) The generalized Hurst exponents ${\rm H}_{\rm q}$ versus 1=q for the price of KOSPI in the scaling regimes t < t_c () and t > t_c () where t_c ' 40 m in. The error bars here and in Figs. 3 (b) and 4 (c) are standard deviations in the slopes of least-square ts to log-log plots of F $_{\rm q}$ (t).

of the stock index due to the dierence between the closing and opening index. In KOSPI, there was no trading after closing and before opening the market until 1995. So, the changes of the index associated with the daily jumps are small during those years. Since 1995, after market trading and before market




Fig. 2. (Color online.) The time series of the Korean stock-market index, KOSPI: original one-minute time series (light gray, green online), the time series with the daily jumps removed (dark gray, red online), and the time series of the daily jumps only (black).

trading have been allowed for one hour each, leading to substantial overnight jumps. There are additional big daily changes during the period of the A sian nancial crisis around N ovem ber, 1997. In Fig. 2, we present three time series: the original time series, the time series with the daily jumps removed, and the time series of the daily jumps only. In the period between 1992 and 1999, there were bubbles in the K orean stock market around 1995, and there were big crashes around 1997 after the A sian nancial crisis. Larger daily changes can be observed around the period of the A sian nancial crisis.

We show the GHCF for the original KOSPI, the stock index with the daily jumps removed, and the time series of the daily jumps only in Fig. 3(a). In the short-time scaling regime $t < t_c$, the deviations between the original and the modiled data sets increase with increasing q. But in the long-time scaling regime $t > t_c$, the deviations are larger for small q. For large q, the GHCF have large uctuations, and there may be additional scaling regimes for very long times. However, practically no change is observed in Hq, as shown in Fig. 3(b). Removing the daily jumps thus does not delete the multifractal properties of the stock index and only slightly changes the generalized Hurst




Fig. 3. (Color online.) (a) Log-log plots of the generalized height-height correlation function F_q (t) versus time in the time series of the original index (solid lines) and the index with the daily jumps removed (dotted lines) for q=0.5;1;2;; 10 from bottom to top. Inset: the generalized height-height correlation function versus time for the daily jumps. (b) The generalized Hurst exponents H_q versus 1=q for the index with the daily jumps removed in the scaling regimes $t< t_c$ () and $t> t_c$ () where t_c '40 m in. For easy comparison we also include the original index in the scaling regimes $t< t_c$ () and $t> t_c$ () (identical to Fig. 1 (b)).

exponents. In the inset in Fig. 3(a) we present the GHCF of the time series of the daily jumps. We observe obvious multifractality, and there are several scaling regimes.

2.2 Gaussian smoothing

Next, as simply removing the discontinuities has little e ect, we consider the e ects of smoothing the stock-index time series by convolution with a Gaussian of standard deviation 5 m inutes. The convolution of two functions f (t) and g (t) is dened by

$$f' = \int_{1}^{2} f(t)g(t u)du;$$
(3)

where g(t) is a Gaussian convolution kernel [40]. In discrete time series the convolution is a sum instead of an integral

$$\mathbf{f}_{i}^{*} = \sum_{j=1}^{X^{n}} \mathbf{f}_{j} \mathbf{g}_{i \quad j} : \tag{4}$$

In Fig. 4(a) we show a short segment of the original time series and the time series smoothed by convolution with a Gaussian. Although this procedure removes short-time uctuations of the stock index, global trends remain. In Fig. 4(b) we show the GHCF before and after the convolution. Before the convolution, the GHCF shows multifractality. After the convolution we observe monofractal behavior in the short-time scaling regime, t< t_c . In this scaling regime, the lines of the log-log plots are all parallel, with the same slope, $H_q(t < t_c) = 0.986(4)$ 1, independent of q, as shown in table 1. $H_q = 1$ means that the convoluted time series is a smooth curve. The result of H_q 1 for $t < t_c$ and unchanged H_q for $t > t_c$ agrees with those obtained from Gaussian smoothing of simulated hydrogel surfaces in Ref. [37]. For $t > t_c$, the convoluted time series yield the same slopes as the original time series. The corresponding values of H_q are shown vs 1=q in Fig. 4(c).

We also checked the elects of varying the standard deviation in the Gaussian function between 1 m in and 100 m in.W hen the standard deviation is close to the time interval of the stock-index time series, we found that the multifractal behaviors are the same as those of the original time series. When the standard deviation is greater than 4 m inutes and less than the crossover time $t_{\rm c}=40$ m inutes, the multifractal behavior is the same as the results obtained with a standard deviation of 5 m inutes in Fig. 4: H $_{\rm q}$ is close to one for t < $t_{\rm c}$, and it depends on q for t > $t_{\rm c}$. When the standard deviation is greater than 40




Fig. 4. (Color online.) (a) A portion of the original stock index (black) and the time series convoluted with a Gaussian (gray, green online). (b) Log-log plots of the generalized height-height correlation function $F_{\rm q}$ (t) versus time for the original time series (solid curves) and the time series smoothed by Gaussian convolution (dotted curves) for q=0.5;1;2; ;10 from bottom to top. The same gray scales (colors online) correspond to the same values of q. (c) The generalized Hurst exponents H $_{\rm q}$ versus 1=q for the time series smoothed by Gaussian convolution in the scaling regimes t < t_c () and t > t_c (), where t_c ' 40 m in. For easy comparison we also include the original index in the scaling regimes t < t_c () and t > t_c () (identical to Fig. 1 (b)).

m inutes, the crossover time increases proportional to the standard deviation. In this case, the convoluted time series deviate from the global trends of the original time series. O ther functions than G aussian in the convolution, such as a Lorentzian, do not change the results signicantly. From these observations, we conclude that them ultifractality in the short-time regime, $t < t_c$, originates from the local actuations of the stock index. The convolution removes the local actuations and smooths out the variations. However, in the long-time regime, $t > t_c$, there are no dierences between the convoluted time series and the original time series in the aspect of multifractality. In this regime multifractality may be caused by the nonlinearities and the herding dynamics of the stock index [27,41,42]. The behavior of the GHCF for the convoluted time series removing the daily jumps is similar to the original convoluted time series keeping the daily jumps. The multifractality in the long-time regime is robust with respect to smoothing of the time series, and it appears to be an intrinsic property of the stock-market dynamics [43].

Table 1 The generalized Hurst exponents of original and convoluted stock index for $t < t_c$

q	original stock index H $_{\rm q}$ (t < t $_{\rm c}$)	convoluted stock index H $_{\rm q}$ (t < t $_{\rm c}$)
0.5	0.80(1)	0.987(2)
1	0.78(1)	0.986(3)
2	0.73(1)	0.988(2)
3	0.60(1)	0.985(4)
4	0.48(1)	0.987(4)
5	0.41(1)	0.985 (5)
6	0.37(1)	0.984(5)
7	0,33 (1)	0.983 (6)
8	0,31 (2)	0.983 (6)
9	0.29(2)	0.983(7)
10	0 28 (2)	0.983 (7)

3 Conclusion

In sum mary, we have studied multifractality of the Korean KOSP I stock index. Multifractality is observed in two scaling regimes in the original time series of the stock index. However, multifractality in the short-time regime can be removed by convoluting the time series by a Gaussian. In the long-time regime the multifractality is unchanged by smoothing the time series. We propose that

multifractality in the short-time regime is caused by the local uctuations in the stock index. However, multifractality in the long-time regime appears to be a result of the complex behaviors in the stock market, such as herding behavior, information outside the market, the long memory of the volatility, and the intrinsic nonlinear dynamics of the market. Understanding the origins of multifractality in the long-time scaling regime remains an active research topic.

A cknow ledgm ents

We appreciate useful correspondence with S.J.M itchell. This work was supported by KOSEF (R05-2003-000-10520-0 (2003)) and by U.S.N ational Science Foundation Grant Nos. DMR-0120310 and DMR-0240078. Work at Florida State University was also supported by the School of Computational Science and the center for Materials Research and Technology.

R eferences

- [1] R. N. M antegna and H. E. Stanley, An Introduction to Econophysics: Correlations and Complexity in Finance, Cambridge University Press, Cambridge, 1999.
- [2] B. Mandelbrot, Fractals and Scaling in Finance, Springer, New York, 1997.
- [3] J.P.Bouchaud and M.Potters, Theory of Financial Risk, Cambridge University Press, New York, 2000.
- [4]D.Somette, Phys. Rep. 378 (2003) 1.
- [5] R.N.M antegna and H.E.Stanley, Nature 376 (1995) 46.
- [6] J.P. Bouchaud and D. Somette, J. Phys. I France 4 (1994) 863.
- [7] L. Bachelier, Ann. Sci. Ecole Norm. Sup. 3 (1900) 21.
- [8] B.M andelbrot, J.Business 36 (1963) 294.
- [9] E.F. Farm a, J. Business 36 (1963) 420.
- [10] P. Gopikrishnan, V. Plerou, L. A. N. Amaral, M. Meyer, and H. E. Stanley, Phys. Rev. E 60 (1999) 5305.
- [11] P. Gopikrishan, M. Meyer, L. A. N. Amaral, and H. E. Stanley, Eur. Phys. J. B. 3 (1999) 139.
- [12] Y. Liu, P. Gopikrishnan, P. Cizeau, M. Meyer, C.-K. Peng, and H. E. Stanley, Phys. Rev. E 60 (1999) 1390.

- [13] P. Gopikrishnan, V. Plerou, Y. Liu, L. A. N. Amaral, X. Gabaix, and H. E. Stanley, Physica A 287 (2000) 362.
- [14] H.E. Stanley, L.A.N. Amaral, P.Gopikrishnan, and V.Plerou, Physica A 283 (2000) 31.
- [15] H.-J. Kim, I.-M. Kim, and B. Khang, J. Kor. Phys. Soc. 40 (2002) 1105.
- [16] G. Bonanno, G. Caldarelli, F. Lillo, and R. N. Mantegna, Phys. Rev. E 68 (2003) 046130.
- [17] Y. Nakajima, J. Kor. Phys. Soc. 40 (2002) 1096.
- [18] S.Y. Park, S.J. Koo, K.E. Lee, J.W. Lee, and B.H. Hong, New Physics 44 (2002) 293.
- [19] K.E.Lee and J.W.Lee, J.Kor. Phys. Soc. 44 (2004) 668.
- [20] A. Z. Gorski, S. Drodz, and J. Septh, Physica A 316 (2002) 496.
- [21] B.H.W ang and P.M. Hui, Eur. Phys. J. B. 20 (2001) 573.
- [22] K. Kim, J.-S. Choi, and S.-M. Yoon, J. Kor. Phys. Soc. 44 (2004) 643.
- [23] J.A. Skjeltorp, Physica A 283 (2000) 486.
- [24] K.Kim and S.M. Yoon, Fractals 11 (2003) 131.
- [25] J. A Lvarez-Ramirez, M. Cisneros, C. Ibarra-Valdez, A. Soriano, and E. Scalas, Physica A 313 (2002) 651.
- [26] Z.X in-Tian, H.X iao-Yuan, and S.Yan-Li, Physica A 333 (2003) 293.
- [27] Z. Eisler and J. Kertesz, Physica A 343 (2004) 603.
- [28] A. Bershadskii, Physica A 317 (2003) 591.
- [29] M. Ausloos and K. Ivanova, Comp. Phys. Comm. 147 (2002) 582.
- [30] K. Ivanova and M. Auslos, Eur. Phys. J. B 8 (1999) 665.
- [31] A. Turiel and C. J. Perez-Vicente, Physica A 322 (2003) 629.
- [32] H. Katuragi, Physica A 278 (2000) 275.
- [33] A. Bershadskii, J. Phys. A 34 (2001) L127.
- [34] K. Matia, Y. Ashkenazy, and H. E. Stanley, Europhys. Lett. 61 (2003) 422.
- [35] Z. Xu and R. Gencay, Physica A 323 (2003) 578.
- [36] G.M. Buend a, S.J. Mitchell, and P.A. Rikvold, Phys. Rev. E 66 (2002) 046119.
- [37] G.M. Buend a, S.J. Mitchell, and P.A. Rikvold, Microelectronics J. in press.
- [38] S.J.M itchell, cond-m at/0210239.

- [39] P. Meakin, Fractals, scaling and growth far from equilibrium, Cambridge University Press, Cambridge, 1998.
- [40] W . H . P ress, S. A. Teukolsky, W . T. Vetterling, and B. P. Flannery, Numerical Recipes in Fortran: The art of scientic computing, Cambridge University Press, Cambridge, 1986, pp 492.
- [41] A.K raw iecki, J.A. Holyst, and D. Helbing, Phys. Rev. Lett. 89 (2002) 158701.
- [42] M . Barotolozzi and A . W . Thom as, Phys. Rev. E 69 (2004) 046112.
- [43] The perform ance of our analysis code was checked with a synthetic, monofractal time series, for which it showed no indication of multifractality.