Nonlinear Dynamics of Quantum Systems and Soliton Theory

E. Bettelheim,¹ A. G. Abanov,² and P. Wiegmann^{1,3}

¹James Frank Institute, University of Chicago, 5640 S. Ellis Ave. Chicago IL 60637.

²Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794-3800.

³Also at Landau Institute of Theoretical Physics.

(Dated: December 23, 2021)

We show that space-time evolution of one-dimensional fermionic systems is described by nonlinear equations of soliton theory. We identify a space-time dependence of a matrix element of fermionic systems related to the *Orthogonality Catastrophe* or boundary states with the τ -function of the modified KP-hierarchy. The established relation allows to apply the apparatus of soliton theory to the study of non-linear aspects of quantum dynamics. We also describe a *bosonization in momentum* space - a representation of a fermion operator by a Bose field in the presence of a boundary state.

I. INTRODUCTION

In the seminal paper by Date, Jimbo, Kashiwara and Miwa [1] Sato's approach [2] to integrable hierarchies of soliton equations has been formulated in terms of auxiliary fermions. In this formulation the hidden symmetries of soliton equations become explicit symmetries of free fermions, the modes of which are labelled by one-dimensional parameter with respect to the algebra of fermionic bilinear forms $-gl(\infty)$. It was shown that the τ -function of soliton theory can be expressed as a certain matrix element of 1D-fermions. The paper [1] and subsequent works of the Kyoto school (see [3] and references therein) laid out a mathematical foundation of *bosonization* — a commonly used representation of fermionic modes in terms of a Bose field. These works established a deep relation between quantum physics and classical nonlinear equations.

In this paper we adopt this theory to a particularly interesting area of electronic physics - dynamics of fermionic coherent states propagating in real space and time generated by the Galilean Hamiltonian

$$H = \sum_{p} \frac{p^2}{2M} \psi_p^{\dagger} \psi_p. \tag{1}$$

We will show that the real time evolution of important objects of electronic physics also obeys classical non-linear soliton equations. Therefore the powerful apparatus of soliton theory can be used to study complex aspects of non-equilibrium physics of effectively one-dimensional quantum systems. We will report some applications of the results of this paper to non-linear electronic transport elsewhere. Here we focus on the formal connection of soliton theory and quantum dynamics. In this course we encounter an alternative *bosonization* with respect to the boundary state, but now in momentum space.

Before formulating our main results, we introduce notations and give a brief review of some important facts on a connection between solutions of soliton equations and fermionic matrix elements. The literature on the subject is enormous, we restrict the reference list to the original papers [1, 3].

A. Bilinear identity

Consider a Fermi-system on a unit circle or, equivalently, 1D fermions subject to periodic boundary conditions. We let ψ_p be a fermionic mode of the fermionic field $\psi(x) = \sum_p e^{ipx}\psi_p$, labelled by integer momenta, p. The conjugated field is given by $\psi^{\dagger}(x) = \sum_p e^{-ipx}\psi_p^{\dagger}$ with basic (anti-) commutation relations $\{\psi_p, \psi_q\} = \{\psi_p^{\dagger}, \psi_q^{\dagger}\} = 0$ and $\{\psi_p, \psi_q^{\dagger}\} = \delta_{p,q}$. The ground state of the system is a filled Fermi-sea such that all modes in the interval between the Fermi-points are filled. We will be interested only in one, say, the right chiral sector where all essential modes are concentrated around the Fermi point $+p_F$. It is convenient to count modes from this Fermi point and change $p \to p - p_F$. Then the vacuum is the state with all non-positive momenta filled by fermions. It is defined as the state $|0\rangle$ such that:

$$\psi_p|0\rangle = 0, \text{ if } p > 0, \qquad \psi_p^{\dagger}|0\rangle = 0, \text{ if } p \le 0.$$
 (2)

A state, where in addition to the filled negative modes, m > 0 (-m > 0) consecutive positive (negative) modes are also filled (empty) is denoted by $|m\rangle (|-m\rangle)$.

$$\langle m|g = \langle m|e^{\sum_{p,q} A_{pq}:\psi_p^{\dagger}\psi_q:}.$$
(3)

A coherent state represents particle-hole wave-packets. It is characterized by a infinite matrix A_{pq} . The colon in (3) and throughout the paper denotes the normal ordering with respect to the vacuum (2), : $\psi_p^{\dagger}\psi_q := \psi_p^{\dagger}\psi_q - \langle 0|\psi_p^{\dagger}\psi_q|0\rangle$.

Within the algebra of fermionic bilinears one distinguishes two commutative subalgebras. One is generated by positive k > 0 (or by negative k < 0) modes of the current

$$J_{k} = \sum_{p} : \psi_{p}^{\dagger} \psi_{p+k} :, \quad [J_{k}, J_{-k'}] = k \delta_{kk'}.$$
(4)

This subalgebra is central in the construction of Ref. [1, 3]. The other commutative subalgebra will be central in the construction of the current paper. It is defined in (13).

An important result [1, 3] is that the matrix element

$$\tau_m(\mathbf{t}) = \langle m | g \; e^{-\sum_{k>0} t_k J_{-k}} | m \rangle \tag{5}$$

considered as a function of an infinite number of continuous parameters $\mathbf{t} = \{t_k\}$ and of an integer parameter m is the tau-function of an integrable hierarchy. τ_m depends also on the state $\langle g |$ (or on the matrix A_{pq}), but this dependence will be omitted in notations throughout the paper. This means that the family of matrix elements $\tau_m(\mathbf{t})$ obeys a bilinear identity:

$$\oint_{0} e^{\sum_{k>0} (t_{k} - t_{k}') z^{k}} \tau_{m} \left(\mathbf{t} - [z] \right) \tau_{m+n} \left(\mathbf{t}' + [z] \right) z^{n} dz = 0, \tag{6}$$

where $\mathbf{t} \pm [z] \equiv \{t_k \pm z^{-k}/k\}\ k = 1, 2, \dots$ and \mathbf{t}, \mathbf{t}' are two independent sets of parameters. The integral goes along a small contour around the origin.

The bilinear identity (6) encodes an infinite set of hierarchically structured non-linear differential equations in the variables (flows) t_k . For a given integer n, this set is called the *n*-MKP (modified KP) hierarchy. The differential equations with respect to the flows t_k in a bilinear (Hirota) form are obtained by expanding the integrand of (6) in powers of z^{-1} and setting $t'_k = t_k$.

The KP-hierarchy (n = 0) consists of a set of differential equations on τ_m . The first equation of the KP-hierarchy reads

$$(D_1^4 + 3D_2^2 - 4D_1D_3)\tau_m \cdot \tau_m = 0, \tag{7}$$

where D_k is the Hirota derivative in the variable t_k defined as $D_k^n a \cdot b = (\partial_{t_k} - \partial_{t'_k})^n a(\mathbf{t}) b(\mathbf{t}')|_{t'_k = t_k}$. This equation is a bilinear form of the Kadomtsev-Petviashvili equation of plasma physics [1]. It gave the name (KP) to the hierarchy.

The MKP-hierarchy involves a discrete evolution in *m*. The first equation of the 1-MKP-hierarchy reads

$$(D_2 + D_1^2)\tau_{m+1} \cdot \tau_m = 0.$$
(8)

B. Evolution in the space of parameters

Some physical applications arise if $t_k = -t_{-k}^*$. We write $t_k = i\delta t A_{-k}$ and notice that the operator $\sum_k t_k J_k = i\delta t \int A(x)\rho(x)(dx/2\pi)$ represents the action of the scalar electromagnetic potential $A(x) = i\sum_k t_{-k}e^{ikx}$

$$e^{-\sum_{k} t_k J_k} = e^{-i\delta t \int A(x)\rho(x) dx}, \quad \rho(x) =: \psi^{\dagger}(x)\psi(x):, \tag{9}$$

which was turned on instantaneously for a short period δt . This matrix element is relevant, e.g., for the process when an electronic system being initially in an excited state $\langle g| = \langle 0|g$ is instantaneously hit by radiation. The tau-function then is the amplitude of the probability to find a system in the vacuum (ground) state.

An alternative interpretation of the formal expression (5), common in the literature, emphasizes the commutativity of flows with respect to parameters t_k . In this interpretation the operator $e^{\sum_k t_k J_{-k}}$ is viewed as an evolution operator with a set of "times" t_k corresponding to a family of mutually commuting "Hamiltonians" J_k . The excited state $\langle g | = \langle m | g$ evolves in "times". The tau-function is an overlap between the "evolved" excited state $\langle g(\mathbf{t}) |$ and the vacuum

$$\tau_m = \langle g(\mathbf{t}) | m \rangle, \quad \langle g(\mathbf{t}) | = \langle m | g \; e^{-\sum_k t_k J_{-k}}. \tag{10}$$

This terminology, although common, is somewhat confusing. Of course, the "Hamiltonians", J_k are not Hermitian and there is no energy associated with this evolution. The "evolution" occurs in the space of parameters. This may be a reason why applications of the powerful mathematical apparatus of soliton theory to electronic physics have been limited.

In the next subsection we introduce the tau-function as a matrix element of the real-time evolution generated by the Hermitian Hamiltonian (1).

C. Evolution in real time

In physics one is interested in the real-time evolution driven by a Hermitian positive Hamiltonian as (1) rather than currents (4).

Can the theory of solitons be used to study the dynamics of fermions in the real space-time, when the evolution is described by a unitary operator

$$e^{-iHt+iPx} \tag{11}$$

corresponding to a positive energy (1) and momentum

$$P = \sum_{p \in I} p : \psi_p^{\dagger} \psi_p : .$$
⁽¹²⁾

In this case the vacuum is selected as a ground state of the Hamiltonian

$$H|0\rangle = 0$$

Indeed, the set of Hermitian Hamiltonians

$$H_k = \sum_{p \in I} p^k : \psi_p^{\dagger} \psi_p :, \quad k = 1, 2, \dots$$
(13)

generates another commutative subalgebra of fermionic bilinears. Here we denote the set of quantized momenta for the system by I.

Surprisingly, this question has never been addressed systematically. The most relevant papers we know are Refs. [4, 5] where the relation of time dependent Green functions of impenetrable bosons and Green functions of Ising model to the Painlevé equations has been established.

In this paper we will show that, indeed, an action of the commutative Hamiltonians (13) generates an integrable hierarchy. Specifically, we show that the matrix element

$$\tau_m(\mathbf{t}) = \langle 0|g \ e^{-\sum_{k>0} t_k H_k} : e^{-(a+m) \,\varphi(x)} : |0\rangle, \quad 0 < |a| < 1/2, \quad m \text{ is integer}$$
(14)

obeys the MKP hierarchy 1 .

Namely, the tau-functions $\tau_m(\mathbf{t})$, similarly to (5), satisfy a set of equations analogous to (7, 8). These equations are generated by the identity:

$$\oint_{0} e^{\sum_{k>0} (t_{k} - t'_{k}) z^{k}} \frac{\Gamma(z + m + n)}{\Gamma(z + m)} \tau_{m} \left(\mathbf{t} - [z] \right) \tau_{m+n} \left(\mathbf{t}' + [z] \right) dz = 0, \tag{15}$$

which is a modified form of (6). The factor $\frac{\Gamma(z+m+n)}{\Gamma(z+m)} = \prod_{j=m}^{m+n-1} (z+j)$ replaces z^n in (6)².

¹ Here we take $a \neq 0$. Although the tau-function obeys the same *a*-independent equations, it degenerates at a = 0 and can be computed by elementary means.

² The difference between (15) and (6) is inessential insofar as the former may be transformed into the latter by the transformation $\tau_m \to \tau_m e^{\sum_{k\geq 0} V(m+k)}$, where $V(z) = \sum_k t_k z^k$, however, this transformation changes analytical properties of the tau-function as a function of the "times".

In the eq. (14) the operator φ is defined as a Bose field

$$\varphi(x) = \sum_{k \neq 0} \frac{1}{k} e^{ikx} J_k + i J_0 x \tag{16}$$

and

$$:e^{a\varphi(x)}:=e^{a\varphi_{-}(x)}e^{iaxJ_{0}}e^{a\varphi_{+}(x)}.$$
(17)

where $\varphi_{\pm}(x) = \sum_{\pm k>0} (e^{ikx}/k) J_k$ contain only positive or negative modes respectively. Note that the flow t_1 just shifts the coordinate x by it_1 as $e^{t_1H_1}\varphi(x)e^{-t_1H_1} = \varphi(x+it_1)$, so we may set x = 0 in some formulas below. In physical applications we identify

time:
$$t = it_2$$
, space coordinate in the Galilean frame: $x - v_F t = -it_1$, (18)

where $v_F = p_F/M$ is a Fermi velocity.

The operator : $e^{a \varphi(x)}$: in (14) is called *a boundary condition changing operator*. This operator acting on a vacuum creates *a boundary state*

$$|B_m(x)\rangle =: e^{-(a+m)\varphi(x)} : |0\rangle.$$
⁽¹⁹⁾

The meaning of the boundary condition changing operator is clarified by the action of the counting operator $n(y) = \int_0^y (:\psi^{\dagger}(y')\psi(y'):-(a+m))dy'$ on the boundary state

$$\langle B_m(x)|n(y)|B_m(x)\rangle = (a+m) \quad \text{if } 0 < x < y, \quad \text{or } 0 \quad \text{otherwise.}$$

$$\tag{20}$$

In other words, the boundary condition changing operator moves particles towards the point x creating an increase of the density $\rho(y) = \partial_y n(y)$ at that point.

Boundary states play an important role in 2D-critical phenomena in systems with boundaries. The connection of the latter to 1D-fermions is following. The fields (e.g. $\psi(x)$ and $\varphi(x)$) are originally defined on the unit circle. One views the circle as a contour embedded in the complex plane z. The fields and their differentials can be analytically extended to the exterior of the circle and to its interior. Instead of considering fields defined at the point z' in the interior of the circle one views them as defined at the point $z = 1/\overline{z'}$ of the circle's exterior (Schwarz reflection). From this point of view the circle $z = e^{-ix}$, $x \in \mathbb{R}$, is seen as a boundary of the holomorphic 2D-field theory defined in its exterior.

In electronic physics boundary states appear in numerous problems related to the Fermi-edge singularities, quantum impurity problems and other phenomena related to Orthogonality Catastrophe ([10], see, e.g., [8]). We will discuss some applications to electronic physics elsewhere [14].

In terms of the boundary state, our tau-function (14) reads

$$\tau_m(\mathbf{t}) = \langle G(\mathbf{t}) | B_m \rangle, \quad \langle G(\mathbf{t}) | = \langle 0 | g \, e^{-\sum_{k>0} t_k H_k}. \tag{21}$$

The tau-function depends on two parameters: a fraction a, and an integer m. Below we fix a and consider m as a running parameter.

One can think of this matrix element as follows. An excited coherent state $\langle g |$ evolves up to a time t and then is measured at a point x by a projection onto a boundary state $|B_a(x)\rangle$.

A comment is in order. The presence of the boundary operator : $e^{i\varphi(x)}$: changes the momentum quantization. The eigenvalues p of the momentum operator (12) appearing in (12, 13) are no longer integer, but are integers shifted by a:

$$p \in \mathbb{Z} + a,\tag{22}$$

that is I in (12) and (13) is given by $I = \mathbb{Z} + a$.

An obvious generalization of this result is a passage from the KP to the 2D Toda hierarchy [9]. The matrix element

$$\langle B_m | e^{-\sum_k \bar{t}_k H_k} g e^{-\sum_k t_k H_k} | B_m \rangle$$

obeys the 2D-Toda hierarchy with respect to two sets of parameters t_k and \bar{t}_k and a discrete parameter m. Other obvious generalization occurs when fermions are placed on a spatial lattice so that the energy spectrum is a periodic function of momentum. We do not discuss these generalizations here.

$$(iD_t + D_x^2 - 2imD_x)\tau_{m+1} \cdot \tau_m = 0.$$
⁽²³⁾

The independent variables x and t in this equation are physical space and time coordinates of a 1D fermionic system. Among the hierarchy, this equation seems to be the most important for the study of the dynamics of electronic systems. The two functions τ_m and τ_{m+1} are not independent but are connected by certain analytical properties, thus making equation (23) closed. In the most interesting situations (corresponding to particular choices of coherent state g) the analytical conditions are explicit and correspond to a certain reduction of the MKP-hierarchy. We consider coherent states having particle-hole symmetry and the corresponding reduction in Appendix A.

We note here that the matrix element (14) give solutions of the MKP hierarchy, which are essentially different from the ones given by (5). For example, we show below that (14) has only positive Fourier modes with respect to all odd flows it_1, it_3, \ldots This analytical condition is restrictive. In particular, it does not admit soliton solutions. In contrast, it is well known that (5) can generate soliton solutions (see Sec.III).

Some mathematical constructions related of the matrix element (14), although devoted to different problems, have appeared recently in papers by A. Orlov [6] and by A. Okounkov, R. Pandharipande, and N. A. Nekrasov [7].

II. HIROTA EQUATION AND BOSONIZATION

Before we proceed, we present another form of the bilinear identity (15) for a particular case n = 1. Setting $\mathbf{t} = \mathbf{t}^{(0)} + [p], \mathbf{t}' = \mathbf{t}^{(0)} - [q]$ the factor $(z+m)e^{\sum_k (t_k-t'_k)z^k}$ becomes $\frac{(z+m)pq}{(q-z)(p-z)}$. The integrand has three poles at $z = 0, p, q, \infty$ and can be computed. It gives the discrete Hirota equation

$$(p+m)\,\tau_m\cdot\tau_{m+1}([p]-[q])-(q+m)\,\tau_m([p]-[q])\cdot\tau_{m+1}-(p-q)\,\tau_m([p])\cdot\tau_{m+1}(-[q])=0,$$
(24)

where we have dropped the term $\mathbf{t}^{(0)}$ in each of the arguments of the τ functions. We stress that p and q are shifted integers $p, q \in \mathbb{Z} + a$.

Hirota's equation, written in the discrete form (24) appears as a particular specification of the integral form of the bilinear identity (6). In fact, they carry the same information as an integral form of the bilinear identity written for an arbitrary n. The latter can be obtained from the former [2]. In the rest of this section we will derive the discrete form of Hirota's equation (24) for tau-functions defined in (14). It will be equivalent to proving (15).

Our main equation (23) directly follows from Hirota equation written in the form (24). It appears in the leading order expansion at large q and p, and setting q = -p.

A. Bosonization in coordinate space and the tau-function $\tau_m = \langle g(\mathbf{t}) | m \rangle$

First, we show how to derive the discrete Hirota equation (25) for the the conventional matrix element matrix (5). Then we derive the discrete Hirota equation (24) for our tau-function (14). The first derivation is well known [1, 2, 3, 12]. We, nevertheless, present it here in order to use it as a guide for the following derivation of Hirota's equation for the real time evolution. The comparison of these two derivations is instructive.

Applying the same trick as is the previous section to (6) we obtain a discrete Hirota equation for the tau-function (5)

$$p \tau_m \cdot \tau_{m+1}([p] - [q]) - q \tau_m([p] - [q]) \cdot \tau_{m+1} - (p-q) \tau_m([p]) \cdot \tau_{m+1}(-[q]) = 0.$$
(25)

To derive it one needs two technical ingredients: bosonization and the Wick's theorem.

Bosonization formulas

The Bosonization formulas express the action of fermion operators in terms of bosons. We will need the following formulas (briefly derived below)

$$\psi^{\dagger}(x)\psi(y)|m\rangle = \frac{1}{1 - e^{i(x-y)}} : e^{\varphi(y) - \varphi(x)} : |m\rangle,$$
(26)

$$\psi(x)|m+1\rangle = e^{ix} : e^{\varphi(x)} : |m\rangle, \qquad \psi^{\dagger}(x)|m\rangle = : e^{-\varphi(x)} : |m+1\rangle.$$
(27)

Let us now use these bosonization formulas to "fermionize" the tau-function $\tau_{m+1}([p] - [q])$. We set $q = e^{ix}$ and $p = e^{iy}$, and write the tau-function $\tau_{m+1}([p] - [q]) = \langle m+1 | g \ e^{-\sum_k t_k^{(0)} J_{-k}} : e^{\varphi(y) - \varphi(x)} : |m+1\rangle$, appearing in (25), in terms of fermion operators:

$$\tau_{m+1}([p] - [q]) = e^{i(m+1)(x-y)} (1 - e^{i(x-y)}) \langle m+1|g(\mathbf{t})\psi^{\dagger}(x)\psi(y)|m+1\rangle, \quad g(\mathbf{t}) = g \ e^{-\sum_{k} t_{k}^{(0)} J_{-k}}.$$
(28)

Wick's theorem

We apply Wick's theorem to four fermion matrix elements. Let χ_i be a linear combination of fermionic modes $\chi_i = \sum_q v_q^{(i)} \psi_q$, and $|G_2\rangle$ and $\langle G_1|$ are coherent states (i.e. states of the form (3)). The general form of Wick's theorem for four fermion matrix elements reads

$$\langle G_1 | \chi_1^{\dagger} \chi_2 \chi_3^{\dagger} \chi_4 | G_2 \rangle \langle G_1 | G_2 \rangle = \langle G_1 | \chi_1^{\dagger} \chi_2 | G_2 \rangle \langle G_1 | \chi_3^{\dagger} \chi_4 | G_2 \rangle + \langle G_1 | \chi_1^{\dagger} \chi_4 | G_2 \rangle \langle G_1 | \chi_2 \chi_3^{\dagger} | G_2 \rangle.$$

$$\tag{29}$$

A proof of the Hirota bilinear identity (25)

We apply Wick's theorem to the matrix element:

$$< m |\psi_{m+1} g(\mathbf{t}) \psi^{\dagger}(x) \psi(y) \psi^{\dagger}_{m+1} |m\rangle \langle m |g(\mathbf{t})|m\rangle = \langle m+1 |g(\mathbf{t})|m+1\rangle \langle m |g(\mathbf{t}) \psi^{\dagger}(x) \psi(y)|m\rangle + \langle m+1 |g(\mathbf{t}) \psi^{\dagger}(x)|m\rangle \langle m |g(\mathbf{t}) \psi(y)|m+1\rangle.$$
(30)

Eq. (25) emerges if one writes each of the matrix elements appearing in this identity as τ -functions with shifted arguments, by further use of the bosonization formulas (26,27)³.

Derivation of bosonization formulas (26,27)

It will be instructive to have a short proof of bosonization formulas (26,27) as we are going to use a similar approach in the next section. In the following we set for simplicity m = 0.

Let us project the r.h.s. and the l.h.s. of (26) onto a general state with a given number of particles $\langle \{x_i, y_i\} | = \langle 0 | \left(\prod_{i=1}^N \psi^{\dagger}(x_i)\psi(y_i)\right)$, consisting of N particle and N hole excitations with coordinates x_i, y_i , and compare the results.

First of all, let us use Wick's theorem to calculate the overlap between the state with N particle-hole pairs and the vacuum. This overlap is a determinant of a single-particle Green's function $\langle 0|\psi^{\dagger}(x)\psi(y)|0\rangle = (1 - e^{i(x-y)})^{-1}$

$$\langle \{x_i, y_i\} | 0 \rangle = e^{i \sum_{i=1}^{N} y_i} \det_{i,j \le N} \left(\frac{1}{e^{iy_i} - e^{ix_j}} \right) = e^{i \sum_{i=1}^{N} y_i} \frac{\prod_{i < j \le N} (e^{iy_i} - e^{iy_j}) (e^{ix_j} - e^{ix_i})}{\prod_{i,j \le N} (e^{iy_i} - e^{ix_j})},$$
(31)

where we used the multiplicative form of the Cauchy determinant

$$\det_{i,j \le N} \left(\frac{1}{q_i - p_j} \right) = \frac{\prod_{i < j \le N} (q_i - q_j)(p_j - p_i)}{\prod_{i,j \le N} (q_i - p_j)}.$$
(32)

The l.h.s. of (26) is computed by extending the previous formula to N+1 particle-hole pairs and setting $x_{N+1} = x$ and $y_{N+1} = y$

$$\langle \{x_i, y_i\} | \psi^{\dagger}(x) \psi(y) | 0 \rangle = \frac{e^{iy}}{e^{iy} - e^{ix}} \prod_{i \le N} \frac{(e^{iy} - e^{iy_i})(e^{ix} - e^{ix_i})}{(e^{iy} - e^{ix_i})(e^{ix} - e^{iy_i})} \,\langle \{x_i, y_i\} | 0 \rangle. \tag{33}$$

³ This proof appeals to a general view on tau functions as Plücker coordinates of the Grassmann manifold [2].

The r.h.s. of (26) can be computed by (i) noticing that $e^{b\varphi}|0\rangle = e^{b\varphi_-}|0\rangle$, (ii) pulling the vertex operator $e^{b\varphi_-}$ to the left using the identity

$$e^{-b\varphi_{-}(x)}\psi^{\dagger}(x_{i})\psi(y_{i})e^{b\varphi_{-}(x)} = \left(\frac{e^{ix} - e^{iy_{i}}}{e^{ix} - e^{ix_{i}}}\right)^{b}\psi^{\dagger}(x_{i})\psi(y_{i})$$
(34)

and then (iii) using $\langle 0|e^{b\varphi_{-}(x)}=0$. As a result we get

$$\langle \{x_i, y_i\} | : e^{\varphi(y) - \varphi(x)} : |0\rangle = \prod_{i \le N} \frac{(e^{iy} - e^{iy_i})(e^{ix} - e^{ix_i})}{(e^{iy} - e^{ix_i})(e^{ix} - e^{iy_i})} \,\langle \{x_i, y_i\} | 0\rangle.$$
(35)

Comparing eqs. (35) and (33) gives the bosonization formula (26). Eq. (27) can be derived in a similar way, or by setting $x = -i\infty$ in (26) and using the fact that $\psi^{\dagger}(-i\infty)|m\rangle = |m+1\rangle$. In all formulas x and y carry small positive imaginary parts.

B. Bosonization in momentum space and the tau-function $\tau_m = \langle G(\mathbf{t}) | B_m \rangle$

Now we show that the matrix element (14) obeys the Hirota equation (24). The steps of the derivation in this section follow the steps of the previous section, but technically different. We repeat that in this section all momenta are shifted integers $p, q \in \mathbb{Z} + a$.

The important property of the boundary state is displayed by the form of its wave-function in momentum space. We have $\langle 0|B_m \rangle = 1$ and two important formulas

$$\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle = f_m^+(p)\frac{1}{p-q}f_m^-(q), \quad q \le a < p,$$
(36)

$$\frac{p+m}{q+m}\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle = \langle 0|\psi_q^{\dagger}\psi_p|B_{m+1}\rangle, \tag{37}$$

where

$$f_m^+(p) = \frac{\Gamma(p+m)}{\Gamma(m+a)\Gamma(p-a)},$$

$$f_m^-(q) = \frac{\Gamma(1-q-m)}{\Gamma(1-m-a)\Gamma(1-q+a)}.$$
(38)

The matrix element (36) is also non zero at $p = q \leq a$, where it is equal to 1⁴. Eq. (37) follows from (36,38) with the help of the formula

$$f_m^+(p)f_m^-(q)\frac{p+m}{q+m} = f_{m+1}^+(p)f_{m+1}^-(q).$$
(39)

We prove these formulas at the end of this section.

The formula (36) is related to a phenomenon referred to as "Orthogonality Catastrophe" [10]. Namely, if the length of the ring is large, i.e., in the limit $p, -q \gg |m|$ the formula (36) reads

$$\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle = (-1)^m \frac{\sin(\pi a)}{\pi} \frac{1}{p-q} \left(\frac{p}{-q}\right)^{a+m}.$$
(40)

The overlap between the excited state and the boundary state vanishes as the momentum of the excited particle approaches the Fermi-point $\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle \sim p^{a+m}$ as $p \to 0$.

⁴ The formula (36) is understood as being analytically continued from the real axis to the complex plane of p and q. In particular, at p = q it should be understood as a limit $p \to q$ at fixed q.

Consider now a generic excited state $\langle \{p_i, q_i\} | = \langle 0 | \prod_{i=1}^N \psi_{q_i}^{\dagger} \psi_{p_i}$ characterized by a set of N-particles with momenta $p_i > a$ and N holes with momenta $q_i \leq a$. The overlap of this excited state with the boundary state is obtained from the Wick's theorem and eq.(36)

$$\langle \{p_i, q_i\} | B_m \rangle = \det_{i,j \le N} \langle 0 | \psi_{q_j}^{\dagger} \psi_{p_i} | B_m \rangle = \det_{i,j \le N} \left(f_m^+(p_i) \frac{1}{p_i - q_j} f_m^-(q_j) \right)$$

$$= \prod_{i \le N} f_m^+(p_i) f_m^-(q_i) \frac{\prod_{i < j \le N} (p_i - p_j) (q_j - q_i)}{\prod_{i,j \le N} (p_i - q_j)},$$

$$(41)$$

where in the last line we again used the Cauchy determinant formula (32).

Let us note that up to the multiplicative factor $\prod_i f_m^+(p_i)f_m^-(q_i)$, the overlap (41) between the excited state and the boundary state written in momentum space is given by the Cauchy determinant, i.e., has the same form as the similar overlap written in real space (31). The determinants become identical under a substitution $p_i = e^{ix_i}$ and $q_i = e^{iy_i}$.

We observe an interesting phenomenon: the boundary state $|B_m\rangle$ interchanges momentum and coordinate spaces. The transmutation

coordinate space
$$\leftrightarrow$$
 momentum space, (42)

Fermi sea \leftrightarrow Boundary state (42) (43)

is the reason why different matrix elements (5, 14) obey similar non-linear equations. There is no one-to-one correspondence between coordinate and momentum space – the space is continuous and finite while the momentum space is discrete and infinite.

Following the logic of the previous section we proceed in two steps. First, we establish the bosonization formulas in momentum space. They will allow us to rewrite tau-functions in eq. (24) as matrix elements of fermionic operators. Then eq.(24) is equivalent to a particular form of the Wick's theorem.

Bosonization formulas

Let us introduce a bosonic field associated with the set of commuting Hamiltonians (13)

$$\Phi(s) = \sum_{k>0} \frac{s^{-k}}{k} H_k = -\sum_p \ln(1 - \frac{p}{s}) : \psi_p^{\dagger} \psi_p : .$$
(44)

Bosonization in the momentum space is summarized by formulas

$$\psi_{q}^{\dagger}\psi_{p}|B_{m}\rangle = \frac{f_{m}^{+}(p)f_{m}^{-}(q)}{p-q}e^{\Phi(q)-\Phi(p)}|B_{m}\rangle, \tag{45}$$

and

$$\lim_{p \to -m} (p+m)\psi_q^{\dagger}\psi_p | B_m \rangle = (-1)^m \frac{\sin \pi a}{\pi} f_{m+1}^-(q) e^{\Phi(q)} | B_{m+1} \rangle.$$
(46)

These formulas will be derived below. We remark here that (45,46) should be understood as analytically continued formulas for matrix elements. For example, one should first calculate the overlap of the l.h.s. of (46) with some bra-state, then analytically continue it from $p \in \mathbb{Z} + a$ to the complex *p*-plane and only then take a limit $p \to -m$.

Once bosonization formulas in momentum space (45,46) are established we are ready to prove the Hirota equation (24).

Wick's theorem

First, we apply the Wick theorem to the 4-fermion matrix element

$$\langle G(\mathbf{t})|\psi_q^{\dagger}\psi_p\psi_Q^{\dagger}\psi_P|B_m\rangle\langle G(\mathbf{t})|B_m\rangle = \langle G(\mathbf{t})|\psi_q^{\dagger}\psi_p|B_m\rangle\langle G(\mathbf{t})|\psi_Q^{\dagger}\psi_P|B_m\rangle - \langle G(\mathbf{t})|\psi_q^{\dagger}\psi_P|B_m\rangle\langle G(\mathbf{t})|\psi_Q^{\dagger}\psi_P|B_m\rangle.$$
(47)

Next we multiply both sides by $\frac{P+m}{f_{m+1}^-(Q)}$, send $P \to -m$, and use eq. (46). We get

$$\frac{q-Q}{p-Q}\langle G(\mathbf{t})|e^{\Phi(Q)}\psi_{q}^{\dagger}\psi_{p}|B_{m+1}\rangle\langle G(\mathbf{t})|B_{m}\rangle$$

$$= \langle G(\mathbf{t})|\psi_{q}^{\dagger}\psi_{p}|B_{m}\rangle\langle G(\mathbf{t})|e^{\Phi(Q)}|B_{m+1}\rangle - \frac{f_{m+1}^{-}(q)}{f_{m+1}^{-}(Q)}\langle G(\mathbf{t})|e^{\Phi(q)}|B_{m+1}\rangle\langle G(\mathbf{t})|\psi_{Q}^{\dagger}\psi_{p}|B_{m}\rangle.$$
(48)

The next step is to use the bosonization formula (45)

$$\frac{q-Q}{p-Q} \frac{p+m}{q+m} \langle G(\mathbf{t}) | e^{\Phi(Q)} e^{\Phi(q)-\Phi(p)} | B_{m+1} \rangle \langle G(\mathbf{t}) | B_m \rangle$$

$$= \langle G(\mathbf{t}) | e^{\Phi(q)-\Phi(p)} | B_m \rangle \langle G(\mathbf{t}) | e^{\Phi(Q)} | B_{m+1} \rangle - \frac{Q+m}{q+m} \frac{p-q}{p-Q} \langle G(\mathbf{t}) | e^{\Phi(q)} | B_{m+1} \rangle \langle G(\mathbf{t}) | e^{\Phi(Q)-\Phi(p)} | B_m \rangle.$$
(49)

We recognize the matrix elements in (49) as tau-functions (14) with shifted arguments, e.g., $\langle G(\mathbf{t})|e^{\Phi(q)}|B_{m+1}\rangle = \tau_{m+1}(-[q])$. The last step is to take a limit $Q \to \infty$. In this limit $\Phi(Q) \to 0$ and (49) brings us to (24), which completes the proof.

Derivation of formulas for matrix elements (36,38)

Now we prove the formulae (36,38). First, we find using (34) that in coordinate representation

$$\langle 0|\psi^{\dagger}(x)\psi(y)|B_{m}\rangle = \langle 0|\psi^{\dagger}(x)\psi(y)|0\rangle \left(\frac{1-e^{ix}}{1-e^{iy}}\right)^{m+a} = \frac{e^{iy}}{e^{iy}-e^{ix}} \left(\frac{1-e^{ix}}{1-e^{iy}}\right)^{m+a}.$$
(50)

We consider $(p-q)\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle$ and pass to the coordinate representation $(p-q)\langle 0|\psi_q^{\dagger}\psi_p|B_m\rangle \rightarrow (i\partial_x + i\partial_y)\langle 0|\psi^{\dagger}(x)\psi(y)|B_m\rangle$. Then we calculate

$$(i\partial_x + i\partial_y) \left[\frac{e^{iy}}{e^{iy} - e^{ix}} \left(\frac{1 - e^{ix}}{1 - e^{iy}} \right)^{m+a} \right] = (m+a)(1 - e^{ix})^{m+a-1}e^{iy}(1 - e^{iy})^{-(a+m+1)}$$

We observe that the right hand size factorizes. Taking an inverse Fourier transform we obtain (36) with

$$f_m^-(q) = \int_0^{2\pi} e^{iqx} (1 - e^{ix})^{m+a-1} \frac{dx}{2\pi}, \qquad f_m^+(p) = (m+a) \int_0^{2\pi} e^{-i(p-1)y} (1 - e^{iy})^{-(m+a+1)} \frac{dy}{2\pi}.$$

The calculation of these integrals gives (38) for f_m^{\pm} . These formulas can also be found in [11].

Derivation of bosonization formulas (45,46)

To prove the formula (45) we evaluate the projection of its r.h.s. on the state $\langle \{p_i, q_i\} \rangle$. With the help of the formula

$$e^{b\Phi(s)}\psi_q^{\dagger}\psi_p e^{-b\Phi(s)} = \left(\frac{s-p}{s-q}\right)^b \psi_q^{\dagger}\psi_p \tag{51}$$

analogous to (34) and using $\langle 0|\Phi(s)=0$ we obtain

$$\langle \{p_i, q_i\} | e^{\Phi(q) - \Phi(p)} | B_m \rangle = \langle \{p_i, q_i\} | B_m \rangle \prod_{i \le N} \frac{(p - p_i)(q - q_i)}{(p - q_i)(q - p_i)}.$$
(52)

On the other hand, using the multiplicative representation for the l.h.s. of (45) similar to (41) we obtain

$$\langle \{p_i, q_i\} | \psi_q^{\dagger} \psi_p | B_m \rangle = \det_{i,j \le N+1} \left(f_m^+(p_i) \frac{1}{p_i - q_j} f_m^-(q_j) \right) = \langle \{p_i, q_i\} | B_m \rangle \frac{f_m^+(p) f_m^-(q)}{p - q} \prod_{i \le N} \frac{(p - p_i)(q - q_i)}{(p - q_i)(q - p_i)}, \tag{53}$$

where $p_{N+1} = p$ and $q_{N+1} = q$. Comparing (53) and (52) we prove (45).

To obtain (46), we first use (39) to obtain

$$\langle \{p_i, q_i\} | B_{m+1} \rangle = \langle \{p_i, q_i\} | B_m \rangle \prod_{i \le N} \frac{p_i + m}{q_i + m}.$$
(54)

Then we use (54) and (53) to proceed as follows

$$\langle \{p_i, q_i\} | e^{\Phi(q)} | B_{m+1} \rangle = \prod_{i \le N} \frac{q - q_i}{q - p_i} \langle \{p_i, q_i\} | B_{m+1} \rangle = \prod_{i \le N} \frac{q - q_i}{q - p_i} \frac{p_i + m}{q_i + m} \langle \{p_i, q_i\} | B_{m+1} \rangle$$

$$= \langle \{p_i, q_i\} | \psi_q^{\dagger} \psi_p | B_m \rangle \frac{p - q}{f_m^+(p) f_m^-(q)} \prod_{i \le N} \frac{p_i + m}{p_i - p} \frac{q_i - p_i}{q_i + m}.$$
(55)

The r.h.s. of (55) effectively does not depend on p and we take its limit as $p \to -m$. The product in the last line disappears and we conclude that

$$e^{\Phi(q)}|B_{m+1}\rangle = -\frac{q+m}{f_m^-(q)} \lim_{p \to -m} \frac{1}{f_m^+(p)} \psi_q^{\dagger} \psi_p |B_m\rangle.$$
 (56)

Finally, using

$$\lim_{p \to -m} (p+m) f_m^+(p) = (-1)^{m+1} (m+a) \frac{\sin(\pi a)}{\pi},$$
(57)

we arrive to (46).

III. MULTI-PERIODIC SOLUTIONS

In this section we present explicit formulas for multi-phase solutions evolving in real time. These formulas reveal the structure of the tau-function and provide another direct proof of Hirota's equation.

A set of excited states $|\{p_i, q_i\}\rangle$ characterized by momenta of N particles $p_i > a$ and holes $q_i \leq a, i = 1, ... N$ form an orthonormal basis for the Fock space of fermions. In this basis the Hamiltonians (13) are diagonal: $H_k|\{p_i, q_i\}\rangle = \sum_i (p_i^k - q_i^k)|\{p_i, q_i\}\rangle$. Using this basis, we decompose

$$\tau_m = \sum_{N \ge 0} \sum_{\{p_i, q_i\}} e^{-\sum_{k, i} t_k(p_i^k - q_i^k)} \langle g | \{p_i, q_i\} \rangle \langle \{p_i, q_i\} | B_m \rangle.$$
(58)

The matrix elements entering this expression are determinants of two-particle matrix elements. The first one is

$$\langle g|\{p_i,q_i\}\rangle = \det_{i,j} A_{p_i,q_j}, \quad A_{p,q} = \langle g|p,q\rangle.$$
(59)

The second one is the Schur function given by eq. (41)

$$\langle \{p_i, q_i\} | B_m \rangle = \det_{i,j \le N} \left(\frac{1}{p_i - q_j} \right) \prod_{i \le N} f_m^+(p_i) f_m^-(q_i).$$

$$\tag{60}$$

Introducing

$$s_m(\{p_i, q_i\}, g) = \det_{i,j} \left(A_{p_i, q_j}^{(m)} \right)$$

where

$$A_{p,q}^{(m)} = f_m^+(p)A_{p,q}f_m^-(q)$$

we write the tau-function in the form

$$\tau_m = \sum_{N \ge 0} \sum_{\{p_i, q_i\}} e^{-\sum_{k,i} t_k(p_i^k - q_i^k)} s_m(\{p_i, q_i\}, g) \det_{ij} \left(\frac{1}{p_i - q_j}\right).$$
(61)

$$\frac{s_{m+1}}{s_m} = \prod_i \left(\frac{p_i + m}{q_i + m}\right). \tag{62}$$

We do not present this proof of (24) here ⁵.

If the matrix A_{pq} defining the state $\langle g |$ by (59) is a finite matrix, the sum in (61) truncates. It represents a multi-phase solution. In this case the determinant formula holds

$$\tau_m(\mathbf{t},a) = \det_{i,j \le N} \left(\delta_{ij} + f_m^+(p_i) e^{-\epsilon(p_i)} \frac{A_i}{p_i - q_j} e^{\epsilon(q_j)} f_m^-(q_j) \right), \quad \epsilon(p) = \sum_k t_k p^k.$$
(63)

Here $A_i = A_{p_i q_i}$. Note that a similar determinant representation holds for (5). Namely,

$$\tau_m(\mathbf{t}) = \det_{i,j \le N} \left(\delta_{ij} + p_i^{-m} e^{-\epsilon(p_i)} \frac{A_i}{p_i - q_j} e^{\epsilon(q_j)} q_j^{m+1} \right), \quad \epsilon(p) = \sum_k t_k p^k.$$
(64)

Although (63) and (64) are very similar there is an essential difference between them. Namely, in (63) the parameters p_i and q_i are momenta of particles and holes and are subject to "Fermi sea" restriction $p_i > a \ge q_i$. On the other hand, in (64) $p = e^{ix}$ and $q = e^{iy}$ represent complex coordinates of particles and holes and are not subject to this restriction. Due to the restriction on the values of p_i and q_i the tau function (63) has only positive Fourier components in all odd flows.

The simplest 1-phase and 2-phase solutions are written below as a function of space $t_1 = ix$ and time $t_2 = -it$.

- 1-phase solution describes one particle-hole excitation:

$$\tau_m = 1 + A \frac{f_m^+(p) f_m^-(q)}{p - q} e^{i(p - q)(x - vt)}.$$
(65)

The solution is characterized by two positive numbers p and -q from $\mathbb{Z} + a$ labelling particle-hole momenta and an arbitrary *m*-independent constant A, which may depend on p, q. It propagates with velocity v = p + q.

- 2-phase solution (two particle-hole excitations):

$$\tau_m = 1 + A_1 \frac{f_m^+(p_1)f_m^-(q_1)}{p_1 - q_1} e^{i(p_1 - q_1)x - i(p_1^2 - q_1^2)t} + A_2 \frac{f_m^+(p_2)f_m^-(q_2)}{p_2 - q_2} e^{i(p_2 - q_2)x - i(p_2^2 - q_2^2)t}$$
(66)
+ $A_1 A_2 \det_{i,j=1,2} \left(\frac{f_m^-(q_j)f_m^+(p_i)}{p_i - q_j} \right) e^{i(p_1 - q_1)(x - v_1t)} e^{i(p_2 - q_2)(x - v_2t)}.$

The multi-phase solutions in (64) allow for rational limit. Keeping $v_i = p_i + q_i$ constant while taking the limit $p_i - q_i \to 0$, one obtains rational solutions corresponding to N-solitons with v_i being asymptotic velocities of solitons. However, in (63) and its particular cases (65, 66) this limit does not exist. Indeed, under the condition $p_i > a \ge q_i$ the limit $p_i - q_i \to 0$ implies $p_i, q_i \to 0$ and therefore it is impossible to have finite v_i in this limit. We see that the matrix element (14) does not contain soliton solutions of the MKP hierarchy.

IV. CONCLUSION

We have shown that the real time dynamics of certain matrix elements (14) of one-dimensional (non-relativistic) fermions obeys non-linear integrable equations of the modified KP hierarchy. This evolution is driven by a commutative set of Hermitian Hamiltonians (13)

,

$$H_k = \int :\psi^{\dagger}(x) \left(-i\frac{\partial}{\partial x}\right)^k \psi(x) : \frac{dx}{2\pi}$$

⁵ An observation that the series (61) for (14) obeys KP-hierarchy, i.e., eq. (15) at n = 0 has been made in Ref.[6].

$$J_k = \int e^{ikx} : \psi^{\dagger}(x)\psi(x) : \frac{dx}{2\pi}.$$

operators (4)

In these two cases the flows have different meaning. The former are coordinates of physical space-time, while the latter are deformation parameters of the coherent state.

The matrix element we studied in this paper is understood as an overlap of the evolved coherent fermionic state $\langle G(\mathbf{t})|$ with a "probe" boundary state $|B_m\rangle$. This matrix element appears in variety of physical problems involving the phenomena referred to as the Orthogonality Catastrophe. Among them are problems related with the Fermi-edge singularity, quantum impurity and tunneling in effectively one dimensional systems.

The fact that questions of quantum dynamics in one dimension are linked to the tau-function, indicates that the kinetics are essentially non-linear and are a subject to instabilities inherited from the non-linear dynamics. In a forthcoming paper [14] we describe the Whitham approach to shock wave type solutions for electronic systems.

An essential element of our proof of the MKP hierarchy is the representation of a fermionic mode by a Bose field (44,46) in the action on a boundary state,

$$\psi_p \sim f^+(p) e^{-\Phi(p)}.$$

This is a representation in momentum space. It is valid in a subspace of the Fock space consisting of excitations on top of the boundary state.

Again it has to be contrasted with the traditional "bosonization" - representation of Fermi-field in terms of a Bose-field with respect to the Fermi-vacuum

$$\psi(x) \sim e^{\varphi(x)}.$$

This is a representation in coordinate space. It is valid in a subspace of the Fock space consisting of excitations on top of the Fermi-vacuum.

We emphasize that although the matrix elements (5) and (14) obey similar MKP equations they give essentially different solutions of those equations. For example, (14) gives only solutions satisfying analyticity requirements in odd flows. The latter condition excludes soliton solutions.

Finally, we emphasize that the Schur decomposition (61) of the tau-functions similar to (14) (an essential element of the proof) has appeared in [7] in the problem of counting of Hurwitz numbers and in [6] in constructions of tau-functions of hypergeometric type. Similar objects arise in supersymmetric gauge theories [7, 15] and also in matrix quantum mechanics.

V. ACKNOWLEDGEMENT

We have benefited from discussions with I. Krichever, A. Orlov, I. Gruzberg, J. Harnad, T. Takebe, A. Zabrodin. We thank A. Zabrodin and J. Harnard who pointed out the paper [6] to us and N. Nekrasov for the information about [7]. P.W. and E.B. were supported by the NSF MRSEC Program under DMR-0213745 and NSF DMR-0220198. E.B. was also supported by BSF 2004128 P.W. acknowledges support by the Humboldt foundation and is grateful to Takashi Takebe for his kind hospitality in Ochanomizu University. The work of AGA was supported by the NSF under the grant DMR-0348358.

APPENDIX A: REDUCTION FROM THE PARTICLE-HOLE SYMMETRY

Let us treat the tau-function as a function of space-time (18). At first glance, the equation (23) is not closed. It is written on two functions τ_m and τ_{m+1} . In fact, both functions are determined by the initial state $\langle g |$ and are not independent. The relation between these functions determined by formulas (61,62) makes (23) a closed equation.

However, in the case of generic g the relations between τ_m and τ_{m+1} are difficult to formulate. The situation is simplified when the initial state $\langle g |$ has some special form, or is restricted by symmetries. One may think about these cases as *reductions* of MKP hierarchy.

The multi-phase solution (61) is the simplest (finite-dimensional) reduction. In this case the tau-function, treated as a function of the variable $z = e^{-ix}$ is a polynomial in z^{-1} .

A more interesting reduction having direct physical applications occurs when the parameter a + m takes a "symmetric" value such that a + m = -(a + m + 1). We set⁶

$$a = -1/2, \quad m = 0.$$

In addition we require the state $\langle g |$ to be invariant under a charge conjugation

$$e^{2ix_0(p-q)}\langle g|p,q\rangle = -\overline{\langle g|-q,-p\rangle},\tag{A1}$$

where x_0 is some real constant. This condition reflects a particle-hole symmetry: $\psi_q^{\dagger} \rightarrow \psi_{-q}, \psi_p \rightarrow \psi_{-p}^{\dagger}$.

Under these conditions the tau-function treated as a function of real space-time coordinates (18) has a reflection property

$$\tau_0(x) = \overline{\tau_1(-x+2x_0)},\tag{A2}$$

where $\overline{\tau}$ stands for complex conjugation of the function. This property makes the equation (23) closed with the second complex function τ_1 being completely determined by τ_0 .

The analytical meaning of the condition (A2) becomes clear if one considers an extension of the tau-function to the complex plane z from the unit circle $z = e^{-i(x-x_0)}$. Then the reflection condition means that the coefficients of the formal series (61) for τ_0 and τ_1 are complex conjugated (we use a notation $\overline{f}(z) = \overline{f(\overline{z})}$)

$$\tau_0(z) = \overline{\tau_1}(z). \tag{A3}$$

This follows from the property

$$f_1^+(p)f_1^-(q) = f_0^+(-q)f_0^-(-p)$$

held at a = -1/2.

The holomorphic tau-function $\tau(z) = \tau_0(z)$ is defined in the exterior of the unit disk, while its Schwarz reflection $\tau(\bar{z}) = \bar{\tau}_1(1/\bar{z})$ s defined in the interior of the disk. The reflection condition (A3) means that the holomorphic tau-function, and its Schwarz reflection, can be glued along the unit circle forming a function analytical in some neighborhood of the circle.

Particle-hole symmetry (A1) occurs when A_{pq} is a Toeplitz matrix, i.e., it depends on the difference p-q. In this case the state $\langle g |$ is given by

$$g = e^{i \int A(x)\rho(x)dx}, \quad A(x-x_0) = -A(-x+x_0),$$
(A4)

where A(x) - the symbol of the Toeplitz matrix i.e., a Fourier transform of A_{pq} . It is an odd real function.

E. Date, M. Jimbo, and T. Miwa. Method for generating discrete Soliton equations. III. Journal of the Physical Society of Japan, 52 (2):388–393, 1983.

Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. In M. Stone, editor, *Bosonization*, Singapore, 1994. World Scientific. RIMS-394.

^[2] M. Sato. Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds. *RIMS Kokyuroku*, 439:30–40, 1981. Sato, Mikio and Sato, Yasuko: Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold. Nonlinear partial differential equations in applied science (Tokyo, 1982), 259–271, North-Holland Math. Stud., 81, North-Holland, Amsterdam, 1983.

 ^[3] E. Date T. Miwa, M. Jimbo. Solitons: Differential Equations, Symmetries and Infinite Dimensional Algebras, volume 135 of Cambridge Tracts in Math. Cambridge university press, London, Great Britain, 1999. Michio Jimbo and Tetsuji Miwa. Solitons and infinite dimensional lie algebras. Publ. Res. Inst. Math. Sci. Kyoto, 19:943, 1983.
 E. Date, M. Jimbo, and T. Miwa. Method for generating discrete Soliton equations. I. Journal of the Physical Society of

E. Date, M. Jimbo, and I. Miwa. Method for generating discrete Soliton equations. I. Journal of the Physical Society of Japan, 51 (12):4116–4124, 1982.

E. Date, M. Jimbo, and T. Miwa. Method for generating discrete Soliton equations. II. Journal of the Physical Society of Japan, 51 (12):4125–4131, 1982.

⁶ This case is known as a "unitary limit" in problems related to Fermi-edge singularities.

- [4] A. R. Its, A. G. Izergin, V. E. Korepin, and N. A. Slavnov. Differential Equations for Quantum Correlation Functions. International Journal of Modern Physics B, 4:1003–1037, 1990.
- [5] M. Jimbo, T. Miwa, Y. Môri, and M. Sato. Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent. *Physica D Nonlinear Phenomena*, 1:80–158, April 1980. M. Sato, T. Miwa, and M. Jimbo, Proc. Jpn. Acad. 53A, 147, 153, 183 (1977); Publ. Res. Inst. Math. Sci. 14, 223 (1978); 15, 201, 577, 871 (1979).
- [6] A Yu Orlov. Hypergeometric tau functions $\tau(\mathbf{t}, \mathbf{t}, \mathbf{t}^*)$ as ∞ -Soliton tau function in t variables. arXiv.org:nlin/0305001.
- [7] A. Okounkov. Toda equations for Hurwitz numbers. arXiv:math/0004128. A. Okounkov and R. Pandharipande. The equivariant Gromov-Witten theory on P¹. arXiv:math.AG/0207233. N. A. Nekrasov and A. Okounkov. Seiberg-Witten theory and random partitions. arXiv:hep-th/0306238.
- [8] I. Affleck and A. W. W. Ludwig. The Fermi edge singularity and boundary condition changing operators. Journal of Physics A Mathematical General, 27:5375–5392, August 1994.
- [9] K.Ueno and K.Takasaki, Toda lattice hierarchy, Adv. Studies in Pure Math. 4 (1984) 1-95.
- [10] P. Nozires and C. T. deDominicis, Phys. Rev. 178, 1097 (1969). G. D. Mahan, Phys. Rev. 163, 612 (1967).
 P. W. Anderson, Phys. Rev. Lett. 18, 1049 (1967).
- [11] I.G.Macdonald, Symmetric Functions and Hall Polynomials, Clarendon Press, Oxford, 1995
- [12] R. Hirota. Exact solution of the kortewegde vries equation for multiple collisions of solitons. Phys. Rev. Lett., 27:1192, 1971.
- [13] J. Satsuma and Y. Ishimori. Periodic-wave and rational Soliton solutions of the Benjamin-Ono equation. J. Phys. Soc. Japan, 46:681–687, 1979.
- [14] E. Bettelheim, A. G. Abanov and P. Wiegmann, Quantum Schock Waves the case for non-linear effects in dynamics of electronic liquid. arXiv:cond-mat/0606778; Orthogonality catastrophe and shock waves in a non-equilibrium Fermi gas. arXiv:cond-mat/0607453.
- [15] Nekrasov N.A.: Seiberg-Witten Prepotential from Instanton Counting, hep-th/0206161 v1; Losev A, S., Marshakov A. and Nekrasov N.A.: Small Instantons, Little Strings and Free Fermions, hep-th/0302191 v3.