# Combinatorial Bethe ansatz and ultradiscrete Riemann theta function with rational characteristics <sup>1</sup>

## Atsuo Kuniba and Reiho Sakamoto

ABSTRACT. The  $U_q(\hat{sl}_2)$  vertex model at q = 0 with periodic boundary condition is an integrable cellular automaton in one-dimension. By the combinatorial Bethe ansatz, the initial value problem is solved for arbitrary states in terms of an ultradiscrete analogue of the Riemann theta function with rational characteristics.

### 1. INTRODUCTION

Solvable vertex models [2] associated with a quantum affine algebra  $U_q$  yield completely integrable one-dimensional cellular automata at q = 0 [9, 8, 6]. The simplest case of  $U_q(\hat{sl}_2)$  with periodic boundary condition is known as the periodic box-ball system [17, 22]. Here is an example of the time evolution pattern:

z = 0:	1112222211112222222211111111111111112222
= 1:	11111111222211111111222222222211111111
z = 2:	22221111111122221111111111111222222222
= 3:	1111222222221111222211111111111111111
= 4:	22221111111122221111222222222211111111
= 5:	111122221111111122221111111112222222222
c = 6:	222211112222111111122221111111111111112222
t = 7:	111122221111222222221111222221111111111
t = 8:	111111112222111111122221111122222222222
t = 9:	222211111111222211111111222211111111112222

The dynamics is described as a motion of balls (letter 2) hopping to the right along the periodical array of boxes. The fusion transfer matrices  $T_1, T_2, \ldots$  in the q = 0vertex model give rise to a commuting family of deterministic time evolutions. The above example is the evolution under  $T_9$ .

In [17], the initial value problem of the periodic box-ball system is solved by an inverse scattering method. It synthesizes the combinatorial versions of the Bethe ansatz [3] at q = 1 [13] and q = 0 [14]. The action-angle variables are constructed from the rigged configurations (q = 1) by imposing a certain equivalence relations specified by the string center equation (q = 0). The time evolution  $T_l^t(p)$  of any state p is obtained by an explicit algorithm whose computational steps are independent of t.

The Bethe ansatz approach [17] captures several characteristic features in the quasi-periodic solutions of soliton equations [4, 5, 11]. It generalizes the connection between the original box-ball system on infinite lattice [20] and soliton solutions via the ultradiscretization [21] to a periodic setting. For instance, the nonlinear dynamics of the periodic box-ball system becomes a straight motion of the Bethe roots (angle variable) which live in an ultradiscrete analogue (4.1) of the Jacobi variety.

 $<sup>^1</sup>$  Mathematics Subject Classification (2000). 81R50, 82B23, 37B15, 37K15, 05E15 Keywords: crystal basis, periodic box-ball system, Bethe ansatz, Riemann theta function

In this paper we express the algorithmic solution of the initial value problem [17] by an explicit formula involving the ultradiscretization of the Riemann theta function with rational characteristics ( $\mathbf{z} \in \mathbb{R}^{g}$ ,  $\mathbf{a} \in (\mathbb{Q}/\mathbb{Z})^{g}$ ):

(1.1) 
$$\Theta_{\mathbf{a}}(\mathbf{z}) = \lim_{\epsilon \to +0} \epsilon \log \vartheta_{\mathbf{a}}(\mathbf{z}) \\ = -\min_{\mathbf{n} \in \mathbb{Z}^g} \{ {}^t(\mathbf{n} + \mathbf{a}) \Omega(\mathbf{n} + \mathbf{a})/2 + {}^t(\mathbf{n} + \mathbf{a}) \mathbf{z} \}.$$

Here  $\vartheta_{\mathbf{a}}(\mathbf{z})$  is the Riemann theta function [19] with a pure imaginary period matrix:

(1.2) 
$$\vartheta_{\mathbf{a}}(\mathbf{z}) = \sum_{\mathbf{n} \in \mathbb{Z}^g} \exp\left(-\frac{t(\mathbf{n} + \mathbf{a})\Omega(\mathbf{n} + \mathbf{a})/2 + t(\mathbf{n} + \mathbf{a})\mathbf{z}}{\epsilon}\right).$$

Here  $\Omega$  is a symmetric positive definite integer matrix (5.1). The function  $\Theta_{\mathbf{a}}(\mathbf{z})$  enjoys the quasi-periodicity analogous to the Riemann theta function:

(1.3) 
$$\Theta_{\mathbf{a}}(\mathbf{z} + \mathbf{v}) = \Theta_{\mathbf{a}}(\mathbf{z}) + {}^{t}\mathbf{v}\Omega^{-1}(\mathbf{z} + \mathbf{v}/2) \text{ for any } \mathbf{v} \in \Omega\mathbb{Z}^{g}.$$

Our main Theorem 5.1 covers *all* the states in the periodic box-ball system, which generalizes the earlier result [15] on those states in which the amplitudes of solitons were assumed to be distinct. The theorem is derived from the recently obtained piecewise linear formula [16] for the Kerov-Kirillov-Reshetikhin (KKR) bijection [13] from rigged configurations to highest paths. It involves the ultradiscretization of the tau function for the KP hierarchy [10]. A state p of the periodic box-ball system is effectively treated as the infinite system  $p \otimes p \otimes \cdots$ , and the ultradiscrete tau function for such states turns out to be expressible in terms of  $\Theta_{\mathbf{a}}(\mathbf{z})$  as in (5.5).

In section 2 we define the periodic box-ball system. In section 3 various data in the combinatorial Bethe ansatz at q = 1 [13] and q = 0 [14] are introduced. In section 4 we recall the solution of the initial value problem by the inverse scattering transform. In section 5 we present our main Theorem 5.1 in this paper.

#### 2. Periodic box-ball system

Let us recall the periodic box-ball system without getting much into the crystal base theory [12]. For a positive integer l, let  $B_l = \{(x_1, x_2) \in (\mathbb{Z}_{\geq 0})^2 \mid x_1 + x_2 = l\}$  and set  $u_l = (l, 0) \in B_l$ . The element  $(x_1, x_2)$  will also be denoted by the sequence

 $\overbrace{1...12...2}^{x_1}$ , which is the semistandard Young tableau (without a frame) containing l

the letter  $i x_i$  times. In this notation  $u_l = \overbrace{1 \dots 1}^{l}$ , and the two elements (1,0) and (0,1) in  $B_1$  are simply written as 1 and 2, respectively. In the following, the symbol  $\otimes$  meaning the tensor product of crystals can just be understood as a product of sets. Define the map  $R : B_l \otimes B_1 \to B_1 \otimes B_l$  by

$$(x_1, x_2) \otimes 1 \mapsto \begin{cases} 1 \otimes (l, 0) & \text{if } (x_1, x_2) = (l, 0) \\ 2 \otimes (x_1 + 1, x_2 - 1) & \text{otherwise,} \end{cases}$$

$$(x_1, x_2) \otimes 2 \mapsto \begin{cases} 2 \otimes (0, l) & \text{if } (x_1, x_2) = (0, l) \\ 1 \otimes (x_1 - 1, x_2 + 1) & \text{otherwise.} \end{cases}$$

*R* is a bijection and called the combinatorial *R*. We write the relation  $R(u \otimes b) = b' \otimes u'$  simply as  $u \otimes b \simeq b' \otimes u'$ , and similarly for any consequent relation of the form  $a \otimes u \otimes b \otimes c \simeq a \otimes b' \otimes u' \otimes c$ .

A state of the periodic box-ball system is an array of 1 and 2, which is regarded as an element  $b_1 \otimes \cdots \otimes b_L \in B_1^{\otimes L}$  with L being the system size. Let the number of  $2 \in B_1$  appearing in  $b_1 \otimes \cdots \otimes b_L$  be N. We assume  $L \ge 2N$ . The other case can be reduced to this case by the exchange  $1 \leftrightarrow 2$  as explained in §3.3 of [17]. Let  $\mathcal{P}$  be the set of such states. Then the time evolution  $T_l : \mathcal{P} \to \mathcal{P}$  is defined by

(2.1) 
$$u_l \otimes p \simeq p^* \otimes v_l, \quad v_l \otimes p \simeq T_l(p) \otimes v_l.$$

In the first relation, one applies the combinatorial  $R \ L$  times to carry  $u_l$  through  $p \in \mathcal{P}$  to the right. It determines  $v_l \in B_l$  and  $p^* \in \mathcal{P}$  uniquely.  $(p^* \text{ will play no role}$  in the sequel.) Then the second relation using the so obtained  $v_l$  specifies  $T_l(p)$ , where the appearance of the same  $v_l$  in the right hand side is a non-trivial claim ([17] (2.10)).  $v_l$  is dependent on p as opposed to  $u_l$ .

The combinatorial R is the identity map on  $B_1 \otimes B_1$ , therefore  $T_1$  is just the cyclic shift  $T_1(b_1 \otimes \cdots \otimes b_L) = b_L \otimes b_1 \otimes \cdots \otimes b_{L-1}$ . The commutativity  $T_l T_k = T_k T_l$  holds for any k, l ([17] Theorem 2.2).

**Example 2.1.** We depict the relation  $R(u \otimes b) = b' \otimes u'$  by a vertex diagram:

$$u \xrightarrow{b} u'$$

Then for  $p = 1121221 \in B_1^{\otimes 7}$  (symbol  $\otimes$  omitted), one has  $v_2 = (1, 1) = 12 \in B_2$  from (2.1). The time evolution  $T_2(1121221)$  is obtained by the composition of the vertex diagrams

$$1 \\ 12 \\ + \\ 11 \\ + \\ 11 \\ + \\ 11 \\ + \\ 12 \\ + \\ 12 \\ + \\ 11 \\ + \\ 12 \\ + \\ 11 \\ + \\ 12 \\ +$$

leading to  $T_2(1121221) = 2112112$ .

#### 3. Combinatorial Bethe Ansatz data

Let  $\mathcal{P}_+$  be the subset of  $\mathcal{P}$  consisting of the  $sl_2$ -highest states. Namely,  $\mathcal{P}_+ = \{p \in \mathcal{P} \mid \tilde{e}_1 p = 0\}$ , where  $\tilde{e}_1$  is a Kashiwara operator [12]. In practice,  $p = b_1 \otimes \cdots \otimes b_L$  is highest if and only if  $b_1 \otimes \cdots \otimes b_k$  has nonnegative weights for any  $1 \leq k \leq L$ .

According to the combinatorial Bethe ansatz at q = 1 [13], there is one to one correspondence between  $\mathcal{P}_+$  and the set of the objects called the *rigged configurations*. We let  $\phi(p_+) = (\mu, J)$  denote the Kerov-Kirillov-Reshetikhin (KKR) bijection. Here  $p_+$  is highest state and the right hand side is a rigged configuration. The data  $\mu$ , called a configurations, is just a Young diagram for a partition of N, for which  $N \leq L/2$  is assumed.

In what follows, we let  $\mathbb{I} = \{i_1 < i_2 < \cdots < i_g\}$  be the set of distinct lengths of rows of  $\mu$ , and the multiplicity of the length  $i \in \mathbb{I}$  rows by  $m_i \geq 1$ . In terms of the non-increasing array of parts we have

(3.1) 
$$\mu = (i_g^{m_{i_g}} \dots i_1^{m_{i_1}}).$$

Obviously, a configuration  $\mu$  is equivalent with the data I and  $m_{i_1}, \ldots, m_{i_n}$ .

To each row of  $\mu$ , there is assigned an nonnegative integer called a rigging, which is represented by the symbol J in  $(\mu, J)$ . See Example 5.2. Let  $J_{i,1}, \ldots, J_{i,m_i}$  be the ones assigned to the  $m_i$  rows of length i in  $\mu$  from the bottom to the top. The data  $(\mu, J)$  with  $J = \{(J_{i,\alpha}) \mid i \in \mathbb{I}, 1 \le \alpha \le m_i\}$  will be called a rigged configuration if  $0 \le J_{i,1} \le \cdots \le J_{i,m_i} \le p_i$  holds for each  $i \in \mathbb{I}$ . Here  $p_i$  is defined by

(3.2) 
$$p_i = L - 2\sum_{j \in \mathbb{I}} \min(i, j) m_j,$$

which is nonnegative if  $\mu$  is a configuration.

We introduce the index set

(3.3) 
$$\tilde{\mathbb{I}} = \{(i,\alpha) \mid i \in \mathbb{I}, 1 \le \alpha \le m_i\}, \quad \gamma = |\tilde{\mathbb{I}}| = m_{i_1} + \dots + m_{i_i}$$

labeling the rows of  $\mu$ . The following vectors and matrix whose components are labeled with  $\tilde{\mathbb{I}}$  will be used:

$$(3.4) \quad \mathbf{h}_j = (\min(i,j))_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathbb{Z}^{\gamma} \quad (j \in \mathbb{I}),$$

(3.5) 
$$\mathbf{\hat{J}} = (J_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathbb{Z}^{\gamma},$$

$$(3.6) \qquad A = (A_{i\alpha,j\beta})_{(i,\alpha),(j,\beta)\in\tilde{\mathbb{I}}}, \quad A_{i\alpha,j\beta} = \delta_{i,j}\delta_{\alpha,\beta}(p_i + m_i) + 2\min(i,j) - \delta_{i,j}$$

We also introduce the vectors and matrices whose indices range over the previously introduced set  $\mathbb{I} = \{i_1, \ldots, i_g\}$  as

(3.7) 
$$\mathbf{h}_j = (\min(i,j))_{i \in \mathbb{I}} \in \mathbb{Z}^g,$$

(3.8) 
$$\mathbf{p} = (p_i)_{i \in \mathbb{I}} = L\mathbf{h}_1 - 2\sum_{j \in \mathbb{I}} m_j \mathbf{h}_j \in \mathbb{Z}^g,$$

(3.9) 
$$\mathbf{J} = (J_{i,1} + \dots + J_{i,m_i})_{i \in \mathbb{I}} \in \mathbb{Z}^g,$$

(3.10) 
$$F = (F_{i,j})_{i,j \in \mathbb{I}}, \quad F_{i,j} = \sum_{\beta=1}^{j} A_{i\alpha,j\beta} = \delta_{ij} p_i + 2\min(i,j)m_j,$$

$$(3.11) M = \operatorname{diag}(m_i)_{i \in \mathbb{I}},$$

The matrices A, F were introduced in the study of q = 0 Bethe equation [14]. In particular, A is the coefficient matrix of the linearized Bethe equation (string center equation), which is known to be positive definite. The result of the sum in (3.10) is independent of  $\alpha$ .

### 4. ACTION-ANGLE VARIABLE AND INVERSE SCATTERING TRANSFORM

Any state  $p \in \mathcal{P}$  can be made highest by applying the cyclic shift appropriately. Namely, one can express p as  $p = T_1^d(p_+)$  with some integer d and highest state  $p_+ \in \mathcal{P}_+$ . Although the choice of d and  $p_+$  is not unique, the configuration of  $\phi(p_+)$  is unique. The set of states is decomposed according to the configuration of  $p_+$  as  $\mathcal{P} = \bigsqcup_{\mu} \mathcal{P}(\mu)$ . We call  $\mu$  the action variable. It is a conserved quantity ([17] Lemma C.3, Corollary 3.5), therefore each subset  $\mathcal{P}(\mu)$  is invariant under any time evolution  $T_l$ . The action variable  $\mu$  is equivalent with the data  $\mathbb{I} = \{i_1, \ldots, i_g\}$  and  $\{m_i \mid i \in \mathbb{I}\}$  via (3.1). In the periodic box-ball system,  $\mathbb{I}$  is the list of amplitudes of solitons and  $m_i$  is the number of solitons with amplitude i.

Given an action variable  $\mu$ , the set of angle variable is given by

(4.1) 
$$\mathcal{J}(\mu) = (\mathcal{I}_{m_{i_1}} \times \dots \times \mathcal{I}_{m_{i_g}})/\Gamma, \quad \Gamma = A\mathbb{Z}^{\gamma}$$

where  $\mathcal{I}_n = (\mathbb{Z}^n - \Delta_n)/\mathfrak{S}_n$  is the *n*-dimensional lattice without the "diagonal points"  $\Delta_n = \{(z_1, \ldots, z_n) \in \mathbb{Z}^n \mid z_\alpha = z_\beta \text{ for some } 1 \leq \alpha \neq \beta \leq n\}$  divided by the action of the symmetric group  $\mathfrak{S}_n$ . Plainly, it is the set of array of *n* distinct

integers whose ordering does not matter. In the sequel, elements in  $\mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$ and its image in  $\mathcal{J}(\mu)$  will be denoted by the same symbol. The set  $\mathcal{J}(\mu)$  is finite and its cardinality is given by [14, 17]

(4.2) 
$$|\mathcal{J}(\mu)| = (\det F) \prod_{i \in \mathbb{I}} \frac{1}{m_i} \binom{p_i + m_i - 1}{m_i - 1} = \frac{L}{p_{i_g}} \prod_{i \in \mathbb{I}} \binom{p_i + m_i - 1}{m_i}.$$

This is a positive integer if  $\mu$  is a configuration. (When  $|\mu| = L/2$ , we have  $p_{i_g} = 0$ . In this case, the factor  $\frac{L}{p_{i_g}} {p_{i_g} + m_{i_g} - 1 \choose m_{i_g}}$  is to be understood as  $\frac{L}{m_{i_g}}$ .) The formula (4.2) tells the number of states containing  $m_i$  solitons with amplitude *i*. The completeness of soliton states is known in the sense that  $\sum_{\mu} |\mathcal{J}(\mu)| = {L \choose N}$  for  $0 \leq N \leq L/2$ , where the sum extends over those  $m_i$  such that  $\sum_i im_i = N$  ([14], Theorems 3.5 and 4.9). The states in section 1 have the length L = 45 and  $\mu$  depicted in Example 5.2. For this  $\mu$  and L, the formula (4.2) tells  $|\mathcal{J}(\mu)| = 316350$ .

Let us specify the angle variable for a given state  $p \in \mathcal{P}(\mu)$ . Suppose p is expressed as  $p = T_1^d(p_+)$  in terms of a highest state  $p_+$  and the cyclic shift  $T_1$ . Let the image of the KKR bijection be  $\phi(p_+) = (\mu, J)$ . From the configuration  $\mu$  we know  $\mathbb{I} = \{i_1 < i_2 < \cdots < i_g\}$  and  $m_{i_1}, \ldots, m_{i_g}$  by (3.1). Then the angle variable is found by applying the definitions (3.4), (3.5), (3.6) to  $J = (J_{i,\alpha})_{(i,\alpha) \in \tilde{\mathbb{I}}}$  as

(4.3) 
$$\begin{aligned} \Phi: \ \mathcal{P}(\mu) &\longrightarrow \mathbb{Z} \times \mathcal{P}_+ \longrightarrow \mathcal{J}(\mu) \\ p &\longmapsto (d, p_+) \longmapsto (\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1) / \Gamma. \end{aligned}$$

Due to the shift  $+\alpha - 1$  in (3.5) and the rigging condition  $0 \leq J_{i,1} \leq \cdots \leq J_{i,m_i} \leq p_i$ , it follows that the components of  $\tilde{\mathbf{J}}$  hence  $\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1$  within each block labeled by  $i \in \mathbb{I}$ are distinct. In (4.3) we regard the  $\tilde{\mathbf{J}} + d\tilde{\mathbf{h}}_1$  as a representative of an element in  $\mathcal{J}(\mu)$ . The non-uniqueness of  $d, p_+$  is cancelled by mod  $\Gamma$  making  $\Phi$  well-defined ([17] Proposition 3.7).

**Theorem 4.1** ([17] Theorem 3.12).  $\Phi$  is a bijection and the following diagram is commutative.

$$egin{array}{cccc} \mathcal{P}(\mu) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(\mu) \ & & & & \downarrow T_l \ & & & \downarrow T_l \ \mathcal{P}(\mu) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(\mu) \end{array}$$

Here the time evolution on  $\mathcal{J}(\mu)$  is defined by  $T_l(\tilde{\mathbf{J}}) = \tilde{\mathbf{J}} + \tilde{\mathbf{h}}_l$ .

The composition  $\Phi^{-1} \circ T_l \circ \Phi$  achieves the solution of the initial value problem by the inverse scattering transform. Namely,  $\Phi^{\pm 1}$  is the direct/inverse scattering map which transforms the dynamics  $T_l$  of the periodic box-ball system into the straight motion on  $\mathcal{J}(\mu)$  with the velocity  $\tilde{\mathbf{h}}_l$ . In this sense,  $\mathcal{J}(\mu)$  serves as an ultradiscrete analogue of the Jacobi variety in quasi-periodic solutions to soliton equations [4, 5, 11].

By using Theorem 4.1, a closed formula for the fundamental period, i.e., the smallest positive integer such that  $T_l^{\mathcal{N}}(p) = p$  for any l and p has been obtained in Theorem 4.9 of [17], which supplements a recursive description of the  $l = \infty$  case [22]. The states in section 1 have the fundamental period 3515. See also [18] for a generalization to  $A_n^{(1)}$  based on the Bethe eigenvalue at q = 0.

We note that the time evolution of the angle variable  $\tilde{\mathbf{J}}$  (3.5) leads to the rule

(4.4) 
$$T_l(\mathbf{J}) = \mathbf{J} + M\mathbf{h}$$

for the "bundled" angle variable  $\mathbf{J}$  (3.9), where M is defined by (3.11).

5. EXPLICIT FORMULA FOR THE SOLUTION OF THE INITIAL VALUE PROBLEM

Fix the action variable  $\mu$  (3.1) and set

(5.1) 
$$\Omega = MF = (\Omega_{i,j})_{i,j \in \mathbb{I}}, \quad \Omega_{i,j} = \delta_{ij}p_im_i + 2\min(i,j)m_im_j$$

where F is defined in (3.11).  $\Omega$  is a positive definite symmetric integer matrix which will play the role of the period matrix. Note that

(5.2) 
$$\Omega \mathbf{h}_1 = LM \mathbf{h}_1$$

In addition to the ultradiscrete Riemann theta function  $\Theta_{\mathbf{a}}(\mathbf{z})$  in (1.1), we need  $\chi(\mathbf{s}; \tilde{\mathbf{I}})$ . It is a function of  $\mathbf{s} = (s_i)_{i \in \mathbb{I}} \in \mathbb{Z}^g$  as well as  $\tilde{\mathbf{I}} = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$  that takes rational values:

(5.3) 
$$\chi(\mathbf{s}; \tilde{\mathbf{I}}) = \chi(\mathbf{s}'; \tilde{\mathbf{I}}) \text{ if } \mathbf{s} \equiv \mathbf{s}' \mod M\mathbb{Z}^g,$$

(5.4) 
$$\chi(\mathbf{s}; \tilde{\mathbf{I}}) = \sum_{i \in \mathbb{I}} \frac{1}{m_i} \sum_{1 \le \alpha \le s_i < \beta \le m_i} (I_{i,\beta} - I_{i,\alpha} - \frac{p_i}{2})$$
  
if  $0 \le s_i < m_i, \quad I_{i,1} \le \dots \le I_{i,m_i}$  for all  $i \in \mathbb{I}$ 

Here the second sum in (5.4) extends over the pairs  $(\alpha, \beta)$ . In particular,  $\chi(\mathbf{0}; \tilde{\mathbf{I}}) = 0$ . The condition  $I_{i,1} \leq \cdots \leq I_{i,m_i}$  is needed to cope with the  $\mathfrak{S}_{m_i}$  symmetry in  $\mathcal{I}_{m_i}$ .

Given  $\mathbf{I} \in \mathbb{Z}^g$  and  $\tilde{\mathbf{I}} = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{I}_{m_{i_1}} \times \cdots \times \mathcal{I}_{m_{i_g}}$ , we define the ultradiscrete tau function by

(5.5) 
$$\tau_r(k) = \max_{\mathbf{s} \in \mathbb{Z}^g / M \mathbb{Z}^g} \left\{ \Theta_{M^{-1}\mathbf{s}} \left( \mathbf{I} + M \left( r \mathbf{h}_{\infty} - k \mathbf{h}_1 - \frac{\mathbf{p}}{2} \right) \right) + \chi(\mathbf{s}; \tilde{\mathbf{I}}) \right\}$$

for  $r \in \{0, 1\}$ ,  $k \in \mathbb{Z}$ . Here  $\mathbf{h}_j$ ,  $\mathbf{p}$  and M are those in (3.7), (3.8) and (3.11).

Let us specify  $\tilde{\mathbf{I}}$  and  $\mathbf{I}$  in (5.5) from a given state  $p \in \mathcal{P}(\mu)$ . As for  $\tilde{\mathbf{I}}$ , it is taken to be the angle variable of  $p \in \mathcal{P}(\mu)$  according to (4.3), and  $\mathbf{I}$  is determined from it as follows:

(5.6) 
$$\mathbf{I} = \Phi(p) = (I_{i,\alpha} + \alpha - 1)_{(i,\alpha) \in \tilde{\mathbb{I}}} \in \mathcal{J}(\mu),$$

(5.7) 
$$\mathbf{I} = (I_{i,1} + \dots + I_{i,m_i})_{i \in \mathbb{I}} \in \mathbb{Z}^g / F \mathbb{Z}^g.$$

This is the same relation as (3.5) and (3.9). Note that the elements  $\mathbf{I}$  and  $\mathbf{I}$  are defined mod  $\Gamma = A\mathbb{Z}^{\gamma}$  and mod  $F\mathbb{Z}^{g}$ , respectively, which is consistent with the relation (3.10) between A and F. Now we state the main result of this paper.

**Theorem 5.1.** The state  $p \in \mathcal{P}(\mu)$  corresponding to the action variable  $\mu$  and the angle variable  $\tilde{\mathbf{I}}$  is expressed as  $p = (1 - x_1, x_1) \otimes \cdots \otimes (1 - x_L, x_L)$ , where  $x_k = 0, 1$  is given by

(5.8) 
$$x_k = \tau_0(k) - \tau_0(k-1) - \tau_1(k) + \tau_1(k-1).$$

The proof is similar to section 3 in [15]. Combined with Theorem 4.1, Theorem 5.1 leads to an explicit formula for the solution of the initial value problem. The state  $(\prod_l T_l^{c_l})(p)$  is obtained just by replacing the angle variable as

(5.9) 
$$\tilde{\mathbf{I}} \mapsto \tilde{\mathbf{I}} + \sum_{l} c_{l} \tilde{\mathbf{h}}_{l}$$

in (5.5). Under the change (5.9), the function  $\chi(\mathbf{s}; \tilde{\mathbf{I}})$  (5.4) is invariant since it only depends on the difference  $I_{i,\beta} - I_{i,\alpha}$ . On the other hand, the variable  $\mathbf{I}$  entering (5.5) changes as

(5.10) 
$$\mathbf{I} \mapsto \mathbf{I} + M(\sum_{l} c_{l} \mathbf{h}_{l})$$

according to the rule (4.4). Especially,  $T_1^L(\mathbf{I}) = \mathbf{I} + LM\mathbf{h}_1 = \mathbf{I} + \Omega\mathbf{h}_1$  by (5.2). Thus the tau function only exhibits the change due to the quasi-periodicity (1.3) leaving (5.8) invariant. This is consistent with the fact that the cyclic shift  $T_1^L$  is trivial and (5.8) is a period L function of k. We remark that our tau function (5.5) is always integer-valued.

From (3.7) we see  $\mathbf{h}_l = \mathbf{h}_{\infty}$  if  $l \geq \max \mathbb{I}(=i_g)$ . Therefore the effect (5.10) of the time evolution  $T_l$  is  $\mathbf{I} \mapsto \mathbf{I} + M\mathbf{h}_{\infty}$ , which is equivalent to setting  $r \mapsto r+1$ in (5.5). In other words, r serves as the time variable for such  $T_l$ , i.e.,  $T_l^t(p) = (1 - x_{1,t}, x_{1,t}) \otimes \cdots \otimes (1 - x_{L,t}, x_{L,t})$  with

(5.11) 
$$x_{k,t} = \tau_t(k) - \tau_t(k-1) - \tau_{t+1}(k) + \tau_{t+1}(k-1)$$

holds for  $l \geq \max \mathbb{I}$ .

**Example 5.2.** Let us consider the t = 0, 1 states in section 1:

which are elements in  $B_1^{\otimes 45}$ . We have  $p = T_1^{29}(p_+)$  with a highest state

which corresponds to the rigged configuration:



Thus  $\mathbb{I} = \{1, 4, 9\}$ ,  $M = \text{diag}(m_1, m_4, m_9) = \text{diag}(1, 2, 1)$ .  $I_{i,\alpha}$  in (5.6) reads  $(I_{1,1}, I_{4,1}, I_{4,2}, I_{9,1}) = (39, 31, 35, 29)$ . The data **p** (3.8), **I** (5.7) and  $\Omega$  (5.1) are given by

$$\mathbf{p} = \begin{pmatrix} p_1 \\ p_4 \\ p_9 \end{pmatrix} = \begin{pmatrix} 37 \\ 19 \\ 9 \end{pmatrix}, \quad \mathbf{I} = \begin{pmatrix} 39 \\ 66 \\ 29 \end{pmatrix}, \quad \Omega = \begin{pmatrix} 39 & 4 & 2 \\ 4 & 70 & 16 \\ 2 & 16 & 27 \end{pmatrix}.$$

The argument of the tau function (5.5) reads

(5.13) 
$$\mathbf{I} + M(r\mathbf{h}_{\infty} - k\mathbf{h}_{1} - \frac{\mathbf{p}}{2}) = \begin{pmatrix} r - k + 41/2\\ 8r - 2k + 47\\ 9r - k + 49/2 \end{pmatrix} =: \mathbf{z}_{r,k} .$$

The function  $\chi(\mathbf{s}; \tilde{\mathbb{I}})$  in (5.4) is non-vanishing at  $\mathbf{s} = (0, 1, 0)$  taking the value  $\frac{1}{m_4}(I_{4,2} - I_{4,1} - \frac{p_4}{2}) = -\frac{11}{4}$ . Thus the tau function is given by

$$\tau_r(k) = \max\{\Theta_{0,0,0}(\mathbf{z}_{r,k}), \ \Theta_{0,\frac{1}{2},0}(\mathbf{z}_{r,k}) - \frac{11}{4}\}$$

The time evolution pattern in section 1 is reproduced according to (5.11). Here is the plot of  $\tau_r(k)$  for  $0 \le r \le 2$  and  $0 \le k \le 45$ .



The boundary values are  $(\tau_0(0), \tau_1(0), \tau_2(0)) = (15, 28, 41)$  and  $(\tau_0(45), \tau_1(45), \tau_2(45)) = (13, 8, 3)$ . In the plot of  $\tau_t(k)$  or  $\tau_{t+1}(k)$ , one observes a change of the gradient at those k corresponding to the front or tail of solitons in  $T_9^t(p)$  (5.12).

The tau function (5.5) simplifies considerably if there are no solitons with the same amplitude, namely,  $\forall m_i = 1$ . We have  $\chi(\mathbf{s}; \tilde{\mathbf{I}}) = 0$ ,  $\tilde{\mathbb{I}} = \mathbb{I}$  and  $\Omega = F = (\delta_{ij}p_i + 2\min(i,j))_{i,j\in\mathbb{I}}$ . Let us write the ultradiscrete Riemann theta function with characteristics  $\mathbf{0}$  as  $\Theta(\mathbf{z}) = \Theta_{\mathbf{0}}(\mathbf{z})$ . Then  $\tau_r(k) = \Theta(\mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1 + r\mathbf{h}_{\infty})$ . Consequently (5.8) reduces to

$$\begin{aligned} x_k &= \Theta \left( \mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1 \right) - \Theta \left( \mathbf{I} - \frac{\mathbf{p}}{2} - (k-1)\mathbf{h}_1 \right) \\ &- \Theta \left( \mathbf{I} - \frac{\mathbf{p}}{2} - k\mathbf{h}_1 + \mathbf{h}_\infty \right) + \Theta \left( \mathbf{I} - \frac{\mathbf{p}}{2} - (k-1)\mathbf{h}_1 + \mathbf{h}_\infty \right), \end{aligned}$$

which was obtained in [15]. This has the same structure as the ultradiscretization of the solution of the periodic Toda lattice by Date and Tanaka [4].

**Acknowledgments** The authors thank Rei Inoue for a discussion. R.S. is a research fellow of the Japan Society for the Promotion of Science. He thanks Miki Wadati for continuous encouragements.

#### References

- Ablowitz, M. J. and Segur, H.: Solitons and the inverse scattering transform, SIAM Studies in Appl. Math. 4. Philadelphia Pa. (1981)
- [2] Baxter, R. J.: Exactly solved models in statistical mechanics, Academic Press, London (1982)
- Bethe, H. A.: Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, Z. Physik 71 205–231 (1931)
- [4] Date, E. and Tanaka, S.: Periodic multi-soliton solutions of Korteweg-de Vries equation and Toda lattice, Prog. Theoret. Phys. Suppl. 59 107–125 (1976)

- [5] Dubrovin, B. A., Matveev, V. B. and Novikov, S. P.: Nonlinear equations of Korteweg-de Vries type, finite-band linear operators and Abelian varieties Russian Math. Surveys **31** 59– 146 (1976)
- [6] Fukuda, K., Okado, M. and Yamada, Y.: Energy functions in box ball systems, Int. J. Mod. Phys. A 15 1379–1392 (2000)
- [7] Gardner, C. S., Greene, J. M., Kruskal, M. D. and Miura, R. M.: Method for solving the Korteweg-de Vries equation, Phys. Rev. Lett. 19 1095–1097 (1967)
- [8] Hatayama, G., Hikami, K., Inoue, R., Kuniba, A., Takagi T. and Tokihiro, T.: The A<sup>(1)</sup><sub>M</sub> automata related to crystals of symmetric tensors, J. Math. Phys. 42 274–308 (2001)
- [9] Hatayama, G., Kuniba, A. and Takagi, T.: Soliton cellular automata associated with crystal bases, Nucl. Phys. B577[PM] 619–645 (2000)
- [10] Jimbo, M. and Miwa, T.: Solitons and infinite dimensional Lie algebras, Publ. RIMS. Kyoto Univ. 19 943–1001 (1983)
- [11] Kac, M. and Moerbeke, P.: A complete solution of the periodic Toda problem, Proc. Nat. Acad. Sci. USA. 72 2879–2880 (1975)
- [12] Kashiwara, M.: On crystal bases of the q-analogue of universal enveloping algebras. Duke Math. J.  ${\bf 63}$  465–516 (1991)
- [13] Kirillov, A. N. and Reshetikhin, N. Yu.: The Bethe ansatz and the combinatorics of Young tableaux, J. Soviet Math. 41 925–955 (1988)
- [14] Kuniba, A. and Nakanishi, T.: The Bethe equation at q = 0, the Möbius inversion formula, and weight multiplicities: I. The sl(2) case, Prog. in Math. **191** 185–216 (2000)
- [15] Kuniba, A. and Sakamoto, R.: The Bethe ansatz in a periodic box-ball system and the ultradiscrete Riemann theta function, J. Stat. Mech. P09005 (2006)
- [16] Kuniba, A. , Sakamoto, R. and Yamada, Y.: Tau functions in combinatorial Bethe ansatz, preprint math.QA/0610505
- [17] Kuniba, A., Takagi, T. and Takenouchi, A.: Bethe ansatz and inverse scattering transform in a periodic box-ball system, Nucl. Phys. B [PM] 354–397 (2006)
- [18] Kuniba, A. and Takenouchi, A.: Bethe ansatz at q=0 and periodic box-ball systems, J. Phys. A: Math. Gen. **39** 2551–2562 (2006)
- [19] Mumford, D.: Tata Lectures on Theta II, Birkhäuser, Boston (1984)
- [20] Takahashi, D. and Satsuma, J.: A soliton cellular automaton, J. Phys. Soc. Jpn. 59 3514–3519 (1990)
- [21] Tokihiro, T., Takahashi, D., Matsukidaira and J. Satsuma, J.: From soliton equations to integrable cellular automata through a limiting procedure, Phys. Rev. Lett. **76** 3247–3250 (1996)
- [22] Yoshihara, D., Yura, F. and Tokihiro, T.: Fundamental cycle of a periodic box-ball system, J. Phys. A: Math. Gen. 36 99–121 (2003)

Atsuo Kuniba:

Institute of Physics, Graduate School of Arts and Sciences, University of Tokyo, Komaba, Tokyo 153-8902, Japan atsuo@gokutan.c.u-tokyo.ac.jp

Reiho Sakamoto:

Department of Physics, Graduate School of Science, University of Tokyo, Hongo, Tokyo 113-0033, Japan

reiho@monet.phys.s.u-tokyo.ac.jp