

Remarks on rational solutions for the Korteveg - de Vries hierarchy

Nikolai A. Kudryashov

Department of Applied Mathematics
Moscow Engineering and Physics Institute
(State University)
31 Kashirskoe Shosse, 115409, Moscow,
Russian Federation

Abstract

Differential equations for the special polynomials associated with the rational solutions of the second Painlevé hierarchy are introduced. It is shown rational solutions of the Korteveg - de Vries hierarchy can be found taking the Yablonskii - Vorob'ev polynomials into account. Special polynomials associated with rational solutions of the Korteveg - de Vries hierarchy are presented.

Keywords: The Korteveg - de Vries hierarchy, Special polynomials, the Yablonskii - Vorob'ev polynomials, Rational solutions,

PACS: 02.30.Hq - Ordinary differential equations

1 Introduction

In this paper our interest is in rational solutions of the Korteveg - de Vries hierarchy [1]

$$u_t + \frac{\partial}{\partial x} L_{N+1}[u] = 0 \quad (1.1)$$

Here the operator L_N is determined by the Lenard recursion formula [1–3]

$$\frac{d}{dz} L_{N+1}[u] = \left(\frac{d^3}{dz^3} + 4u \frac{d}{dz} + 2u_z \right) L_N[u], \quad L_0[u] = \frac{1}{2} \quad (1.2)$$

From (1.2) we have

$$L_1[u] = u, \quad (1.3)$$

$$L_2[u] = u_{zz} + 3u^2, \quad (1.4)$$

$$L_3[u] = u_{zzzz} + 10u u_{zz} + 5u_z^2 + 10u^3, \quad (1.5)$$

$$L_4[u] = u_{zzzzzz} + 14u u_{zzzz} + 28u_z u_{zzz} + 21u_{zz}^3 + 70u^2 u_{zz} + 70u u_z^2 + 35u^4 \quad (1.6)$$

Hierarchy (1.1) is integrable by the inverse scattering transform [4]. The rational solutions of the Korteveg - de Vries equation (hierarchy (1.1) at $N = 1$) are studied in [5–7].

The aim of this paper is to introduce the hierarchy for the special polynomials associated with rational solutions of the second Painlevé hierarchy (the Yablonskii - Vorob'ev polynomials [8–13]). These special polynomials can be used to find the rational solutions of the Korteveg - de Vries hierarchy. As additional result of this paper is special polynomials associated with rational solutions of the Korteveg - de Vries hierarchy.

2 Hierarchy for the Yablonskii - Vorob'ev polynomials

Suppose $u(x, t)$ in (1.1) takes the form

$$u = 2 \frac{\partial^2 \ln(\varphi(t)y(z))}{\partial x^2}, \quad z = \frac{x}{((2N+1)t)^{\frac{1}{2N+1}}} \quad (2.1)$$

where $\varphi(t)$ is arbitrary function of t and $y(z)$ is a new function of z . Then from equation (1.1) we have the hierarchy

$$\frac{d}{dz} \left(z \frac{d \ln y}{dz} \right) = \frac{1}{2} L_{N+1} \left[2 \frac{d^2 \ln y}{dz^2} \right] \quad (2.2)$$

It can be shown that special polynomials $Q_n^{(N)}(z)$ associated with the rational solutions of the second Painlevé hierarchy satisfy hierarchy (2.2).

Assuming $N = 1$, $N = 2$ and $N = 3$ in (2.2) we have differential equations for $y(z) \equiv Q_n^{(1)}(z)$ in (2.3), $y(z) \equiv Q_n^{(2)}(z)$ in (2.4) and $y(z) \equiv Q_n^{(3)}(z)$ in (2.5)

$$yy_{zzzz} - 4y_z y_{zzz} + 3y_{zz}^2 - y y_z - z y_{zz} y + z y_z^2 = 0 \quad (2.3)$$

Table 2.1: The Yablonskii - Vorob'ev polynomials $Q_n^{(1)}(z)$

$$\begin{aligned}
Q_0^{(1)}(z) &= 1, \\
Q_1^{(1)}(z) &= z, \\
Q_2^{(1)}(z) &= z^3 + 4, \\
Q_3^{(1)}(z) &= z^6 + 20z^3 - 80, \\
Q_4^{(1)}(z) &= (z^9 + 60z^6 + 11200)z, \\
Q_5^{(1)}(z) &= z^{15} + 140z^{12} + 2800z^9 + 78400z^6 - 313600z^3 - 6272000, \\
Q_6^{(1)}(z) &= z^{21} + 280z^{18} + 18480z^{15} + 627200z^{12} - 17248000z^9 + 1448832000z^6 \\
&\quad + 19317760000z^3 - 38635520000, \\
Q_7^{(1)}(z) &= (z^{27} + 504z^{24} + 75600z^{21} + 5174400z^{18} + 62092800z^{15} + 13039488000z^{12} - \\
&\quad - 828731904000z^9 - 49723914240000z^6 - 3093932441600000)z
\end{aligned}$$

$$\begin{aligned}
y^2 y_{zzzzzz} - 6 y y_z y_{zzzzz} + 5 y y_{zz} y_{zzzz} + 10 y_z^2 y_{zzzz} - 20 y_z y_{zz} y_{zzz} + \\
+ 10 y_{zz}^3 - y_z y^2 - z y_{zz} y^2 + z y_z^2 y = 0
\end{aligned} \tag{2.4}$$

$$\begin{aligned}
y^4 y_{zzzzzzzz} - 8 y^3 y_z y_{zzzzzzz} + 28 y^2 y_z^2 y_{zzzzzz} + 7 y^3 y_{zzzz}^2 - \\
- 56 y y_z^3 y_{zzzzz} + 112 y y_z^2 y_{zzz}^2 + 28 y^2 y_{zz}^2 y_{zzzz} + 28 y y_{zz}^4 + \\
+ 56 y_z^2 y_{zz}^3 + 56 y_z^4 y_{zzzz} - 112 y_z^3 y_{zz} y_{zzz} + 28 y y_z^2 y_{zz} y_{zzzz} - \\
- 56 y^2 y_z y_{zzz} y_{zzzz} - 112 y y_z y_{zz}^2 y_{zzz} - y_z y^4 - z y_{zz} y^4 + z y_z^2 y^3 = 0
\end{aligned} \tag{2.5}$$

Equation (2.3) for $Q_n^{(1)}(z)$ was found in [10] but equations (2.4) and (2.5) for special polynomials $Q_n^{(2)}(z)$ and $Q_n^{(3)}(z)$ are new. Hierarchy (2.2) for the special polynomials associated with rational solutions of the second Painlevé hierarchy is new as well and we need to study this hierarchy in future.

Denote by $w = \frac{d \ln y}{dz}$ then equation (2.2) is reduced to equation

$$\frac{d(zw)}{dz} = L_{N+1} \left[\frac{dw}{dz} \right] \tag{2.6}$$

Assuming $N = 1$, $N = 2$ and $N = 3$ from (2.6) we obtain the differential equations

$$w_{zzz} + 3w_z^2 - w - zw_z = 0, \tag{2.7}$$

Table 2.2: The Yablonskii - Vorob'ev polynomials $Q_{(n)}^{(2)}(z)$

$Q_0^{(2)}(z) = 1,$
$Q_1^{(2)}(z) = z,$
$Q_2^{(2)}(z) = z^3,$
$Q_3^{(2)}(z) = z(z^5 - 144),$
$Q_4^{(2)}(z) = z^{10} - 1008z^5 - 48384,$
$Q_5^{(2)}(z) = z^{15} - 4032z^{10} - 3048192z^5 + 146313216,$
$Q_6^{(2)}(z) = z(z^{20} - 12096z^{15} - 21337344z^{10} - 33798352896z^5 - 4866962817024),$
$Q_7^{(2)}(z) = z^3(z^{25} - 30240z^{20} - 55883520z^{15} - 1182942351360z^{10} + 701543488297107456)$

$$w_{zzzzz} + 10 w_z w_{zzz} + 5 w_{zz}^2 + 10 w_z^3 - w - z w_z = 0, \quad (2.8)$$

$$w_{zzzzzz} + 14 w_z w_{zzzz} + 28 w_{zz} w_{zzz} + 21 w_{zzz}^3 + 70 w_z^2 w_{zzz} + 70 w_z w_{zz}^2 + 35 w_z^4 - w - z w_z = 0 \quad (2.9)$$

Equation (2.7) for $\frac{d \ln(Q_n^{(1)}(z))}{dz}$ was studied in [10] but equations (2.8) and (2.9) for special polynomials $\frac{d \ln(Q_n^{(2)}(z))}{dz}$ and $\frac{d \ln(Q_n^{(3)}(z))}{dz}$ are new.

Multiplying (2.7) on $(2 w_{zz} - \frac{1}{2})$ we have the first integral of this equation in the form [10]

$$w_{zz}^2 - \frac{1}{2} w_{zz} + 2 (2 w_z^2 - w) \left(w_z - \frac{z}{4} \right) = C_1 \quad (2.10)$$

Taking $C_1 = \frac{n(n+1)}{4}$ in (2.10) we have solutions of (2.10) in the form of the Yablonskii - Vorob'ev polynomials. Using $w(z) = \frac{z^2}{8} + p(z)$ in (2.10) we obtain the equation in the form [10]

$$p_{zz}^2 + 4 (p_z)^3 + 2 z (p_z)^2 - 2 p p_z - \frac{1}{4} \left(n + \frac{1}{2} \right)^2 = 0 \quad (2.11)$$

Hierarchy (2.6) is integrable because this is obtained from the Korteveg - de Vries hierarchy. The study of equation (2.7) by the Painlevé test [14] yields the Fuchs indices: $j_1 = -1$, $j_2 = 1$ and $j = 6$. These indices correspond to

Table 2.3: The Yablonskii - Vorob'ev polynomials for $Q_n^{(3)}(z)$

$Q_0^{(3)}(z) = 1$
$Q_1^{(3)}(z) = z$
$Q_2^{(3)}(z) = z^3$
$Q_3^{(3)}(z) = z^6$
$Q_4^{(3)}(z) = z^3(z^7 + 14400)$
$Q_5^{(3)}(z) = z(z^{14} + 129600z^7 - 373248000)$
$Q_6^{(3)}(z) = z^{21} + 648000z^{14} - 24634368000z^7 - 35473489920000$
$Q_7^{(3)}(z) = z^{28} + 2376000z^{21} - 825251328000z^{14} - 30436254351360000z^7$ $+ 43828206265958400000$

the arbitrary constants z_0 , a_1 and a_6 in the Laurent series

$$\begin{aligned}
 w(z) = & \frac{2}{(z - z_0)} + a_1 + \frac{z_0}{6} (z - z_0) - \left(\frac{z_0^2}{360} + \frac{a_1}{30} \right) (z - z_0)^3 - \\
 & - \frac{z}{72} (z - z_0)^4 + a_6 (z - z_0)^5 - \left(\frac{z_0^2}{4320} + \frac{a_1}{360} \right) (z - z_0)^6 + \dots
 \end{aligned} \tag{2.12}$$

Special polynomials $Q_n^{(N)}(z)$ were found in works [11, 13]. It is very important to remark that these polynomials can be found taking into account the differential - difference hierarchy [15]

$$Q_{n+1}^{(N)} Q_{n-1}^{(N)} = z (Q_n^{(N)})^2 - 2 (Q_n^{(N)})^2 L_N \left[2 \frac{d^2}{dz^2} (\ln Q_n^{(N)}) \right], \quad n \geq 1 \tag{2.13}$$

Assuming $Q_0^{(N)}(z) = 1$ and $Q_0^{(N)}(z) = z$ from (2.13) we have special polynomials $Q_n^{(N)}(z)$ at $n \geq 1$. Some of few special polynomials $Q_n^{(1)}(z)$, $Q_n^{(2)}(z)$ and $Q_n^{(3)}(z)$ are given in tables (2.1), (2.2) and (2.3).

3 Rational solutions of the Korteveg - de Vries hierarchy

Using the special polynomials associated with the rational solutions of the second Painlevé hierarchy (the Yablonskii - Vorob'ev polynomials) we can

find the rational solutions of the Korteveg - de Vries hierarchy (1.1) by the formula

$$u(x, t) = 2 \frac{\partial^2 \ln Q_n^{(N)}(z)}{\partial x^2}, \quad z = \frac{x}{((2N+1)t)^{\frac{1}{2N+1}}} \quad (3.1)$$

Table 3.1: Polynomials $P_n^{(1)}(x, t)$

$$\begin{aligned}
P_1^{(1)}(x, t) &= x, \\
P_2^{(1)}(x, t) &= x^3 + 12t, \\
P_3^{(1)}(x, t) &= x^6 + 60x^3t - 720t^2, \\
P_4^{(1)}(x, t) &= (x^9 + 180x^6t + 302400t^3)x, \\
P_5^{(1)}(x, t) &= x^{15} + 420x^{12}t + 25200x^9t^2 + 2116800x^6t^3 - \\
&\quad -2540166000x^3t^4 - 1524096000t^5, \\
P_6^{(1)}(x, t) &= x^{21} + 840x^{18}t + 166320x^{15}t^2 + 16934400x^{12}t^3 - 1397088000x^9t^4 + \\
&\quad + 352066176000x^6t^5 + 14082647040000x^3t^6 - 84495882240000t^7, \\
P_7^{(1)}(x, t) &= x(x^{27} + 1512x^{24}t + 680400x^{21}t^2 + 139708800x^{18}t^3 + \\
&\quad + 5029516800x^{15}t^4 + 3168595584000x^{12}t^5 - 604145558016000x^9t^6 - \\
&\quad - 108746200442880000x^6t^7 - 60897872248012800000t^9)
\end{aligned}$$

The four rational solutions of the Korteveg - de Vries equation (hierarchy (1.1) at $N = 1$) take the form

$$u_1(x, t) = -\frac{2}{x^2}, \quad (3.2)$$

$$u_2(x, t) = 6 \frac{x(-x^3 + 24t)}{(x^3 + 12t)^2}, \quad (3.3)$$

$$u_3(x, t) = -12 \frac{x(x^9 + 43200t^3 + 5400x^3t^2)}{(-x^6 - 60x^3t + 720t^2)^2}, \quad (3.4)$$

$$\begin{aligned}
u_4(x, t) &= -\frac{20}{x^2(x^9 + 180x^6t + 302400t^3)^2} [x^{18} + 144x^{15}t - \\
&\quad - 2116800x^9t^3 + 22680x^{12}t^2 - 152409600x^6t^4 + 9144576000t^6]
\end{aligned} \quad (3.5)$$

The four rational solutions of the fifth-order Korteveg - de Vries equation (hierarchy (1.1) at $N = 2$) can be written in the form

$$u_1(x, t) = -\frac{2}{x^2}, \quad (3.6)$$

Table 3.2: Polynomials $P_n^{(2)}(x, t)$

$$\begin{aligned}
P_1^{(2)}(x, t) &= x, \\
P_2^{(2)}(x, t) &= x^3, \\
P_3^{(2)}(x, t) &= x(x^5 - 720t), \\
P_4^{(2)}(x, t) &= x^{10} - 5040x^5t - 1209600t^2, \\
P_5^{(2)}(x, t) &= x^{15} - 20160x^{10}t - 76204800x^5t^2 + 18289152000t^3, \\
P_6^{(2)}(x, t) &= x(x^{20} - 60480x^{15}t - 533433600x^{10}t^2 - 4224794112000x^5t^3 - \\
&\quad - 3041851760640000t^4), \\
P_7^{(2)}(x, t) &= x^3(x^{25} - 151200x^{20}t - 1397088000x^{15}t^2 - 147867793920000x^{10}t^3 + \\
&\quad + 2192323400928460800000t^5)
\end{aligned}$$

$$u_2(x, t) = -\frac{6}{x^2}, \quad (3.7)$$

$$u_3(x, t) = -\frac{12(x^{10} + 2160x^5t + 86400t^2)}{(-x^5 + 720t)^2x^2} \quad (3.8)$$

$$u_4(x, t) = -\frac{20x^3(x^{15} + 5040x^{10}t + 23587200x^5t^2 - 12192768000t^3)}{(x^{10} - 5040x^5t - 1209600t^2)^2} \quad (3.9)$$

The four rational solutions of the seventh-order Kortevæg - de Vries equation (hierarchy (1.1) at $N = 3$) can be given as the following

$$u_1(x, t) = -\frac{2}{x^2}, \quad (3.10)$$

$$u_2(x, t) = -\frac{6}{x^2}, \quad (3.11)$$

$$u_3(x, t) = -\frac{12}{x^2}, \quad (3.12)$$

$$u_4(x, t) = -\frac{20(x^{14} - 362880x^7t + 3048192000t^2)}{x^2(x^7 + 100800t)^2} \quad (3.13)$$

The rational solutions of the Kortevæg - de Vries hierarchy can be presented in the form of special polynomials taking into account the arbitrary function

Table 3.3: Polynomials $P_n^{(3)}(x, t)$

$P_1^{(3)}(x, t) = x,$
$P_2^{(3)}(x, t) = x^3,$
$P_3^{(3)}(x, t) = x^6,$
$P_4^{(3)}(x, t) = x^3 (x^7 + 100800 t),$
$P_5^{(3)}(x, t) = x (x^{14} + 907200 x^7 t - 18289152000 t^2),$
$P_6^{(3)}(x, t) = x^{21} + 4536000 x^{14} t - 1207084032000 x^7 t^2 - 12167407042560000 t^3$
$P_7^{(3)}(x, t) = x^{28} + 16632000 x^{21} t - 40437315072000 x^{14} t^2 -$ $-10439635242516480000 x^7 t^3 + 105231523244566118400000 t^4$

$\varphi(t)$ in (2.1). These polynomials are determined by means of formulae

$$P_n^{(N)}(x, t) = ((2N + 1)t)^{\binom{n^2+n}{4N+2}} Q_n^{(N)}(z), \quad z = \frac{x}{((2N + 1)t)^{\frac{1}{2N+1}}} \quad (3.14)$$

Using these polynomials we obtain rational solutions from (2.4) of the Korteweg - de Vries hierarchy (1.1) by formula

$$u_n^{(N)}(x, t) = 2 \frac{\partial^2 \ln P_n^{(N)}(x, t)}{\partial x^2} \quad (3.15)$$

Some of these polynomials $P_n^{(1)}(x, t)$, $P_n^{(2)}(x, t)$ and $P_n^{(3)}(x, t)$ are given in tables (3.1), (3.2) and (3.3).

4 Acknowledgments

This work was supported by the International Science and Technology Center under Project B 1213.

References

- [1] *P.D. Lax*, Integrals of nonlinear equations of evolution and solitary waves, Comm. Pure Appl. Math., v.21, (1968), 467 - 490

- [2] *N.A. Kudryashov*, The first Painleve and the second Painleve equations of higher order and some relations between them, Physics Letters A, v.224, (1997), 353 - 360
- [3] *N.A. Kudryashov*, Analytical theory of nonlinear differential equations, Institute of Computer Investigations, Moscow-Izhevsk, (2004), 360 p. (in Russian)
- [4] *C.S. Gardner, J.M. Greene, M.D. Kruskal,, R.M. Miura*, Method for solving the kortevog - de Vries equation, v.19, (1967), 1095 - 1097
- [5] *M.J. Ablowitz, H. Segur*, Solitons and the Inverse Scattering TraOn classes of integrable systems and the painleve property, J. Math. th. (SIAM, Philadelphia,1981)
- [6] *H. Airault, H.P. McKean, J. Moser*, Rational and Elliptic Solutions of the Korteveg - de Vries Equation and a related Many- Body Problem, Comm. Pure Appl. Math. 30, (1977) 95
- [7] *J. Weiss*, On classes of integrable systems and the painleve property, J. Math. Phys., 25(1) (1984) 13-24
- [8] *A.I. Yablonskii*, On rational solutions of the second Painleve equations, Vesti Akad. Nauk BSSR, Ser. Fiz. Tkh. Nauk, 3 (1959), 30 - 35 (in Russian)
- [9] *A.P. Vorob'ev*, On rational solutions of the second Painleve equations, Differential equations 1 (1965) 79 - 81 (in Russian)
- [10] *P.A. Clarkson*, Remarks on the Yablonskii - Vorob'ev Polynomials, Physics Letters A, 319, (2003) 137 - 144
- [11] *P.A. Clarkson, E.L. Mansfield*, The second Painleve equation, its hierarchy and associated special polynomials, Nonlinearity 16 (2003) R1
- [12] *P.A. Clarkson*, Painleve equations - Nonlinear Special Functions, (2004) Report of IMC, 15 december 2004
- [13] *M.V. Demina, N.A. Kudryashov*, The Yablonskii - Vorob'ev polynomials for the second Painleve hierarchy, Chaos, Solitons and Fractals, 32(2), (2007), 526 -537
- [14] *R. Conte*, The Painleve approach to Nonlinear Ordinary Differential equations, In "The Painlevé property. One century later, CRM Series in Mathematical Physics", Springer, (1999), New York, 177 - 180

- [15] *M.V. Demina, N.A. Kudryashov*, Special polynomials and rational solutions of the second Painlevé hierarchy, submitted to Theoretical and Mathematical Physics, (2006)