

# Nuclear model effects in Charged Current neutrino–nucleus quasielastic scattering

C. Maieron, M.C. Martínez and J.A. Caballero  
*Departamento de Física Atómica, Molecular y Nuclear*  
*Universidad de Sevilla, Apdo.1065, E-41080 Sevilla, SPAIN*

J.M. Udías  
*Departamento de Física Atómica, Molecular y Nuclear*  
*Universidad Complutense de Madrid, E-28040 Madrid, SPAIN*  
(Dated: March 30, 2022)

The quasielastic scattering of muon neutrinos on oxygen 16 is studied for neutrino energies between 200 MeV and 1 GeV using a relativistic shell model. Final state interactions are included within the distorted wave impulse approximation, by means of a relativistic optical potential, with and without imaginary part, and of a relativistic mean field potential. For comparison with experimental data the inclusive charged–current quasielastic cross section for  $\nu_\mu$ – $^{12}\text{C}$  scattering in the kinematical conditions of the LSND experiment at Los Alamos is also presented and briefly discussed.

PACS numbers: 25.30.Pt; 13.15.+g; 24.10.Jv

In the past few years the observation of neutrino oscillations at Super–Kamiokande [1] and the subsequent proposal and realization of new experiments, aimed at determining neutrino properties with high accuracy [2], have renovated the interest towards neutrino scattering on complex nuclei. In fact, neutrino detectors usually contain carbon or oxygen nuclei, and for a proper interpretation of the experimental results the description of the  $\nu$ –nucleus interaction must be accurate [3].

At intermediate neutrino energies, ranging from some hundreds MeV to a few GeV,  $\nu$ –nucleus quasielastic scattering has been studied within several approaches [4]. Relativistic and non relativistic studies of Random Phase Approximation have shown nuclear structure effects to be relevant only at low momentum transfers, but indications have been found that for future and precise data analyses of, e.g., atmospheric neutrino measurements, more accurate theoretical estimates may be needed. Additionally, very recently new attention has been drawn towards final state interaction (FSI) effects, which, contrary to what is often assumed, may still be relevant even at the relatively high energy  $E_\nu = 1$  GeV [5].

In this contribution we study charged–current (CC) neutrino–nucleus quasielastic scattering within the framework of a Relativistic Shell Model (RSM), already successfully employed to study exclusive electron scattering [6] and Neutral Current neutrino scattering [7]. We compute inclusive  $\nu_\mu$ – $^{16}\text{O}$  quasielastic cross sections for three values of the incident neutrino energy, namely 200 MeV, 500 MeV, and 1 GeV, which are representative of the kinematical range where quasielastic scattering gives the main contribution to the inclusive  $\nu$ –nucleus process.

We describe the CC quasielastic scattering of neutrinos on a nuclear target within the Impulse Approximation (IA), assuming that the incident neutrino exchanges one vector boson with only one nucleon, which is then emitted, while the remaining (A–1) nucleons in the target are spectators. The nuclear current is assumed to be the sum of single nucleon currents, for which we employ the usual

free nucleon expression (see [7]) with the axial form factor parameterized as a dipole with cut–off mass  $M_A = 1.026$  GeV [8], and the states of the target and residual nuclei to be adequately described by an independent particle model wave function.

To describe the bound nucleon states we use Relativistic Shell Model wave functions, obtained as the self–consistent (Hartree) solutions of a Dirac equation, derived, within a Relativistic Mean Field approach, from a Lagrangian containing  $\sigma$ ,  $\omega$  and  $\rho$  mesons [9]. As the single–particle binding energies determine the threshold of the cross section for every shell, in the numerical calculations, we have used the experimental values corresponding to the binding energies of the different shells.

For the outgoing nucleon the simplest choice is to use plane wave spinors, i.e., no interaction is considered between the ejected nucleon and the residual nucleus (PWIA). For a more realistic description, FSI effects should be taken into account. In our formalism this is done by using distorted waves which are given as solutions of a Dirac equation containing a phenomenological relativistic optical potential (ROP), consisting of a real part, which describes the rescattering of the ejected nucleon and of an imaginary part, that accounts for the absorption of it into unobserved channels. In this work we use the ROP corresponding to the EDAD–1 single–nucleon parameterization presented in ref. [10]. The use of this phenomenological ROP leads to an excellent agreement between theoretical calculations and data for exclusive ( $e, e'N$ ) observables [6]; however some caution should be taken in extending the conclusions drawn from the analysis of exclusive reactions to inclusive ones. In the latter, unless a selection of the single nucleon knockout contribution is experimentally feasible, all final channels are included and thus the imaginary term in the optical potential leads to an overestimation of FSI effects. This is clearly illustrated in fig. 1 where we compare our theoretical results with the experimental cross section corresponding to inclusive quasielastic electron scattering on




FIG. 1: Differential cross section ( $d\sigma/d\Omega d\omega$ ) versus the energy transfer for inclusive quasielastic electron scattering on  $^{12}\text{C}$  for a momentum transfer  $q \simeq 400$  MeV/c. The solid curve corresponds to the RSM with FSI described by the RMF, while the long-dashed curve corresponds to the real ROP and the short-dashed curve to the complex ROP. Finally, the dotted curve does not include FSI. Experimental data are from ref. [11].

$^{12}\text{C}$  for a momentum transfer equal to 400 MeV/c (similar results are obtained for 300 and 500 MeV/c). In our calculation besides the complex ROP, we also consider the potential obtained by setting the imaginary part of the ROP to zero. Additionally one may also use distorted waves which are obtained as the solutions in the continuum of the same Dirac equation used to describe the initial bound nucleon. We refer to this approach, that should be adequate to describe FSI at moderate energy transfer, as relativistic mean field (RMF). As shown in fig. 1, the complex ROP results clearly underestimate the data, while the reverse occurs for the PWIA calculation. On the other hand, the RMF and purely real ROP agree much better with experiment, particularly RMF for small transfer energy and real ROP for higher  $\omega$ . We believe that the RMF and real ROP results indicate a reasonable “band” where FSI effects should lie.

Let us now present our results for neutrino scattering on  $^{16}\text{O}$ . To better illustrate our model, we start by neglecting FSI and comparing RSM–PWIA results with inclusive cross sections obtained within the relativistic Fermi gas (RFG) [7]. This is done in fig. 2, which shows the differential cross section ( $d\sigma/dT_\mu$ ) as a function of the outgoing muon kinetic energy. With respect to the RFG curve calculated with no binding energy we observe that the RSM cross section is reduced and shifted towards lower  $T_\mu$  values, in a way which is similar to the effect of an average binding energy in the RFG. In addition the RSM cross section has a different shape, due to the different momentum distributions of the single nucleon shells contributing to the process. Since the various shells have different binding energies, the corresponding contributions to the cross section go to zero at differ-




FIG. 2: Differential cross section ( $d\sigma/dT_\mu$ ) versus the outgoing muon kinetic energy, for the quasielastic scattering of muon neutrinos on  $^{16}\text{O}$  and for three choices of the incident neutrino energy:  $E_\nu = 200$  MeV (upper panel), 500 MeV (middle) and 1 GeV (lower panel). The solid curves correspond to the RSM with no final state interaction, while the remaining curves are calculated within the RFG, with  $k_F = 225$  MeV and  $e_B = 0$  (dot-dashed) and  $e_B = 20$  MeV (dotted).

ent values of  $T_\mu$  and this gives rise to the structure of ( $d\sigma/dT_\mu$ ) observed at large  $T_\mu$ . Fig. 2 shows that nuclear model effects on the cross sections can be rather large at low neutrino energy, but become less relevant as  $E_\nu$  increases, practically disappearing at  $E_\nu = 1$  GeV.

On the other hand, the behavior of FSI effects is quite different, as illustrated in fig. 3. Here the results obtained with the RSM in PWIA are compared with the cases where FSI are described within the RMF, the real ROP and the complex ROP approaches. The use of real




FIG. 3: Same as fig.2, but including FSI effects. All curves are calculated within the RSM model in PWIA (solid), and within the RMF (dashed), real ROP (dotted) and complex ROP (dot-dashed) approaches.

potentials (RMF and real ROP) for describing the final nucleon states leads to the resonant structure observed for relatively high  $T_\mu$  (that is, small energy transfer  $\omega$ ). Note that in this work we only include single-particle excitations within a mean field picture. Including residual interactions would make the width and number of resonances to be considerably larger.

We observe that FSI effects produce a reduction of the cross section, particularly important in the case of the complex ROP model due to the absorption introduced by the imaginary term: about 60% for  $E_\nu = 200$  MeV and 50% for  $E_\nu = 500$  MeV and  $E_\nu = 1$  GeV in the region close to the maximum. For the RMF and real ROP, the




FIG. 4: Integrated cross section  $\sigma(E_\nu)$  for the quasielastic scattering of muon neutrinos on  $^{16}\text{O}$  as a function of the incident neutrino energy. The curves are calculated within the RFG model with  $k_F = 225$  MeV and binding energy  $e_B = 0$  (solid line) and  $e_B = 20$  MeV (dashed). The points correspond to RSM calculations without FSI (stars) and with FSI effects taken into account within the RMF (empty squares), real ROP (full squares) and complex ROP (circles) approaches.

reduction, similar in both cases, is about 30 ÷ 40% for  $E_\nu = 200$  MeV and 20% for the other energy values.

Nuclear model effects on integrated cross sections are studied in fig. 4, where the cross section  $\sigma(E_\nu)$  is plotted as a function of the incident neutrino energy. Here the contributions coming from the RMF and ROP resonances have been included in the calculation, in order to respect the completeness of the set of final states predicted by the model. These contributions are important at  $E_\nu = 200$  MeV, where they amount to about 10% of the integrated cross section, while at higher energies these effects are about 2% (500 MeV) and 1% (1 GeV). Again we see that within the PWIA the discrepancy between different nuclear models is relatively small and decreases with increasing neutrino energy. On the contrary FSI effects remain sizeable even at large  $E_\nu$ . As in the previous figure, the imaginary term in the ROP leads to a too large reduction ( $\sim 50\%$ ) of the integrated cross section. The results for the RMF and real ROP models, which are more reliable, show a smaller, but still sizeable ( $\sim 15\%$  at  $E_\nu = 1$  GeV) reduction.

The results shown in figs. 3 and 4 lead us to conclude that FSI effects should be carefully considered in neutrino experiments which use oxygen based detectors. Analogous conclusions can be drawn for the case of a  $^{12}\text{C}$  target.

For the purpose of comparison with the experiment, let us consider the inclusive  $^{12}\text{C}(\nu_\mu, \mu^-)X$  cross section measured by the LSND collaboration at Los Alamos, using a pion-Decay-in-Flight  $\nu_\mu$  beam, with energies rang-




FIG. 5: Observed distribution of muon kinetic energies  $T_\mu$  compared with the flux-averaged predictions of our RSM, in PWIA (dotted line) and including FSI within the RMF (solid) and purely real ROP (dashed) frameworks. The theoretical distributions have been normalized to give the same integrated values as the experimental points, and have been folded in energy with a bin size of 5 MeV, the same employed for the experimental data. Data are from Albert et al. [12].

ing from muon threshold to 300 MeV, and a large liquid scintillator detector [12]. This experiment has been compared to very different theoretical approaches [13], but an important discrepancy still remains. Although at these low energies processes different from the quasielastic nucleon knockout affect the inclusive cross section, we con-

sider our FSI approach to be useful to estimate integrated cross sections.

In fig. 5 we show the observed [12] and calculated distribution of events, averaged over the 1994 Los Alamos neutrino spectrum  $\phi(E_\nu)$  within the energy range  $E_\nu = 123.1 \div 300$  MeV. The shape and position of the maximum of the experimental distribution are approximately reproduced by the three calculations, but the results that include FSI with the RMF potential are clearly favored by the data. This is consistent with the fact that including FSI with the mean field potential should be adequate at moderate kinetic energy of the ejected nucleon. However the values we obtain for the the flux-averaged integrated cross section overestimate the measured cross section by approximately 50%. More precisely, in the RSM, we obtain  $\langle\sigma\rangle = 20.5$  (PWIA), 16.8 (RMF) and 15.1 (real ROP)  $10^{-40}$   $cm^2$ . Additional corrections due to the outgoing muon Coulomb distortion, evaluated within the effective momentum approximation [14], further increase these numbers by  $5 \div 10\%$ . The corresponding final measured experimental value is  $10.6 \pm 0.3 \pm 1.8$   $10^{-40}$   $cm^2$  [12].

#### Acknowledgements

This work was partially supported by DGI (Spain): BFM2002-03315, FPA2002-04181-C04-04 and BFM2000-0600 and by the Junta de Andalucía. C.M. acknowledges MEC (Spain) for a postdoctoral stay at University of Sevilla (SB2000-0427). M.C.M. acknowledges support from a fellowship from the Fundación Cámara (University of Sevilla).

- 
- [1] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81**, 1562 (1998); C. K. Jung, C. McGrew, T. Kajita and T. Mann, Ann. Rev. Nucl. Part. Sci. **51**, 451 (2001);
- [2] see for example A. De Santo, Int. J. Mod. Phys. A **16**, 4085 (2001);
- [3] “Neutrino Nucleus Interactions In The Few Gev Region”. Proceedings, 1st International Workshop, Nuint01, Tsukuba, Japan, December 13-16, 2001.
- [4] T. K. Gaisser and J. S. O’Connell, Phys. Rev. D **34**, 822 (1986); H. c. Kim, J. Piekarewicz and C. J. Horowitz, Phys. Rev. C **51**, 2739 (1995); H. c. Kim, S. Schramm and C. J. Horowitz, Phys. Rev. C **53**, 2468 (1996), *ibid.* **53**, 3131 (1996); S. K. Singh and E. Oset, Phys. Rev. C **48**, 1246 (1993); J. Engel, E. Kolbe, K. Langanke and P. Vogel, Phys. Rev. D **48**, 3048 (1993); J. Marteau, J. Delorme and M. Ericson, Nucl. Instrum. Meth. A **451**, 76 (2000).
- [5] C. Bleve *et al.*, Astropart. Phys. **16**, 145 (2001); G. Cò, C. Bleve, I. De Mitri and D. Martello, Nucl. Phys. Proc. Suppl. **112**, 210 (2002).
- [6] J.M. Udías *et al.*, Phys. Rev. C **48**, 2731 (1993); **51**, 3246 (1995); **53**, R1488 (1996); **64**, 024614-1 (2001).
- [7] W. M. Alberico *et al.*, Nucl. Phys. A **623**, 471 (1997).
- [8] V. Bernard, L. Elouadrhiri and U. G. Meissner, J. Phys. G **28**, R1 (2002).
- [9] C.J. Horowitz and B.D. Serot, Nucl. Phys. A **368**, 503 (1981); Phys. Lett. B **86**, 146 (1979); B.D. Serot and J.D. Walecka, Adv. Nucl. Phys. **16**, 1 (1986).
- [10] E.D. Cooper, S. Hama, B.C. Clark and R.L. Mercer, Phys. Rev. C **47**, 297 (1993).
- [11] P. Barreau *et al.*, Nucl. Phys. A **402**, 515 (1983).
- [12] M. Albert *et al.* [LSND Collaboration], Phys. Rev. C **51**, 1065 (1995). C. Athanassopoulos *et al.* [LSND Collaboration], Phys. Rev. C **56**, 2806 (1997); L. B. Auerbach *et al.* [LSND Collaboration], Phys. Rev. C **66**, 015501 (2002).
- [13] see N. Jachowicz, K. Heyde, J. Ryckebusch and S. Rombouts, Phys. Rev. C **65**, 025501 (2002); C. Volpe *et al.*, Phys. Rev. C **62**, 015501 (2000) 015501; E. Kolbe *et al.*, Phys. Rev. C **52**, 3437 (1995); E. Kolbe *et al.*, Nucl. Phys. A **652**, 91 (1999); Y. Umino, J. M. Udías and P. J. Mulders, Phys. Rev. Lett. **74**, 4993 (1995); Y. Umino and J. M. Udías, Phys. Rev. C **52**, 3399 (1995);
- [14] C. Giusti and F. D. Pacati, Nucl. Phys. A **473**, 717 (1987) and refs. therein.