## $v_4$ : A sm all, but sensitive observable for heavy ion collisions

Peter F.Kolb

Department of Physics and Astronomy, SUNY, Stony Brook, New York, 11794-3800, USA

(Septem ber 23, 2003)

H igher order Fourier coe cients of the azim uthally dependent single particle spectra resulting from noncentral heavy ion collisions are investigated. For interm ediate to large transverse m om enta, these anisotropies are expected to become as large as 5 %, and should be clearly measurable. The physics content of these observables is discussed from two diment extrem e but complementary view points, hydrodynam ics and the geometric limit with extrem e energy loss.

Collisions of large nuclei at nonzero in pact param eter exhibit the special feature of a strongly deform ed overlap region. The subsequent dynam ical evolution of the collision zone converts the initial spatial eccentricity into an azim uthal anisotropy of the nal state observables. In particular anisotropies in momentum space [1] have been analyzed experimentally in great detail in recent years, as they have been shown to be generated during the earliest and hottest stages of the reaction [2]. As m om entum an isotropies are generated, the eccentricity in coordinate space is smoothed out, bringing further generation of momentum anisotropies to a stall, a process that occurs within about the st 5 fm /c of the total 15 fm /c lifetime of the reball [2{4]. In Au+Au collisions at R H IC (the Relativistic Heavy Ion Collider at Brookhaven National Laboratory), those an isotropies have been found [5] to be as large as predicted by hydrodynam ic calculations that assume a short equilibration 1 fm /c [4,6]. Further theoretical investigatime <sub>equ</sub> tion has revealed the sensitivity of an isotropic ow on the equation of state of the expanding medium [7{9], indicating the necessity of a hard phase at high tem peratures with a soft transition region of width  $e 1 \text{ GeV} / \text{fm}^3$  to lower tem peratures in order to describe the data in more detail. For these reasons an isotropic ow became one of the most attractive tools to study the nuclear equation of state of the matter created in the collision by utilizing the m ost abundant particles, alternatively to study statistically disadvantaged rare probes with which one tries to infer the properties of the bulk by m odeling their m utual interaction with the medium they traverse. O ther predictions from hydrodynam ic concepts such as centrality and m ass dependence of m om entum an isotropies [8] have subsequently been con med to be in qualitative and quantitative agreem ent with experim ents [10,11], as long as the applicability of hydrodynam ics is not overstretched, i.e. for impact parameters not larger than b 7 fm and transversem om enta not to exceed pr 1:5 G eV.Beyond these values clear deviations from the hydrodynam ic predictions start to occur. This is expected as the smaller system s do not su ciently equilibrate and the most rapid particles escape the reball without fully participating in the collective dynam ics.

To date, an isotropies at m idrapidity are generally only characterized in terms of the second  $\cos c$  cient  $v_2$  of the Fourier decomposition of the azim uthally sensitive mo-

mentum spectrum. W ith the z direction given by the beam axis and the x direction de ned by the direction of the in pact parameter (and y perpendicular to both), the general expression for the Fourier decomposition with respect to the azim uthal angle  $' = \arctan(y=x)$  is

$$\frac{dN}{dp_{T} dyd'} = \frac{1}{2} \frac{dN}{dp_{T} dy} \quad 1 + \sum_{n \text{ ; even}}^{X} 2 v_{n} (p_{T}) \cos(n') \quad ; (1)$$

where no sine terms appear due to the sym m etry with respect to the reaction plane, and allodd harm onics vanish at m idrapidity due to the sym m etry ' ' + . (Note that the latter is not true for noncentral d-A u collisions, where v<sub>1</sub>, v<sub>3</sub>,... should be useful quantities to classify and study the measured spectra { even at m idrapidity). In the following we will elaborate on the centrality and m om entum dependence of higher anisotropy coe cients, which turn out to exhibit interesting features that, with the currently available data from RHIC's run in 2002, m ight already be accessible.

Generally, higherm om entum anisotropies are expected to be sm all, and in fact no experim ental data have been published until this day. An early hydrodynam ic calculation [12] predicted the momentum integrated value of  $v_4$  not to exceed 0.5 % at SPS energies even in the most peripheral collisions, and shows only a slight increase with beam energy up to R H IC energies. In another study Teaney and Shuryak [13] calculated 4, which is the fourth coe cient of the momentum integrated di erential particle spectra weighted by an additionalm om entum squared (which was applied to enhance the e ects of transverse ow). This quantity is thus strongly biased to larger  $p_T$ . From their hydrodynam ic calculation they obtain 4 1%; clearly the momentum integrated value of  $v_4$  is still sm aller.

Today, experiments at RH IC achieve su cient statistics to measure the elliptic ow coe cienty dimensional provide the entral of the entral collisions that up to 10 GeV [14]. In middentral collisions this coe cient is found to saturate starting at  $p_T = 2$  GeV at a large value of 20 { 25%, indicating that three times as many particles are emitted into the reaction plane than out of the reaction plane. Such a large value also means that the rest order deformation of Eq. (1) is not elliptic anymore, but that a polar plot of the azim uthal distribution resembles more a peanut than an elliptical shape, aligned with its

longer axis into the reaction plane, see Fig. 1. W ithout any higher order coe cients, the elliptic deform ation starts to develop such a waist when  $v_2$  becomes larger than 10 %, which can be shown explicitly by discussing the term 1+cos2' in Cartesian coordinates (x;y) around  $x\,=\,0.$ 



FIG.1. (Color online) Polar plot of the azim uthal distribution of di erent Fourier expansions. Left, distortion of the unit circle through an elliptic component  $v_2$ , shown for 10% (green) and 20% (red). Right, fourth component  $v_4$  for the same values.

This raises the interesting question, whether at high  $p_T$  Fourier coe cients of higher order m ight become suf-

ciently large to restore the elliptically deformed shape of the particle distribution. To restore the elliptic shape as much as possible only through introduction of a fourth anisotropy coe cient,  $v_4 = \frac{1}{34} (10 v_2 = 1)$  is required, as is derived straightforwardly from Eq. (1). This, in consideration of the large experimental values for  $v_2$ , m ight indicate that  $v_4$  could reach values around 3{5 %.

To investigate higher ow coe cients in more detail, we rst revisit hydrodynam ic calculations, although they nd their limitations for pions of transverse momenta beyond 2 GeV, and for heavier baryons at m om enta 3{ 4 GeV (for a recent review see Ref. [15]). Later on we will discuss a complementary model in terms of a sim ple picture for the limit of very large transverse momentum. The results presented in the following were achieved by a more modern hydrodynamic calculation than in Ref. [12], as it becomes in portant at RHIC energies to correctly account for features of chem ical nonequilibrium during the hadronic evolution of the system [16,17], a feature that signi cantly in vences the relation of energy-density and tem perature [18,19]. At tem peratures beyond  $T_{crit}$  = 165 M eV the underlying equation of state turns into a hard ideal gas equation of state via a strong rst order phase transition in order to m in ic the transition from a gas of interacting resonances

to a hard plasm a phase. The initial conditions of this calculation were determined to t single particle spectra m easured in Au+Au collisions at P = 200 GeV. For further details, see Ref. [17].

Figure 2 shows the momentum dependence of the Fourier coe cients of pion spectra up to order 8. \Elliptic ow," v dom inates the emission anisotropy at all momenta. (To use the same vertical scale for all coe - cients, the second harm onic  $v_2$  was divided by 10.) A santicipated,  $v_4$  increases with momentum, reaching a value of about 2.8 % at  $p_T = 2 \text{ GeV}$  (thus not quite enough to restore the particle distribution to an elliptic shape, as  $v_2$  ( $p_T = 2 \text{ GeV}$ ) 29%). Also  $v_6$  still has a sizable negative value of about 1.2%. The mass dependence of higher

ow harm onics does not bring any surprises. Due to their larger m asses, heavy particles pick up larger transverse m om enta when they participate in the established collective ow. As for the spectra and elliptic ow, also the higher anisotropies are shifted out toward larger transverse m om enta. Thus at a given transverse m om entum below 2 GeV, the anisotropies of heavy resonances are smaller than of pions. Com paring hydrodynam ic calculations to experim ental results, it appears how ever that deviations from hydrodynam ic behavior occur later for heavy particles, and therefore an isotropies of baryons can eventually surpass an isotropies of light m esons [20]. From these calculations one can expect a fourth ow component of (anti-)protons of about 7 % at 4 G eV transverse m om entum, a sizable quantity that should be measurable.



FIG.2. (Coloronline) Fourier coe cients of the pion spectra resulting from a hydrodynamic calculation to describe particle spectra at  $P \frac{1}{S_{N,N}} = 200 \text{ GeV}$  at impact parameter  $b = 7 \text{ fm} \cdot v_2$  is scaled by a factor 0.1.

Supprisingly it turns out that  $v_4$  is highly sensitive to the initial conditions of the calculations: initializing the calculation with a prehydrodynamic, isotropic radial ow eld as in R ef. [17] does not change the magnitude of the

ow anisotropies by a large am ount. However we nd that it completely changes the sign of  $v_4$ , while the other coe cients remain largely the same. Clearly  $v_4$  at low

The de nition of restored ellipticity is am biguous. Here we chose to nd the value of  $v_4$  where the second derivative of y(x) vanishes at x = 0, thus giving rise to a very smooth transition across ' = = 2.

and interm ediate  $p_T$  has therefore a strong potential to constrain m odel calculations and carries valuable information on the dynam ical evolution of the system. To study how the higher ow anisotropies develop in the course of the hydrodynam ic evolution we calculate moments of the transverse ow eld  $\langle y; v_v \rangle$  through

R

$$f_{n}(x) = \frac{\frac{dxdye(x;y; x) v_{T} \cos(n'_{v})}{R}}{dxdye(x;y; x)}; \qquad (2)$$

where e(x;y; ) is the energy density in the transverse plane at time and  $v_T = (v_x^2 + v_y^2)^{1=2}$  is the transverse

ow velocity at a given point (x;y), = (1)  $v_{T}^{2}$ ) <sup>1=2</sup>, and  $v_{v} = \arctan(v_{v} = v_{x})$  is the angle of the local ow velocity with respect to the reaction plane. (Note that for n = 0 this reduces to the de nition of the mean radial velocity used in earlier studies [4], but for n = 2it is slightly di erent from the momentum an isotropy p which is de ned in terms of the di erence in the diagonal elements of the energy momentum tensor.) Figure 3 displays the time evolution of the ow coe cients and clearly shows that radial ow 0 increases throughout the lifetime of the system (which is about 15 fm /c), with a som ew hat reduced acceleration while most of the system is in the mixed phase at around 4 fm /c. Quite oppositely anisotropy components quickly saturate and remain constant, as the initial geom etric deform ation of the source is lost. About 2/3 of the nalvalue of  $_4$  is generated during the earliest stages of the evolution, where the system is governed by the hard equation of state of the quark gluon plasma. 6 quickly achieves small negative values of the order of -0.2% (as expected from the negative value of  $v_6$ ). Due to its small values, it is subject to num erical uctuations and therefore not shown. It is expected that the ow coe cients are monotonically related to the anisotropy coe cients in the nalparticle spectra. Details of this relation are subject to future study.



FIG.3. (Color online) T in e evolution of the anisotropy coe cients in the transverse velocity eld as given by the hydrodynamic calculation.  $_0$  characterizes the radial ow,  $_2$  the elliptic deform ation and  $_4$  the quadrupole deform ation.

In a model complementary to the hydrodynamic description which is constrained to the low momen-

tum, therm alized part of the spectrum, one can study anisotropies that result from opacity e ects that the deform ed initial geom etry exerts on particles of high m om enta [21{23]. A ssum ing extrem e opacity, or extrem e jet quenching, particles of large transversem on enta from any part of the surface can only be em itted within 90 of the norm al vector to the surface, which for this exploratory study is assumed to be given by the overlap of two hard spheres (radius R).<sup>y</sup> In this picture, the nth Fourier coe cient of the resulting expansion is found to be (for n even)

$$v_n = (1)^{n=2} \frac{1}{1 n^2} \frac{\sin(n)}{n};$$
 (3)

where  $= \arccos(b=2R)$  characterizes the centrality in term s of the opening angle between the line through the center of one nucleus to the intersection point of the two nuclear spheres and the reaction plane. Note that as b increases from 0 to 2R, decreases from =2 to 0. For n = 2,  $v_2 = sin(2) = (6)$ , a formula rst given by Voloshin [26]. For n > 2, Eq. 3 becomes signi cantly m ore interesting (unfortunately, how ever, the size of the expected signal drops with  $n^3$ ). Note rst the oscillating sign of the prefactor which will manifest itself in an oscillation of the coe cients at large b (sm all ). Second, one notices that for larger n the sine function starts changing sign upon scanning im pact parameters (b \$ ), whereas for n = 2, the Fourier coe cient increases linearly with impact parameter from 0 to 1/3, and is thus always positive. In particular for n = 4 one observes a transition from positive to negative values at b = 2R, reaching the limiting value  $v_4 = 1=15$ 6:67% for b! 2R, again a rather large number. The local maxin un of  $v_4$  is at b 0:865R with v4 1:45 % . The centrality dependences of the second to eighth Fourier coe cients are shown in Fig. 4 (as before  $v_2$  is divided by a factor of 10).

C learly, the expected values for higher expansion coefcients from this calculation are small. A pplying a more realistic model does, how ever, not necessarily lead to a further reduction of the signal, but might in contrast increase, similarly as observed for calculations of elliptic ow [24]. This is also suggested by the experimental data on  $v_2$  which seem to exceed the limits given even by the most optimistic assumptions for anisotropies from geom – etry and energy loss. In any case, such a distinctive signal as a sign change upon variation of centrality should be preserved when model calculations get rened and should be accessible within the statistics reached by today's experiments.

<sup>&</sup>lt;sup>y</sup>R elaxing this geom etrical constraint by adopting a nite di useness of the nuclei reduces the resulting asym m etries, far below the experim ental results, and novele ects are required to explain those large limiting values [24,25].



FIG. 4. (Color online) Centrality dependence of the anisotropy coe cients in the limit of pure surface emission from the deformed overlap region to describe the saturation value of the anisotropies at large transverse momenta.  $v_2$  is scaled by a factor of 0.1.

As an aside we would like to discuss a new but naive experimental way of measuring higher order ow coefcients, in particular y. Apart from the \straightforward" method to determ ine each event's reaction plane and averaging all particles'  $\cos(4')$  with respect to the reaction plane and the more recently developed cum ulant technique [27], sim ply measuring the particle spectra with respect to the reaction plane might be su cient (theoretically). Having the azim uthally sensitive particle spectra given, Eq. (1) shows directly that [abbreviating dN =pr dpr dyd' (pr;') = n(')]

$$n (0) + n (=2) 2n (=4) =$$

$$= \frac{1}{2} \frac{dN}{p_{T} dp_{T}} [8 v_{4} (p_{T}) + 8 v_{12} (p_{T}) + :::]: (4)$$

From the experience gathered so far we expect  $v_{12}$  to be negligibly sm all compared to  $v_4$ . Finite opening angles for the azim uthally sensitive distribution will reduce the signalagain, how ever, the  $v_4$  is in this approach enhanced by a factor of 8, which m ight still render it m easurable. W e propose this method thus as an interesting independent approach to measure  $v_4$  (but also  $v_2$ , by sim ply subtracting spectra in and out of the reaction plane).

In sum m ary, we have shown that  $v_4$  ( $p_T$ ) m ight achieve signi cantly large values at interm ediate and large transverse m om enta in noncentral heavy ion collisions and contains in portant physics of heavy ion collisions, a fact largely overlooked in the past. Num erical estim ates for its size were given both in the lim it of full them alization w ith a subsequent hydrodynam ic expansion as well as in the geom etrical lim it of extrem e \jet quenching." For the latter case a strong centrality dependence and a change of the sign of the signal for higher anisotropies was found. From both approximations we found that  $v_4$ m ight reach values of the order of 5%, clearly observable w ith todays equipm ent and statistics.  $v_4$  has been found to be highly dependent on the initial conguration of the system and its evolution and is thus an important new tool to constrain m odel calculations and analyze details of the system 's history. Consequently it deserves to be studied in great detail both experim entally as well as theoretically.

A cknow ledgm ents: D iscussions with J.C asalderrey-Solana, U. Heinz, B. Jacak, J. Jia, A. Poskanzer, E. Shuryak, and R. Snellings are greatly appreciated. This work was supported by the U.S. Department of Energy under G rant No. D E-FG 02-88ER 40388 and by the A lexander von H um boldt Foundation.

- [1] J.-Y.Ollitrault, Phys. Rev. D 46, 229 (1992).
- [2] H. Sorge, Phys. Rev. Lett. 78, 2309 (1997).
- [3] B. Zhang, M. Gyulassy, C. M. Ko, Phys. Lett. B 455, 45 (1999).
- [4] P F. Kolb, J. Sollfrank, and U. Heinz, Phys. Rev. C 62, 054909 (2000).
- [5] K H.Ackerm ann et al. (STAR Collaboration), Phys. Rev. Lett. 86, 402 (2001).
- [6] P.F. Kolb, P. Huovinen, U. Heinz, and H. Heiselberg, Phys.Lett. B 500, 232 (2001).
- [7] D. Teaney, J. Lauret, and E. Shuryak, nucl-th/0110037; Phys. Rev. Lett. 86, 4783 (2001).
- [8] P.Huovinen, PF.Kolb, U.Heinz, P.V.Ruuskanen and SA.Voloshin, Phys.Lett. B 503, 58 (2001).
- [9] P.F.Kolb and U.Heinz, hep-ph/0204061.
- [10] C.Adler et al. (STAR Collaboration), Phys. Rev. Lett. 87, 182301 (2001); Phys. Rev. Lett. 89, 132301 (2002).
- [11] K.Adcox et al. (PHENIX Collaboration), Phys.Rev. Lett.89,212301 (2002).
- [12] P F.Kolb, J.Sollfrank, and U.Heinz, Phys.Lett.B 459, 667 (1999).
- [13] D. Teaney and E. Shuryak, Phys. Rev. Lett. 83, 4951 (1999).
- [14] K. Film onov et al. (STAR Collaboration), Nucl. Phys. A 715, 137c (2003).
- [15] PF.Kolb and U.Heinz, nucl-th/0305084.
- [16] R.Rapp, Phys.Rev.C 66, 017901 (2002).
- [17] P.F.Kolb and R.Rapp, Phys. Rev. C 67, 044903 (2003).
- [18] D. Teaney, nucl-th/0204023.
- [19] T.Hirano and K.Tsuda, Phys. Rev. C 66, 054905 (2002).
- [20] S.S. Adler et al. (PHENIX Collaboration), nuclex/0305013; J.Adam set al. (STAR Collaboration), nuclex/0306007.
- [21] M. Gyulassy, I. V itev, X.-N. W ang, and B.-W. Zhang, nucl-th/0302077.
- [22] P.Huovinen, PF.Kolb, and U.Heinz, Nucl. Phys. A 698, 475 (2002).
- [23] E. Shuryak, Phys. Rev. C 66, 027902 (2002).
- [24] J.Casalderrey-Solana and E.Shuryak, hep-ph/0305160.
- [25] D.Molnar and S.A. Voloshin, nucl-th/0302014;
- R.J.Fries, B.Muller, C.Nonaka, and S.Bass, nuclth/0306027.
- [26] S.Voloshin, Nucl. Phys. A 715, 379 (2003).
- [27] N. Borghini, P. Dinh, and J.-Y. Ollitrault, Phys. Rev. C 64, 054901 (2001).