# The sigm a - nucleon form factor in a chiral topological model with anom alous dimension.

A bdulla R akhim ov Institute of Nuclear Physics, Tashkent, U zbekistan

## Abstract

The chiral soliton m odel including ; ;! m esons is extended for the case when the scalar – isoscalar m eson has an anom alous dimension – d. The form factor of the sigm a – nucleon interaction calculated and to be found not highly sensitive to the value of d.

E-m ail: rakhim ov@ rakhim ov.ccc.uz

#### I. IN TRODUCTION

It is well known that, the scalar isoscalar -m eson plays an important role in nucleon nucleon (NN) interaction. In one boson exchange model (OBE) of NN interaction one has totake into account the form <math>-factor of NN vertex which is still poorly known. In our recent paper [1] we considered the form -factor in the fram ework of ! soliton model includinga dilaton eld as a sigm a meson with the ordinary scale dimension <math>d = 1. On the other hand, there are some versions of chiral elective Lagrangians for the nite nuclear matter and nite nuclei where light scalar degrees of freedom have an anom alous scale dimension [2].

Note that, anom alous, fractal, scale dimensions are ingredient feature of modern theories. In particular it has been recently shown that, fractal scale dimension is essential to describe superconductivity [3]. In present brief note we study the role of anom alous scale dimension in NN interaction. The paper is organized as follows. In Sect. II we outline the scaling properties of basic elds; in Sect. III we shall extend the ; ;! Lagrangian for the case of anom alous dimension; in Sect. IV we calculate the vertex form factor of N interaction. The results and brief summary are presented in Sect. V

II. SCALING BEHAVIOR OF SCALAR AND SPINOR FIELDS

In this section, to make the point clearer and for further references we brie y discuss the scaling properties of various elds. For illustration we consider kinetic term s of ferm ion and boson elds. The kinetic term s of a ferm ion eld (e.g. quark eld in QCD, or nucleon eld) is given by

$$L_{kin}(x) = i(x)0$$
 (2.1)

and a boson ( or ) eld

$$L_{kin}(r) = \frac{1}{2}0 \quad 0 \quad :$$
 (2.2)

The scale invariance dem ands that the action  $S = {}^{R} d^{4}xL$  should be invariant under scale transform ations  $x = e x^{0}$ :

$$S = \int_{-\infty}^{Z} d^{4}x L_{kin}(x) = \int_{-\infty}^{Z} d^{4}x^{0} L_{kin}^{0}(x^{0})$$
 (2.3)

Let the ferm ion and boson elds scale as  $(x) = \exp(n)^{0}(x^{0})$  and  $(x) = \exp(m)^{0}(x^{0})$ ; respectively. For ferm ions we get

$$S = i d^{4}x (x) (x) = i d^{4}x^{0} (x^{0}) (x^{0}) (x^{0}) = e^{2n} e^{4} = d^{4}x^{0} L^{0}_{kin} (x^{0})$$
(2.4)

and hence 4 + 2n = (2n - 3) = 0. Since is arbitrary then n = 3=2. Similarly, for boson elds one can show that m = 1. Therefore we may conclude that

x ! e x; (x) ! 
$$e^{3} = 2$$
 (x); (x) ! e (x); (2.5)

i.e. the scale dimension dequals to  $d_f = 3=2$  and  $d_b = 1$  for fermions and bosons respectively. Thus, the term "anom alous scale dimension" e.g. for a boson eld means  $d_b \in 1$ .

In Skyme like models the pion eld is involved through a chiral nonlinear eld  $U = \exp(i \sim -f)$  which has trivial scale dimension  $d_U = 0$ . For this reason the four derivative term of the Skyme Lagrangian given by

$$L_{4} = \frac{1}{32e^{2}} Tr[(@ U)U^{+}; (@ U)U^{+}]^{2}$$
(2.6)

is scale invariant by itself:

$${}^{Z} d^{4}xL_{4}(x) ! {}^{Z} d^{4}x^{0}e {}^{4} L_{4}^{0}(x^{0})e^{4} = {}^{Z} d^{4}x^{0}L_{4}^{0}(x^{0});$$
(2.7)

whereas the kinetic term

$$L_{2}(x) = \frac{f^{2}}{4} Tr[0 U(x)0 U^{+}(x)]$$
(2.8)

is not:

$$\begin{array}{c} {}^{Z} d^{4}xL_{2}(x) ! \frac{f^{2}}{4} d^{4}x^{0}e^{-4} e^{2} \operatorname{Tr} \left[ 0 U^{0}(x^{0}) \left( U^{0+}(x^{0}) \right) \right] = \\ = e^{-2} d^{4}x^{0}L_{2}^{0}(x^{0}) \end{array}$$

$$(2.9)$$

How can the invariance be restored? A coording to Shehter [4] an additional boson eld (x) should be included into the Skyrm e Lagrangian. We shall use this prescription in the next Section.

W e start with extended Skyme - like Lagrangian including vector m esons explicitely [5]

$$L = \frac{f^{2}}{4} \operatorname{Tr} L \quad \frac{f^{2}}{2} \operatorname{Tr} [1 + r + ig \sim - + ig!]^{2} + \frac{3}{2} g! \quad B \quad \frac{1}{4} (! ! + - - )$$
$$+ \frac{f^{2} m^{2}}{2} \operatorname{Tr} (U \quad 1); \qquad (3.1)$$

where left/right{handed currents are given by  $L = U^+ (0, 1) = ($ 

given by B = " TrL L L = (24<sup>2</sup>). Note that, for the case of in nite and ! meson masses the Lagrangian in Eq. (3.1) becomes equal to the usual Skyrme -W itten Lagrangian with two, four and six derivative terms and, hence, is not scale invariant. As it has been mentioned above this shortcom ing can be restored by inclusion of a dilaton eld - sigm a meson. Thus, including a {meson by means of the scale invariance and trace anomaly of QCD into the Lagrangian one obtains following chiral Lagrangian of the coupled ! system

$$L = \frac{S_0^2 e^{2 - d}}{2} @ @ \frac{f^2 e^{2 - d}}{4} TrL L \frac{f^2 e^{2 - d}}{2} Tr[L + r + ig \sim \sim +$$
  
+ ig!  $j^2 + \frac{3}{2}g! B \frac{1}{4}(! + \sim \sim) + \frac{f^2 m^2 e^{3 - d}}{2} Tr(U - 1)$  (3.2)  
$$\frac{d^2 S_0^2 m^2}{16} [L - e^{4 - d}(\frac{4}{d} + 1)];$$

Here the scale dimension of eld (d 1) is included explicitly. In Eq.(3.2)  $S_0$  is the vacuum expectation value of scalar eld in free space matter, f is the pion decay constant (f = 93 M eV) and g = g is determined through the K SFR relation g = m = 2f. The model assumes the masses of and ! mesons to be equal,  $m_q = m_1 = m$ . The mass of the is related to the gluon condensate in the usual way  $m_s = 2$  C  $_q = (dS_0)$  [2].

Nucleons arise as soliton solutions from the Lagrangian Eq.(3.2) in the sector with baryon number B = 1. To construct them one goes through a two step procedure. First, one nds the classical soliton which has neither good spin nor good isospin. Then an adiabatic rotation of the soliton is performed and it is quantized collectively. The classical soliton follows from Eq.(3.2) by virtue of a spherical sym metrical ansatze for the meson elds:

$$U(\mathbf{r}) = \exp(i_{r} \hat{\mathbf{r}} (\mathbf{r})); \quad i_{i}^{a} = "_{iak} \hat{\mathbf{r}}_{k} \frac{G(\mathbf{r})}{gr}; \quad ! \quad (\mathbf{r}) = ! \quad (\mathbf{r}) \quad _{0}; \quad (\mathbf{r}) = (\mathbf{r}) : \quad (3.3)$$

In what follows we call (r), G (r), ! (r), and (r) the pion{, {, ! {, and {m eson prole functions, respectively. The pertinent boundary conditions to ensure baryon num berone and nite energy are, (0) = ;G (0) = 2;!<sup>0</sup>(0) = <sup>0</sup>(0) = 0; (1) = G (1) = ! (1) = (1) = 0. To project out baryonic states of good spin and isospin, we perform a time { independent SU (2) rotation

$$U(\mathbf{x};t) = A(t)U(\mathbf{x})A^{+}(t); \quad (\mathbf{x};t) = A(t)(\mathbf{x})A^{+}(t)$$

$$(\mathbf{x};t) = (\mathbf{r}); \quad ! (\mathbf{x};t) = \frac{(\mathbf{r})}{\mathbf{r}} [\mathcal{K} \hat{r}]$$

$$\sim {}_{0} \cdot (\mathbf{x};t) = \frac{2}{g} A(t) \sim \mathcal{K}({}_{1}(\mathbf{r}) + \hat{r}\mathcal{K} - \hat{r}_{2}(\mathbf{r}))A^{+}(t);$$

$$\sim {}_{i} \cdot (\mathbf{x};t) = A(t) \sim {}_{i} \cdot (\mathbf{r})A^{+}(t)$$
(3.4)

with 2K the angular frequency of the spinning mode of soliton, i~  $K = A^+A$ . This leads to the time (dependent Lagrange function

$$L(t) = dr L = M_{H}(;G;!;) + (;G;!;;;_{1};_{2})Tr(AA^{+}): (3.5)$$

M inimizing the classical mass M<sub>H</sub> (;G;!; ) leads to the coupled di erential equations for ;G;! and subject to the aforem entioned boundary conditions. In the spirit of the large N<sub>c</sub> (expansion, one then extrem izes the moment of inertia (;G;!;;;;1;2) which gives the coupled di erential equations for  $_1$ ,  $_2$  and in the presence of the background proles ;G;! and . The pertinent boundary conditions are (0) = (1) = 0;  $_1^0(0) = _1(1) = 0$ ;  $_2^0(0) = _2(1) = 0$ ;  $_1^0(0) + _2(0) = 2$ : The masses of nucleon M<sub>N</sub> and the mass of ,M, are then given by M<sub>N</sub> = M<sub>H</sub> + 3=8 and M = M<sub>H</sub> + 3=15.

### IV.VERTEX FORM FACTOR OF N INTERACTION

The meson - nucleon form factors can be derived by a well known procedure, proposed long years ago by C ohen [6,7]. A lthough they were derived in a microscopical and consequent way, these form factors could not be directly used in standard OBE schemes of nucleon nucleon interaction. The reason is that the OBE schemes [8] in momentum space use form factors de ned for the elds propagating on a at metric, whereas the de nition of form factors in C ohen's procedure involve a nontrivial metric. Hence, before using them in OBE scheme one should modify the procedure by rede ning meson elds. Note that, the modi cation for pion nucleon form factors in ! model is clearly outlined in refs. [1,9].

Rede nition of a meson eld i.e. introduction of a at metric is based on a canonical form for the kinetic part of the Lagrangian, which determines the dynamics of the eld uctuation. The kinetic term of the scalar meson in Eq. (3.2)

$$L^{kin} = S_0^2 e^{2 - d} [0 \quad (0 = 2)$$
(4.1)

can be easily rewritten in a usual way:

$$L^{KIN} = 0 \sim 0 \sim = 2$$
 (4.2)

by the following rede nition of the basic sigma eld:

$$\sim (r) = S_0 d[1 e^{(r)=d}]$$
: (4.3)

Now the new eld ~ may be identied with a real sigma eld. It can be also easily shown that the above rede nition does not modify the mass:  $m_{-} = m_{-}$ .

Now we apply Cohen's procedure to derive NN form factor. K lein -G ordon equation for the scalar-isoscalar meson interacting with nucleon with the Lagrangian L  $_{NN} = g_{NN} \sim$  is given by:

$$(\tilde{r}^{2} + m^{2} \sim) = j$$
 (4.4)

with the source  $j(x) = g_{NN}$  (x) (x). The rst step of the procedure assumes taking matrix elements of both sides of Eq. (4.4) between nucleon states N ( $\mathcal{P}^0$ ) and N ( $\mathcal{P}$ ). Evaluating the matrix element of the source in Breit frame, ( $\mathcal{P}^0 = \mathcal{P}, q = \mathcal{P} - \mathcal{P}^0$ ) one can nd

$$\begin{array}{l} \text{hN} \ (\mathcal{P}^{0}) \, \widetilde{j}_{1}(\mathbf{r}) \, \widetilde{j}_{1} \ (\mathcal{P}) \, i = \frac{\exp \left( i \mathbf{q} \, \mathbf{r} \right)}{(2 \ )^{3}} \, \text{hN} \ (\mathcal{P}^{0}) \, \widetilde{j}_{1}(0) \, \widetilde{j}_{1} \ (\mathcal{P}) \, i = G_{\text{NN}} \ (t) \, u \, (\mathcal{P}^{0}) \, u \, (\mathcal{P}) \\ \\ = \frac{E_{\text{q}}}{2M} \, \left[ \mathbb{I} + \frac{\mathbf{q}^{2}}{4 \, (E_{\text{q}} + M_{\text{q}})^{2}} \, \left] G_{\text{NN}} \ (t) \ G_{\text{NN}} \ (t) \, \text{T} \ (\mathbf{q}^{2}) \right]; \end{array}$$

$$(4.5)$$

where  $t = q^2$ , and u (p) is D irac spinor

$$u(p) = \frac{E_p + M}{2M} \stackrel{1=2}{(\sim p) = (E_p + M)} s;$$
 (4.6)

with the normalization u(p)u(p) = 1. The similar matrix element for the eld with collective coordinates r is given by:

hN 
$$(\mathcal{P}^{0})$$
 j~  $(\mathbf{r} \ \mathcal{R})$  j)  $(\mathcal{P})$  i =  $\frac{1}{(2)^{3}}^{Z}$  d $\mathcal{R}$  exp (iq $\mathcal{R}$ ) u  $(\mathcal{P}^{0})$  ~  $(\mathbf{r} \ \mathcal{R})$  u  $(\mathcal{P})$   
=  $\frac{\exp(iqr)T(q^{2})}{(2)^{3}}^{Z}$  d $\mathbf{x}$  exp ( iq $\mathbf{x}$ ) ~ ( $\mathbf{x}$ ); (4.7)

where  $\mathcal{R} = r$  x. Now using Eqs. (4.5) and (4.7) in Eq. (4.4) one obtains

$$dx \exp(iqx)(q^2 + m^2) \sim (x) = G_{NN}(t)$$
: (4.8)

For the spherical symmetric ansatz with  $\sim (x) = \sim (jxj)$  the left hand side of Eq. (4.8) can be easily evaluated:

<sup>Z</sup>  

$$d\mathbf{x} \exp(i\mathbf{q}\mathbf{x}) (\mathbf{q}^{2} + \mathbf{m}^{2}) \sim (\mathbf{x}) = {}^{Z} d\mathbf{x} d\mathbf{x} \mathbf{x}^{2} \exp(i\mathbf{q}\mathbf{x}) [\hat{\mathbf{r}}^{2} + \mathbf{m}^{2}] \sim (\mathbf{x})$$

$$= 4 {}^{Z_{1}}_{0} d\mathbf{x} \mathbf{f} \mathbf{j}_{0} (\mathbf{q}\mathbf{x}) [2\mathbf{x} \sim^{0} (\mathbf{x}) + \mathbf{x}^{2} \sim^{00} (\mathbf{x})] + \mathbf{j}_{0} (\mathbf{q}\mathbf{x}) \mathbf{x}^{2} \mathbf{m}^{2} \sim (\mathbf{x}) \mathbf{g};$$
(4.9)

where  $j_0(x) -$ spherical Bessel function.

Ζ

F inally, inserting this into the Eq. (4.8) we get following expression for the NN form factor

$$G_{NN}(t) = 4 \int_{0}^{Z_{1}} dx j_{0}(qx) fx^{2} \sim^{00}(x) + 2x \sim^{0}(x) x^{2}m^{2} \sim (x)g \qquad (4.10)$$

Note that, ~ (x) is de ned in Eq. (4.3), where (x) is the solution of the coupled nonlinear equations of the motion, obtained by minimizing M  $_{\rm H}$ .

### V.RESULTS AND SUMMARY.

Before doing actual calculations the parameters of the model should be claried. As can be seen from Eq.(3.2), the Lagrangian has no free parameters in the ! sector. So, in actual calculations the parameters m; m; f are xed at their emperical values, m = 138 MeV,  $m = m = m_{\perp} = 770 \text{ MeV}$ , f = 93 MeV, g = m = 2f = 5.85. In the {m eson sector there

are in general three free parameters:  $m \, , S_0$  and the scale dimension d. The values for  $S_0 - w$  ere found to be  $S_0 = 90.6 \, 95.6 \, \text{MeV} \, [2]$ . So we put  $S_0 = f = 93 \, \text{MeV}$ . As to the mass of sigm a meson m we set m = 550M eV to get the best description of static nucleon properties. To study the role of the scale dimension we shall consider two cases: d = 1 and d = 2.4 corresponding to the norm aland anom abus cases respectively.

The NN form factor G<sub>NN</sub> (t) given in Eq. (4.10) is presented in Fig.1. The solid and dashed curves are for d = 2.4 and d = 1 cases respectively. It is seen that both curves coincide qualitatively, so that, being used in OBE model they would lead to the same NN potential. Thus, the sensitivity of G<sub>NN</sub> (t) to the value of d need not to be large.

We point out that the sigm a meson {nucleon form factor found in the present model could be useful in a wider context of calculations of nucleon {nucleon observables (phase shifts, deuteron properties etc) and may give more information on meson {nucleon and nucleon { nucleon dynamics.

To sum marize, we have developed a topological chiral soliton model with an explicit light scalar (isoscalar meson eld with anom alous dimension, which plays a central role in nuclear physics, based on the chiral sym metry and broken scale invariance of QCD. We have shown that, the anom alous value of scale dimension would not lead to substantial modi cation of the NN vertex form - factor.

#### ACKNOW LEDGMENTS

This work is supported by Uzbek Science Foundation 2:1:25.

## REFERENCES

[1] UlfG.Meiner, A.Rakhim ov and U.Yakhshiev PhysLett.B473, (2000), 200.

[2] R.J. Fumstahl, Hua-Bin Tang, Brian D. Serot, Phys. Rev. C 52, (1995), 1368.

[3] C.K.Kim, A.Rakhim ov and J.H.Yee, Phys. Rev. B71, 024518 (2005).

[4] J. Schechter PhysRev D 21, (1980), 3393.

[5] Ulf-G.Meiner, N.Kaiser and W.Weise, Nucl. Phys. A 466, (1987), 685.

[6] T D. Cohen, Phys. Rev., D 34, (1986), 2187.

[7] N.Kaiser, U.Vogl, W.Weise and UlfG.Meiner, Nucl. Phys. A 484, (1988), 593.

[8] R.Machleidt, K.Holinde and C.Elster, Phys. Rep. 149, (1987), 1.

[9] G. Holzwarth and R. Machleidt, Phys. Rev. C 55, (1997), 1088.



FIG.1. The normalized NN form factor, G  $_{NN}$  (t). The solid and dashed lines are for d = 2:4 and d = 1 cases respectively.