

## Comment on "Constraints on proton structure from precision atomic physics measurements"

In a recent letter [1] Brodsky *et al.* used experimental and theoretical information on the hyperfine structure (hfs) of hydrogen and muonium to derive the proton structure correction. In our previous work [2] on this topic we used only information on the hydrogen hfs to calculate a corresponding correction, which was consistent with that of [1]. The structure correction can be decomposed into a dominant static part (the Zemach term, proportional to the Zemach moment,  $\langle r \rangle_{(2)}$ ) and a dynamic part (the polarization correction). In [2] we directly determined the Zemach term from the (experimental) electron-scattering form factors that define it. From this result we were able to *infer* the polarization correction. Reference [1] used a value of the polarization correction calculated in [3] to infer the Zemach term. The two results for  $\langle r \rangle_{(2)}$  are in severe disagreement and we strongly disagree with the conclusions of [1].




FIG. 1: (a) Correlation between  $r_{rms}$  and  $\langle r \rangle_{(2)}$ ; the parametrizations of [1] are labeled by #, and are fit by the dotted line. (b)  $\chi^2$  of the parametrizations employed in [1].

Although the resulting  $\langle r \rangle_{(2)}$  of Ref. [1] disagrees with the precise value from electron scattering [2], Brodsky *et al.* claim that combinations of model form factors  $G_e^p(q)$  and  $G_m^p(q)$  exist that are compatible with their  $\langle r \rangle_{(2)}$ . This conclusion is not justified. Values of the proton charge rms-radius  $r_{rms}$  and  $\langle r \rangle_{(2)}$  for 10 different combinations of  $G_e^p$  and  $G_m^p$  are listed in Table 1 of [1]. All values of  $\langle r \rangle_{(2)}$  and several values of  $r_{rms}$  are numerically incorrect; the correct values are displayed in Fig. (1a). This figure shows that no combination of proton form factors is compatible with the upper green symbol, which depicts the  $\langle r \rangle_{(2)}$  of Ref. [1] and the  $r_{rms}$  derived from the hydrogen Lamb shift [4] (which agrees with electron scattering [5]). The figure also clearly shows that there is a tight correlation between  $r_{rms}$  and  $\langle r \rangle_{(2)}$ . The  $\langle r \rangle_{(2)}$  of Brodsky *et al.* is incompatible with all other infor-

mation if any modern value of  $r_{rms}$  is correct. Our result in red is compatible.

Table 1 of [1] and the spread of values in Fig. (1a) might suggest that electron scattering does not accurately determine  $\langle r \rangle_{(2)}$ , notwithstanding the precise result of  $1.086 \pm 0.012$  fm in [2]. Figure (1b) shows that this conclusion is not justified. For the range of momentum transfers relevant for the determination of  $\langle r \rangle_{(2)}(q < 4 fm^{-1})$ , most of the  $G_e^p, G_m^p$ -combinations of [1] give a disastrous  $\chi^2$  when compared to the *world* electron scattering data; they simply do not fit that data. The only exceptions are fits 3 and 4, which differ from the one of Refs. [2,5] mainly by the omission of Coulomb distortion, which explains the slightly low value of  $r_{rms}$ .

Figure (1a) also points out the sensitivity of Ref. [1]'s determination of  $\langle r \rangle_{(2)}$  to the polarization correction (omitting it leads to the lower green symbol). This correction has been calculated by Faustov *et al.* [3]. It depends on the proton spin structure functions  $g_1(q, x)$  and  $g_2(q, x)$  at low  $q$  and low energy loss  $\nu$  (large  $x$ ), where these structure functions are poorly known. Modeling with the constituent quark model (as in [3]) might introduce errors. The uncertainty attributed in Ref. [3] to the polarization correction was a subjective theoretical estimate (unlike the uncertainty in [2]), and we believe that it has been substantially underestimated.

We conclude that the technique of Brodsky *et al.* to use hydrogen and muonium hfs (together with estimates of the polarization correction) to calculate  $\langle r \rangle_{(2)}$  will remain inaccurate until the polarization correction can be calculated using more and better inelastic electron scattering data. Until that time the most reliable value for  $\langle r \rangle_{(2)}$  will come from elastic electron scattering.

J.L. Friar

Los Alamos National Laboratory  
Los Alamos, NM 87545

I. Sick

University of Basel  
CH4056 Basel, Switzerland

DOI

PACS numbers: 14.20Dh, 13.40.Gp, 31.30.-i

- [1] S.J. Brodsky *et al.*, Phys. Rev. Lett. 94(2005)033002
- [2] J.L. Friar and I. Sick, Phys. Lett. B 579(2004)285
- [3] R.N. Faustov *et al.*, Eur. Phys. J. C24(2002)281
- [4] P.J. Mohr *et al.*, Rev. Mod. Phys. 77(2005)1
- [5] I. Sick, Phys. Lett. B576(2003)62