E ects of jet quenching on the hydrodynam ical evolution of quark-gluon plasm a

A.K.Chaudhuri¹, and Ulrich $Heinz^2$

¹Variable Energy Cyclotron Centre, 1-AF, Bidhan Nagar, Kolkata - 700 064, India ²Department of Physics, The Ohio State University, Columbus, OH 43210, USA

(Dated: April 16, 2024)

W e study the e ects of jet quenching on the hydrodynam ical evolution of the quark-gluon plasm a (QGP) uid created in a heavy-ion collision. In jet quenching, a hard QCD parton, before fragmenting into a jet of hadrons, deposits a fraction of its energy in the medium, leading to suppressed production of high- p_T hadrons. A ssum ing that the deposited energy quickly thermalizes, we sim – ulate the subsequent hydrodynam ic evolution of the QGP uid. For partons moving at supersonic speed, $v_p > c_s$, and su ciently large energy loss, a shock wave form s leading to conical ow [1]. The PHENIX Collaboration recently suggested that observed structures in the azim uthal angle distribution [2] m ight be caused by conical ow . We show here that, for phenom enologically acceptable values of parton energy loss, conical ow e ects are too weak to explain these structures.

PACS num bers: PACS num bers: 25.75.-q, 13.85 Hd, 13.87.-a

Recent Au+Au collision experiments at the Relativistic Heavy Ion Collider (RHIC) saw a dramatic suppression of hadrons with high transverse momenta (high- p_T suppression") [3], and the quenching of jets in the direction opposite to a high- p_T trigger particle [4, 5], when compared with p+p and d+Au collisions. This is taken as evidence for the creation of a very dense, color opaque medium of decon ned quarks and gluons [6]. Independent evidence for the creation of dense, therm alized quark-gluon matter, yielding comparable estimates for its initial energy density (rei $> 10 \text{ GeV}/\text{fm}^3$ at time them < 0.6 fm/c [7]), comes from the observation of strong elliptic ow in non-central Au+Au collisions [3], consistent with ideal uid dynam ical behaviour of the bulk of the matter produced in these collisions.

These two observations raise the question what happens, in the sm all fraction of collision events where a hard scattering produces a pair of high- p_T partons, to the energy lost by the parton travelling through the medium . The STAR Collaboration has shown that, while in central Au+Au collisions there are no hard particles left in the direction opposite to a high- p_T trigger particle, one sees enhanced production (com pared to p+p) of soft (low $-p_T$) particles, broadly distributed over the hem isphere diam etrically opposite to the trigger particle [8]. This show s that the energy of the fast parton originally emitted in the direction opposite to the trigger particle is not lost, but severely degraded by interactions with the medium . As the impact parameter of the collisions decreases, the average m om entum of the particles em itted opposite to the trigger particle approaches the mean value associated with all soft hadrons, i.e. the hp_T i of the thermalized medium [8]. This suggests that the energy lost by the fast parton has been largely therm alized. Nevertheless, this energy is deposited locally along the fast parton's trajectory, leading to local energy density inhom ogeneities which, if therm alized, should in turn evolve hydrodynam ically. This would modify the usual hydrodynam ic expansion of the collision reball as observed in the overwhelm ing number of soft collision events where no high- p_T partons are created.

Since the fast parton m oves at supersonic speed, it was suggested in Ref. [1] that a M ach shock (\sonic boom ") should develop, resulting in conical ow and preferred particle emission at a speci c angle away from the direction of the fast parton which lost its energy. This M ach angle is sensitive to the m edium 's speed of sound c_s and thus o ers the possibility to measure one of its key properties. A recent analysis by the PHEN IX Collaboration [2] of azim uthal di-hadron correlations in 200 A G eV Au+Au collisions revealed structures in the angular distribution which m ight be suggestive of conical ow.

The idea of M ach shock waves travelling through com pressed nuclear matter was rst advocated 30 years ago [9, 10], but RHIC collisions for the rst time exhibit [7] the kind of ideal uid behaviour which might make an extraction of the speed of sound conceivable. An alternate scenario, in which the color wake eld generated by the fast colored parton travelling through a quark-gluon plasm a accelerates soft colored plasm a particles in the direction perpendicular to the wake front [11, 12], leads to an emission pattern which is sensitive to the propagation of plasm a rather than sound waves [12]. In a strongly coupled plasm a with overdam ped plasm a oscillations, which seem s to be the preferred interpretation of RHIC data [13, 14], the wake eld scenario should reduce to the hydrodynam ic M ach cone picture. W e here study the dynam ical consequences of the latter, going beyond the discussion of linearized hydrodynam ic equations in a static background o ered in [1].

W e assume that just before hydrodynam ics become applicable, a pair of high- p_T partons is produced near the surface of the reball. One of them moves outward and escapes, forming the trigger jet, while the other enters into the reballalong, say, the x direction. The reball is expanding and cooling. The ingoing parton travels at

the speed of light and loses energy in the reball which therm alizes and acts as a source of energy and momentum for the reball medium. We model this medium as an ideal uid with vanishing net baryon density. Its dynam ics is controlled by the energy-momentum conservation equations

$$Q T = J ;$$
 (1)

where the energy-m om entum tensor has the ideal uid form T = ("+p)u u pg, with energy density " and pressure p being related by the equation of state (EOS) $p = p("), u = (1;v_x;v_y;0)$ is the uid 4-velocity, and the source current J is given by

$$J(x) = J(x) 1; 1;0;0;$$
 (2)

$$J(x) = \frac{dE}{dx}(x) \frac{dx_{jet}}{dt}^{3}(r r_{jet}(t)): \qquad (3)$$

M assless partons have light-like 4-m om entum, so the current J describing the 4-m om entum lost and deposited in the medium by the fast parton is taken to be light-like, too. r_{jet} (t) is the trajectory of the jet moving with speed jdx_{jet}=dtj= c. $\frac{dE}{dx}$ (x) is the energy loss rate of the parton as it moves through the liquid. It depends on the

uid's local rest fram e particle density. Taking guidance from the phenom enological analysis of parton energy loss observed in Au + Au collisions at RH IC [15] we take

$$\frac{dE}{dx} = \frac{s(x)}{s_0} \frac{dE}{dx}_0$$
(4)

where s(x) is the local entropy density without the jet. Them easured suppression of high- $p_{\rm T}$ particle production in Au+ Au collisions at RHIC was shown to be consistent with a parton energy loss of $\frac{dE}{dx}_{0} = 14 \, {\rm GeV} / {\rm fm}$ at a reference entropy density of $s_{0} = 140 \, {\rm fm}^{-3}$ [15]. For com parison, we also perform simulations with ten times larger energy loss, $\frac{dE}{dx}_{0} = 140 \, {\rm GeV} / {\rm fm}$.

For the hydrodynam ic evolution we use AZHYDRO [7, 16], the only publicly available relativistic hydrodynam ic code for anisotropic transverse expansion. This algorithm is formulated in (;x;y;) coordinates, where $p = \frac{p}{t^2 - z^2}$ is the longitudinal propertime, $= \frac{1}{2} \ln \frac{t+z}{t-z}$ is space-time rapidity, and $r_2 = (x;y)$ denes the plane transverse to the beam direction z. AZHYDRO employs longitudinal boost invariance along z but this is violated by the source term (3). We therefore modify the latter by replacing the -function in (3) by

³ (r r_{jet} (t)) !
$$\frac{1}{2}$$
 (x x_{jet} ()) (y y_{et} ())
! $\frac{1}{2} \frac{e^{(r_{?} r_{?,jet}())^{2} = (2^{-2})}}{2^{-2}}$ (5)

with = 0.35 fm. Intuitively, this replaces the \needle" (jet) pushing through the medium at one point by a

 $\$ while " cutting the medium along its entire length along the beam direction. The resulting $\$ wedge ow " is expected to have a stronger signal in the azim uthal particle distribution dN =d than $\$ on ical ow " induced by a single parton, since in the latter case one perform s an im plicit -average when sum m ing over all directions of the cone norm alvector. W hile a com plete study of this would require a full (3+1)-dim ensional hydrodynam ic calculation, the present boost-invariant sim ulation should give a robust upper lim it for the expected angular signatures. We show that the angular structures predicted from wedge ow are too weak to explain the experim entally observed -m odulation [2].

Them odi ed hydrodynam ic equations in (;x;y;) co-ordinates read [16]

- $e_{x} \mapsto e_{x} (v_{x} \perp) \mapsto e_{y} (v_{y} \perp) = e_{x} p = 0, \quad (7)$

$$(d T^{\prime} T^{\prime} + (d_{x} (v_{x} T^{\prime}) + (d_{y} (v_{y} T^{\prime})) = (d_{y} p;$$

$$(8)$$

where T = T, $v_i = T^{i} = T$, p = p, and J = J.

To simulate centralAu+Au collisions at RHIC, we use the standard initialization described in [7] and provided in the downloaded AZHYDRO input le [16], corresponding to a peak initial energy density of " $_0$ = 30G eV/fm 3 at $_0$ = 0.6 fm/c. We use the equation of state EOS-Q described in [7, 16] incorporating a rst order phase transition and hadronic chemical freeze-out at a critical tem – perature $T_{\rm c}$ = 164M eV. The hadronic sector of EOS-Q is soft with a squared speed of sound $c_{\rm s}^2$ 0.15.

In our study the quenching jet starts from $x_{jet} = 6.4$ fm at $_0 = 0.6$ fm, moving left towards the center with constant speed $v_{jet} = c$. For an upper limit on conical ow e ects, the fast parton is assumed to have su cient initial energy to emerge on the other side of the reball. To simulate cases where the fast parton has insuicient energy to fully traverse the medium we have also done simulations where the parton loses all its energy within an (arbitrarily chosen) distance of 6.4 fm. We further compared with a run where the parton moved at (constant) subsonic speed ($v_{jet} = 0.2 c < c_s$).

The resulting evolution of the energy density of the QGP uid is shown in Figs. 1 and 2. The left column of Fig. 1 shows results for a phenom enologically acceptable value $\frac{dE}{dx_0} = 14 \text{ GeV} / \text{fm}$ [15] for the reference parton energy loss whereas in all other columns we use a ten times larger energy loss. The width of the Gaussian source (see Eq. (5)) is = 0.7 fm. In the left column of Fig. 1 the e ects of the energy deposition from the fast parton are hardly visible. Only for a much larger energy loss (right column) we recognize a clear conical ow pattern. The accumulating wave fronts from the expanding energy density waves build up a \sonic boom " shock front which creates a M ach cone. The right columns in Figs. 1 and 2 show that the cone norm alvector form san angle M with the direction of the quenching jet that is qualitatively



FIG.1: Contours of constant local energy density in the x-y plane at three di erent times, = 4.6; 8.6, and 12.6 fm /c. In each case the position of the fast parton, along with the integrated energy loss E = J(x)dxdyd up to this point, is indicated at the top of the gure. Diagrams (a)-(c) in the left column were calculated with a reference energy loss dE = dx j = 14 G eV / fm, those in the right column (panels (d)-(f)) with a 10 times larger value.

consistent with expectations from the theoretical relation $\cos_{M} = c_{s} = v_{jet}$. However, this angle is not sharply de ned since the cone surface curves due to inhom ogene-ity and radial expansion of the underlying medium. This di ers from the static hom ogeneous case [1].

When the parton travels at a subsonic speed $v_{jet} = 0.2 \text{ c} < c_s$ (left column in Fig.2), it doesn't get very far before the reball freezes out due to longitudinal expansion. In this case one only observes an accumulation of energy around the parton but no evidence of M ach cone formation. When the parton travels with $v_{jet} = c$ but looses all its energy after 6.4 fm before fully traversing the reball (right column in Fig.2), the reball evolution beyond = 7 fm is not a ected by the fast parton directly but only indirectly through the propagation of earlier deposited energy. Still, Figs. 1f and 2f show that the late time evolution of the reball is quite similar in both cases, dem onstrating that energy deposition by the



FIG.2: A s in Fig.1 but for a parton m oving at subsonic speed $v_{jet} = 0.2 c$ (left colum n) and for a fast parton ($v_{jet} = c$) which loses all its energy within the rst 6.4 fm (right colum n).

fast parton during the late reball stages is small, due to dilution of the matter, and can alm ost be neglected.

Tests with di erent values for the width of the Gaussian pro le in Eq. (5) for the deposited energy show that the cone angle gets better de ned for sm aller source size . Note that the quenching jet destroys the azim uthal sym m etry of the initial energy density distribution but leaves the azim uthally sym m etric energy contours to the left of the jet una ected.

W e close with a discussion of observable consequences of conical ow. One expects [1] azim uthally an isotropic particle em ission, peaking at angles = M relative to the trigger jet where M is the Mach angle. Using the standard Cooper-Frye prescription, we have com puted the angular distribution of directly emitted pions at a freeze-out tem perature $T_{fo} = 100 \text{ MeV}$ [7]. Figure 3 shows the azim uthal distribution of from a variety of di erent simulations. [The -independent constant =dyd)_{no jet} = 27 from central collisions without (dN jets has been subtracted.] For $dE = dx_{1} = 14 \text{ GeV} / \text{fm}$ the azim uthalm odulation is very sm all. In none of the cases studied we nd peaks at the predicted M ach angle with an associated dip in the direction of the quenched jet at



FIG. 3: A zim uthal distribution dN =dyd of negative pions per unit rapidity. In the upper panel we integrate over all p_T while the lower panel shows only pions with $p_T > 1 \text{ GeV}/\text{c}$. D i erent sym bols refer to di erent param eters as indicated. For better visibility the -independent rate in the absence of the quenching jet has been subtracted. Filled sym bols show the realistic case dE =dx j = 14 G eV /fm, enhanced by a factor 10 for visibility.

= . One rather sees a peak at = , broadened by shoulders on both sides which turn into small peaks at (relative to the quenching jet) backward angles if them all sm earing is reduced by considering only high- p_T pions. The peak at = is absent when the parton loses all its energy halfway through the reball or is too slow to get to the other side before freeze-out, suggesting that it re-

ects the directed m om entum im parted on the m edium by the fast parton. It is slightly more accentuated for smaller and higher p_T . We also found that the width of the shoulders is almost independent of the speed of sound of the medium and can not be used to diagnose the sti ness of its equation of state. The shoulders exist even for subsonic parton propagation ($v_{iet} = 0.2 c$, upright triangles), showing that otherm echanisms (such as backsplash from the hard parton hitting the uid and a generalbias of the energy deposition towards the right side of the reball due to the higher density of the medium at early times) have a strong in uence on the angular distribution of the emitted particles which interferes with the position of the M ach peaks. The shoulder resulting from this combination of e ects is much broader than the angular structures seen in the data [2]. The absence of a clear dip at = in our simulations is all the more troubling since it should have been stronger for the \wedge

ow "studied here than for real \conical" ow .

O ur calculation does not average over the initial production points of the trigger particle, i.e. it ignores that in m ost cases its quenching partner does not travel right through the middle of the reball cylinder, but traverses it sem i-tangentially. This should further decrease the prom inence of the shoulders in dN =d . We conclude that conical ow may be able to explain the broadening of the away-side peak in the hadron angular correlation function around = pointed out by the STAR Collaboration [8], but is unlikely to be responsible for the relatively sharp structures near = 1 seen by PHEN IX [2]. This conclusion extends to other conical ow phenom ena, such as those generated by color wake elds [11, 12].

This work was supported by the U S.D epartm ent of Energy under contract DE-FG 02-01ER 41190.

Correspond to akc@veccalemet.in

- J.Casalderrey-Solana, E.V. Shuryak and D. Teaney, J. Phys.Conf.Ser.27,22 (2005) [hep-ph/0411315].
- [2] S. S. Adler et al. [PHENIX Collaboration], nuclex/0507004.
- [3] BRAHM S Collaboration, I.A reene et al., Nucl. Phys. A 757,1 (2005); PHOBOS Collaboration, B.B.Back et al., ibid. 757, 28 (2005); STAR Collaboration, J.A dam s et al., ibid.757,102 (2005); PHENIX Collaboration, K.Adcox et al., ibid.757, 184 (2005).
- [4] STAR Collaboration, C. Adler et al., Phys. Rev. Lett. 90, 082302 (2003); STAR Collaboration, J. Adams et al., Phys. Rev. Lett. 91, 072304 (2003); and Phys. Rev. Lett. 93, 252301 (2004).
- [5] PHENIX Collaboration, J. Rak et al., J. Phys. G 30, S1309 (2004).
- [6] M.Gyulassy, I.V itev, X.-N.W ang, and B.-W. Zhang, in Quark-Gluon Plasm a 3, edited by R.C.Hwa and X.-N. W ang (W orld Scientic, Singapore, 2004), p.123.
- [7] P.F.Kolb and U.Heinz, in Ref. [6], p. 634.
- [8] STAR Collaboration, J. Adam s et al., Phys. Rev. Lett. 95, 152301 (2005).
- [9] G.F.Chapline, M.H.Johnson, E.Teller and M.S.W eiss, Phys. Rev. D 8 (1973) 4302.
- [10] W .Scheid, H.M uller, and W .G reiner, Phys. Rev. Lett. 32, 741 (1974).
- [11] H. Stocker, Nucl. Phys. A 750, 121 (2005).
- [12] J.Ruppert and B.Muller, Phys.Lett.B 618, 123 (2005).
- [13] U.Heinz and P.F.Kolb, Nucl. Phys. A 702, 269 (2002).
- [14] M.Gyulassy and L.D.McLerran, Nucl. Phys.A 750, 30 (2205); E.Shuryak, Nucl. Phys.A 750, 64 (2005).
- [L5] X.N.W ang, Phys.Rev.C 70,031901 (2004), and private communication.
- [16] P.F.Kolb, J.Sollfrank, and U.Heinz, Phys.Rev.C 62, 054909 (2000); P.F.Kolb and R.Rapp, Phys.Rev.C 67, 044903 (2003). The code can be downloaded from URL http://nt3.phys.colum.bia.edu/people/m.olnard/OSCAR/
- [17] P.Amold, G.D.Moore and L.G.Ya e, JHEP 0112, 009 (2001)