

Study on the One-Proton Halo Structure in ^{23}Al

FANG De-Qing^{1, *}, MA Chun-wang^{1,2}, MA Yu-Gang¹, CAI Xiang-Zhou¹, CHEN Jin-Gen^{1,2}, CHEN Jin-hui^{1,2}, GUO Wei¹, TIAN Wen-Dong¹, WANG Kun^{1,2}, WEI Yi-Bin^{1,2}, YAN Ting-Zhi^{1,2}, ZHONG Chen¹, ZUO Jia-xu^{1,2}, and SHEN Wen-Qing¹

¹Shanghai Institute of Applied Physics, Chinese Academy of Sciences, P. O. Box 800-204, Shanghai 201800 and

²Graduate school of Chinese Academy of Sciences

(Dated: April 16, 2024)

The Glauber theory has been used to investigate the reaction cross section of proton-rich nucleus ^{23}Al . A core plus a proton structure is assumed for ^{23}Al . HO-type density distribution is used for the core while the density distribution for the valence proton is calculated by solving the eigenvalue problem of Woods-Saxon potential. The transparency function in an analytical expression is obtained adopting multi-Gaussian expansion for the density distribution. Coulomb correction and finite-range interaction are introduced. This modified Glauber model is apt for halo nuclei. A dominant s-wave is suggested for the last proton in ^{23}Al from our analysis which is possible in the RMF calculation.

PACS numbers: 25.60.Dz, 24.10.-i

Interest in reaction cross section (σ_R) has grown over the last decades with the development of radioactive ion beams (RIBs). σ_R for light exotic nuclei has been studied extensively both theoretically and experimentally. Up to now much effort has been devoted to the study of neutron halo structure, such as the discovery of neutron halo nuclei ^6He , ^{11}Li , $^{11,14}\text{Be}$ [1] and the recent work on ^{19}C [2, 3]. At the same time, some work has been performed on proton halo nuclei, e.g. ^8B and ^{17}Ne . Due to Coulomb effect, the formation of proton halo is more difficult and complicated compared to neutron halo structure. And discrepancies were found in proton halo nuclei, especially on ^8B [4, 5, 6, 7]. Recently the σ_R was measured for the proton halo candidate ^{23}Al , whose separation energy of the last proton is very small (0.125 MeV) [8]. Enhancement in σ_R was observed for ^{23}Al compared to its neighbors [9, 10]. According to the shell model, the last proton in ^{23}Al is in the $1d_{5/2}$ orbit. But the RMF calculation shows that it is possible to have the inversion of the $1s_{1/2}$ and $1d_{5/2}$ orbit in ^{23}Al [11]. Thus it is very interesting and important to know which orbit the last proton occupies.

Several methods are available to study the total reaction cross section, i.e. the multi-step scattering theory of Glauber [12], the transport model method of Ma et al. [13] and the semi-empirical formula of Kox et al. [14] and Shen et al. [15] etc. For the first one, the optical limit approximation of the Glauber approach is the most common used method [12]. In the high energy domain, the Glauber model is based on the individual nucleon-nucleon collisions along the classical straight-line trajectory in the overlap volume of the colliding nuclei [12, 16].

This model has been extended to low energies by taking into account the Coulomb distortion of the eikonal trajectory [17]. Recently, it has been found that the Glauber model always underestimates σ_R by 10-50% at intermediate energies, if one assumes HO-type nucleon density distributions and determines the width parameter by reproducing the interaction cross section at relativistic energies [18]. To improve the calculation of the Glauber model at low energies, the finite-range interaction was considered and its energy dependence was investigated by fitting the σ_R of $^{12}\text{C} + ^{12}\text{C}$ from low to high energies [19]. And also the few-body Glauber model was widely used to study nuclei with halo structure, which assumes core plus one-(two)-nucleon structure [20]. However, all these calculations are done by solving a multi-dimensional numerical integration or Monte-Carlo simulation. To simplify the calculation, the Glauber model in an analytical form has been obtained if we use Gaussian function for density distributions [17]. But this method can only be used to describe stable nuclei. In order to study nuclear reaction involving exotic nuclei, the separation energy dependent diffuseness was introduced into this method [21]. Here we present a more general analytical form by expanding the density distributions of projectile and target using multi-Gaussian functions, which can be used to calculate σ_R for both stable nuclei and nuclei with halo structure.

To begin with, we describe the formulation of σ_R in the framework of Glauber model in the optical limit approximation. If we distinguish neutrons and protons in the projectile and target, the reaction cross section can be expressed as

$$\sigma_R = 2 \int_0^{\infty} b [1 - T(b)] db \quad (1)$$

where $T(b)$ is the transparency function at impact parameter b

$$T(b) = \exp \left[-i \sum_{i=1}^Z \sum_{j=1}^Z \int_{-\infty}^{\infty} f(r_i, r_j) dr \right] \quad (2)$$

Supported by the Major State Basic Research Development Program in China Under Contract No. G2000774004, the National Natural Science Foundation of China (NNSFC) under Grant No 10405032, 10328259 and 10135030 and the Phosphor program in Shanghai under contract No. 03QA14066.

*Corresponding author. E-mail address: fangdq@sinr.ac.cn

$$\rho_{ij}(r) = \int_{-b}^b \int_{-b}^b \rho_{ij}^z(r) \rho_{ij}^z(r') dr' \quad (2)$$

here $f(r)$ is the finite-range function

$$f(r) = \frac{1}{2} \exp\left(-\frac{r^2}{2}\right) \quad (3)$$

with ρ_0 being the finite-range parameter. ρ_{ij} is the nucleon-nucleon collision cross section. ρ_{ij}^z is the neutron (proton) density distribution in the projectile (target) with the z axis integrated [17]. With the input density distribution of the projectile and target, the reaction cross section can be obtained after the multidimensional numerical integration.

If we assume Gaussian distributions for the density of projectile and target, which is suitable for light stable nuclei. Then we can obtain an analytical transparency function [17]. To extend this method for both stable nuclei and halo nuclei with a long tail in the density distribution, we express the density distribution by N Gaussian functions

$$\rho_{ij}(r) = \sum_{k=1}^N \rho_{ij}^k(0) \exp\left(-\frac{r^2}{a_{ij}^k}\right) \quad (i=P, T; j=n, p) \quad (4)$$

where $\rho_{ij}^k(0)$ and a_{ij}^k are the amplitude and width parameters of the Gaussian functions. Then the transparency function can be written in an analytical form similarly [17, 21]

$$T(b) = \exp\left[-\sum_{i=P, T} \sum_{j=n, p} \sum_{k=1}^N \rho_{ij}^k(0) \exp\left(-\frac{b^2}{a_{ij}^k}\right)\right] \quad (5)$$

where b^0 is the impact parameter after considering Coulomb correction

$$b^0 = \frac{b^2}{1 - \frac{b^2}{V_c E_{CM}}} = \frac{b^2}{1 - \frac{1.44 Z_P Z_T}{R_{int} E_{CM}}} \quad (6)$$

here Z_P (Z_T) refers to the charge number of the projectile (target), R_{int} is the interaction radius and E_{CM} is the incident energy in the center of mass frame [17].

To investigate the R of ^{23}Al , we assume a core (^{22}Mg) plus one proton structure since the weak bond between the last proton and the core. HO-type distribution is used for the core density. The density distribution for the valence proton was calculated by solving the eigenvalue problem of Woods-Saxon potential [22]

$$\frac{d^2 R(r)}{dr^2} + \frac{2}{h^2} [E - U(r)] \frac{1}{R(r)} = 0 \quad (7)$$

where $f(r) = [1 + \exp(\frac{r-R}{a})]^{-1}$ with $R = r_0 A_c^{1/3}$ ($V_{ls} = 17\text{ MeV}$). V_0 is the depth of potential, V_{Coul} is the Coulomb potential. In the calculation the diffuseness (a) and radius parameter (r_0) were chosen to be 0.67 fm and 1.27 fm [23].

First we assume HO-type distribution for the ^{12}C target and determine the HO width parameter by reproducing the interaction cross section of $^{12}\text{C} + ^{12}\text{C}$ at relativistic energy. Then the R for $^{12}\text{C} + ^{12}\text{C}$ at different energies are calculated and compared them with experimental data. In the calculation, the finite-range parameter was taken from Ref. [19].

$$\frac{R}{\text{fm}} = 2.2 \exp\left(\frac{E}{106.679}\right) + 0.089 \quad (8)$$

where E is the incident energy in the laboratory frame in MeV. With the consideration of the energy dependent finite-range interaction, the underestimation at low energies in the zero-range Glauber model was removed as described in Ref. [19]

TABLE I: The reaction cross sections for ^{22}Mg and ^{23}Al with carbon target.

Nuclei	energy (AMeV)	R (mb)	reference
^{22}Mg	33.4	1531 ± 125	[9]
^{23}Al	35.9	1892 ± 145	[9]

Then we assume HO distribution for the ^{22}Mg core, and adjust the HO width parameter to fit the measured R at 33.4 AMeV as given in Table. I.




FIG. 1: Density distributions of ^{22}Mg and ^{23}Al . The dotted line is the density distribution of ^{22}Mg . The solid and dashed line are the density distribution of ^{23}Al in s- and d-wave, respectively. For comparison, the density distribution of ^{23}Al in the s-wave calculated by multi-Gaussian expansion (Eq.(4)) is also plotted as the dotted-dashed line. It is almost overlapped with the original density distribution.




FIG. 4: Density distributions of ^{23}Al . The solid and dashed line are the density distribution of ^{23}Al in s- and d-wave, respectively. The thick line is the extracted density distribution assuming HO plus Yukawa distribution. The shaded area is the error from the experimental R data.

where r_c is the crossing point of these two functions, the factor Y is to keep the equality of the two distributions at r_c . The width parameter of the HO-type core is fitted to the R of ^{22}Mg . Then the parameters Y and r_c are fitted to the R of ^{23}Al given in Table. I with an addi-

tional normalization condition $\int_0^R \rho(r) dr = Z$ and Z is the charge number of ^{23}Al .

In Fig.4, the extracted density distribution was compared with the calculated density for ^{23}Al in s- and d-wave. The extracted density is closer to s-wave than d-wave. It demonstrates a long tail in the density distribution and supports the conclusion that the s-wave is dominant for the last proton in ^{23}Al .

In summary, the Glauber theory has been used to investigate the reaction cross section. Adopting multi-Gaussian expansion for the density distribution of projectile and target, an analytical form for the transparency function was deduced. The energy dependent finite-range interaction and Coulomb correction were considered. It has been pointed out that this modified Glauber model is suitable for studying the R of both stable nuclei and nuclei with exotic structure. We used this model to study the reaction cross section of proton-rich nucleus ^{23}Al . A core plus proton structure was assumed for ^{23}Al . HO-type density distribution was used for the core while the density distribution for the valence proton was calculated by solving the eigenvalue problem of Woods-Saxon potential. The s- and d-wave mixing configuration was studied. Density distribution was extracted for ^{23}Al assuming HO plus Yukawa-square tail. A dominant s-wave was suggested in our analysis which indicates a proton halo structure in ^{23}Al .

-
- [1] Tanihata I et al. 1985 Phys. Rev. Lett. 55 2676; Tanihata I et al. 1985 Phys. Lett. B 160 380; Tanihata I et al. 1992 Phys. Lett. B 287 307
 - [2] Bazin D et al. 1995 Phys. Rev. Lett. 74 3569; Bazin D et al. 1998 Phys. Rev. C 57 2156
 - [3] Ozawa A et al. 2002 Eur. Phys. J. A 13 163
 - [4] Warner R E et al. 1995 Phys. Rev. C 52 R1166
 - [5] Negoita F, Borcea C and Carstoiu F 1996 Phys. Rev. C 54 1787
 - [6] Fukuda M et al. 1999 Nucl. Phys. A 656 209
 - [7] Obuti M M et al. 1996 Nucl. Phys. A 609 74
 - [8] Audi G and Wapstra A H 1993 Nucl. Phys. A 565 66
 - [9] Cai X Z et al. 2002 Phys. Rev. C 65 024610
 - [10] Zhang H Y et al. 2002 Nucl. Phys. A 707 303
 - [11] Chen J G et al. 2004 Chin. Phys. Lett. 21 2140
 - [12] Glauber R J 1959 Lectures in theoretical physics. Vol. 1 (Interscience, New York, 1959)
 - [13] Ma Y G et al. 1993 Phys. Lett. B 302 386; Ma Y G et al. 1993 Phys. Rev. C 49 850.
 - [14] Kox S et al. 1987 Phys. Rev. C 35 1678.
 - [15] Shen W Q et al. 1989 Nucl. Phys. A 491 130.
 - [16] Karol P J 1975 Phys. Rev. C 11 1203
 - [17] Charagi S K and Gupta S K 1990 Phys. Rev. C 41 1610
 - [18] Ozawa A et al. 1996 Nucl. Phys. A 608 63
 - [19] Zheng T et al. 2002 Nucl. Phys. A 709 103
 - [20] Ogawa Y, Suzuki Y and Yabana K 1994 Nucl. Phys. A 571 784
 - [21] Feng J et al. 1993 Phys. Lett. B 305 9
 - [22] Speth J et al. 1977 Phys. Rep. 33 127
 - [23] Yamaguchi T et al. 2003 Nucl. Phys. A 724 3
 - [24] Zhao Y L, Ma Z Y, Chen B Q and Shen W Q 2003 Chin. Phys. Lett. 20 53