Form ation of K aonic A tom s and K aonic N uclei in In-F light (K ;p) R eactions

Junko Yamagata¹, Hideko Nagahiro², Yuko O kumura^{1;)} and Satoru Hirenzaki¹

¹D epartm ent of P hysics, N ara W om en's U niversity, N ara 630-8506, Japan ²Research C enter for N uclear P hysics (R C N P), O saka U niversity, Ibaraki, O saka 567-0047, Japan

(Received March 11, 2005)

We theoretically study kaonic atom and kaonic nucleus form ation in in- ight (K ;p) reactions for C, O, Si and C a targets. Deeply bound kaonic atom swere previously predicted to exist as quasi-stable states, and it is expected that they can be observed using certain well-suited experimental methods. K aonic nuclear states have also been predicted to exist with large decay widths. We evaluate the form ation cross sections of kaonic atom s and kaonic nuclei using an elective number approach. We show that indications of kaonic bound states can be observed in the outgoing proton energy spectra.

x1. Introduction

K aonic atom s and kaonic nuclei carry in portant information concerning the K -nucleon interaction in nuclear medium. This information is very important to determ ine the constraints on kaon condensation in high density matter. The properties of kaons in nuclei are strongly in uenced by the change undergone by (1405) in nuclear medium, because (1405) is a resonance state just below the kaon-nucleon threshold. In fact, there are studies of kaonic atom s carried out by modifying the properties of (1405) in nuclear medium.^{1);2);3)} These works reproduce the properties of kaonic atom s very well, which come out to be as good as the phenom enological study of Batty.⁴⁾

Recently, there have been signi cant developments in the description of hadron properties in terms of the SU (3) chiral Lagrangian. The unitarization of the chiral Lagrangian allows the interpretation of the (1405) resonance state as a baryon-meson coupled system $.^{5);6}$ Subsequently, the properties of (1405) in nuclearmedium using the SU (3) chiral unitary model were also investigated by W aas et al.⁷⁾ Lutz,⁸⁾ R am os and 0 set,⁹⁾ and C ieply et al.¹⁰⁾ Allof these works considered the Paulie ect on the intermediate nucleons. In addition, in Ref. 8), the self-energy of the kaon in the intermediate states is considered, and in Ref. 9), the self-energies of the pions and baryons are also taken into account. These approaches lead to a kaon self-energy in nuclear medium that can be tested with kaonic atoms and kaonic nuclei. There are also K potential studies based on meson-exchange Julich K N interaction.^{11);12)}

In a previous work,¹³⁾ we adopted the scattering amplitude in nuclear medium calculated by Ram os and $0 \sec^{9)}$ for studies of kaonic atom s, and demonstrated

⁾ P resent address: R itto H igh School, R itto, Shiga, Japan

the ability to reproduce the existing kaonic atom data as accurately as the optical potential studied by Batty.⁴⁾ We then calculated the deeply bound kaonic atoms for ¹⁶O and ⁴⁰Ca, which have narrow widths and are believed to be observable with well-suited experimental methods.^{14);15)} We also obtained very deep kaonic nuclear states, which have large decay widths, of the order of several tens of MeV. The (K ;) reaction was studied for the form ation of the deeply bound kaonic atom is states,¹³⁾ which could not be observed with kaonic X-ray spectroscopy, using the form ulation developed in Ref. 16) for the form ation of deeply bound pionic atom s with the (;) reaction.

A nother very in portant developm ent in recent years is that in the study of kaonic nuclear states, which are kaon-nucleus bound systems determined mainly by the strong interaction. Experimental studies of the kaonic nuclear states using in-ight (K;N) reactions were proposed and performed by K ishim oto and his collaborators.^{17);18)} Experiments employing stopped (K;N) reactions were carried out by Iw asaki, T. Suzuki and their collaborators and reported in Refs. 19) and 20). In these experiments, they found some indications of the existence of kaonic nuclear states related to these experimental activities.²¹⁾ It should be noted that these theoretical studies predict the possible existence of ultra-high density states in kaonic nuclear system s.^{21);22)}

In this paper, we study in-ight (K ;p) reactions system atically with regard to their role in populating deeply bound kaonic states and the observation of their properties in experiments. We have found the usefulness of direct reactions in the form ation of deeply bound pionic atoms using the $(d_i^{3}He)$ reactions.^{23);24);25);26)} However, in the present case, K⁺ must be produced in addition in this $(d_i^{3}He)$ reaction, and there would be a large momentum m ism atch. For this reason, the (K ;) reaction was considered rst in Ref.13). Here we theoretically study another reaction, (K ;p), and present system atic results that elucidate the experimental feasibility of the reaction. The (K ;p) reaction was proposed in Refs.15) and 17). However, realistic spectra have not yet been calculated. We calculate the spectra theoretically using the approach of Ref.24) for the deeply bound pionic atom form ation reaction. We believe that this theoretical evaluation will be interesting and in portant for studies of kaon properties in nuclear medium.

In x2, we describe the theoretical model of the structure of kaon-nucleus bound system s and present the num erical results. The theoretical form alism and num erical results for the (K ; p) reactions are discussed in x3. We give sum m ary in x4.

x2. Structure of kaonic atom s and kaonic nuclei

2.1. Form alism

 ${\tt W}$ e study the properties of kaonic bound system s by solving the K ${\tt kin}\mbox{-}{\tt G}$ ordon equation

$$[r^{2} + {}^{2} + 2 V_{opt}(r)] (r) = [E V_{coul}(r)]^{2} (r) : (2.1)$$

Here, is the kaon-nucleus reduced m ass and $V_{coul}(r)$ is the C oulom b potential with a nite nuclear size:

$$V_{coul}(\mathbf{r}) = e^{2} \frac{p(\mathbf{r}^{0})}{j\mathbf{r} - \mathbf{r}^{0}j} d^{3}\mathbf{r}^{0}; \qquad (2.2)$$

where $_{\rm p}$ (r) is the proton density distribution. We employ the empirical W oods-Saxon form for the density and keep the shapes of the neutron and proton density distributions xed as

$$(\mathbf{r}) = {}_{n}(\mathbf{r}) + {}_{p}(\mathbf{r}) = \frac{0}{1 + \exp[(\mathbf{r} - \mathbf{R}) = \mathbf{a}]};$$
 (2.3)

where we use $R = 1:18A^{1=3}$ 0:48 [fm] and a = 0:5 [fm] with A, the nuclear mass number. It is noticed that the point nucleon density distributions are deduced from in Eq. (2.3) by using the same prescription described in Sect. 4 in Ref. 29) and

are used to evaluate the kaon-nucleus optical potential.

The kaon-nucleus optical potential V_{opt} is given by

$$2 V_{opt}(r) = 4 a_e(r)(r);$$
 (2.4)

where a_e () is a density dependent e ective scattering length and = $1 + m_K = M_N$. In this paper, we use two kinds of e ective scattering lengths, that obtained with the chiral unitary approach ⁹ and that obtained with a phenom enological t.²⁷ Here, we do not introduce any energy dependence for the e ective scattering lengths, and we use the scattering lengths at the K N threshold energy. The elective scattering length a_e of the chiral unitary approach is described in Ref. 13) in detail. It is dened by the kaon self-energy in nuclear matter, with the local density approximation. The form of a_e obtained in a phenom enological t is one of the results reported in Ref. 27), and it is parameterized as

$$a_e$$
 () = (0:15 + 0:62i) + (1:66 0:04i) (= 0)^{0:24} [fm]: (2.5)

The reason we consider these two potentials is that they provide equivalently good descriptions of the observed kaonic atom data, even though they have very di erent potential depths, as we will see in next subsection. Thus, it should be extrem ely interesting to compare the results obtained with these potentials in the (K ;p) reaction spectra, including the kaonic nuclear region.

W e solve the K lein-G ordon equation num erically, following the method of 0 set and Saleedo.²⁸⁾ The application of this method to pionic atom studies are reported in detail in Ref. 29).

2.2. Num erical results

We show the kaon nucleus potential for the 39 K case in Fig. 1 as an example. Because the realpart of a_e () changes sign at a certain nuclear density in both the chiral unitary and phenom enological models, the kaon nucleus optical potential is attractive, while keeping the repulsive sign for the kaon-nucleon scattering length in free space.

The real part of the scattering length for the phenom enological t depends on the density much more strongly than the results of the chiral unitary model and yields Re a_e ($_0$) = 1:51 [fm]. Hence, as we can see in Fig. 1, the depths of the real optical potentials of these models di er signi cantly. On the other hand, the density dependence of the imaginary part of the phenom enological scattering length is rather at, and its strength is similar to that of the chiral unitary model.



Fig.1. The kaon-nucleus optical potential for ³⁹K as a function of the radial coordinate r. The left and right panels show the real and in aginary part, respectively. The solid line indicates the potential strength of the chiral unitary approach and the dashed line of the phenom enological t.

The calculated energy levels for the atom ic states and nuclear states in 39 K are shown in Fig. 2, where the results of the chiral model and the phenom enological model, Eq. (2.5), are compared. We see that the results obtained with the two potentials are very similar for the atom ic states. We not that the deep atom ic states, such as atom ic 1s in 39 K (still unobserved), appear with narrower widths than the separation between levels and are predicted to be quasi-stable states. Similar results are reported in previous works.^{13);14)} Because several model potentials predict the states experimentally. On the other hand, the predicted binding energies and widths are very stable and alm ost identical for all of the potential models considered here. Hence, it is very di cult to distinguish the theoretical potentials from only the observation of atom ic levels.

In the lower panels of F ig. 2, we also show the energy levels of the deep nuclear kaonic states of 39 K using the chiral unitary model potential and the phenom enological model potential. The deep nuclear states are represented by the solid bars with numbers, which indicate their widths in units of M eV. These nuclear states have extremely large widths in all cases and would not be observed as peak structures in experiments if they indeed do have such large widths. We should, however, mention here that the level structures of these potential models dier signi cantly. In the chiral unitary potential, only two nuclear states are predicted, while eight states are predicted with the phenom enological model. This dierence presents the opportunity to distinguish the theoretical potentials in observations of kaonic nuclear states.

C alculated density distributions of nuclear 1s and 2s and atom ic 1s kaonic states in 39 K are shown in Fig. 3 for the case of the phenom enological optical potential.



Fig. 2. (Upper panel) Energy levels of kaonic atom s of ³⁹K obtained with the optical potentials of the chiral unitary m odel (left) and the phenom enological t (right). The hatched areas indicate the level widths. (Low er panel) Energy levels of kaonic nuclear states of ³⁹K obtained with the optical potentials of the chiral unitary m odel (left) and the phenom enological t (right). The level width is indicated by the num ber appearing at each level, in units of M eV.

It is seen that the wavefunctions of the deep nuclear kaonic states remain almost entirely inside the nuclear radius, which is about 3.5 fm for 39 K. Hence, the widths become extremely large, of the order of 100 MeV. The wavefunctions of the atom ic states are pushed outward by the imaginary part of the strong interaction. It should be noted that the atom ic 1s state corresponds to the 4-th s state in the solutions of the K lein-G ordon (KG) equation, Eq. (2.1). We divided the series of KG solutions into two categories, 'atom ic states' and 'nuclear states', since the properties of these states are very di erent, and there are no ambiguities in this classi cation, as can be seen in Figs. 2 and 3.

We have also calculated the kaon-nucleus binding energies and widths for both atom ic and nuclear states in other nuclei. The obtained results are compiled in Tables I and II for the phenom enological optical potential and for the chiral unitary potential cases, respectively. We selected ¹¹B, ¹⁵N, ²⁷A land ³⁹K nuclei, which appear in the nal states of the form ation reactions for ¹²C, ¹⁶O, ²⁸Si and ⁴⁰C a targets, as described in the next section. In all cases, we found kaonic atom states and kaonic nuclear states. The results for the atom ic states are sim ilar for the two potentials and are known to reproduce existing data reasonably well. On the other hand, certain di erences are found in the energy spectra of kaonic nuclear states, as expected, and they should be investigated experimentally.

We should mention here that the kaonic nuclear 2p state in ²⁷Al described in Table II provides a negative value for the binding energy. This state, however, is interpreted as a bound state, since the sign of the corresponding eigenenergy in the non-relativistic Schrödinger equation is opposite to that of the K lein-G ordon

| N uclear State | ¹¹ B | | ¹⁵ N | | ²⁷ Al | | ³⁹ K | |
|----------------|-----------------|---------------------|-----------------|---------------------|------------------|---------------------|---------------------|----------------|
| (MeV) | в£. | | в£. | | в£. | | ВЕ. | |
| 1s | 132.5 | 183.0 | 155.7 | 205.5 | 190.8 | 239.5 | 206.7 | 253.8 |
| 2s | - | - | 19.0 | 96.2 | 69.8 | 142.3 | 101.3 | 165.6 |
| 3s | - | - | - | - | - | - | 2.1 10 ¹ | 88.8 |
| 2p | 58.4 | 127.0 | 86.9 | 151.1 | 136.3 | 191,2 | 161.7 | 211.3 |
| Зp | - | - | - | _ | 16.5 | 103.3 | 50.2 | 131.6 |
| 3d | - | - | 22.9 | 108.3 | 80.4 | 152.2 | 113.3 | 175.2 |
| 4d | - | - | - | _ | - | _ | 3.9 | 98.7 |
| 4f | - | - | - | _ | 26.2 | 119.5 | 64.3 | 144.9 |
| Atom ic State | | | | | | | | |
| (keV) | | | | | | | | |
| 1s | 192.4 | 40.6 | 338.1 | 68.3 | 844.9 | 243.1 | 1408.0 | 355.4 |
| 2s | 60.5 | 7.3 | 111.8 | 13.3 | 319.0 | 58.3 | 580.6 | 97.6 |
| 3s | 29.2 | 2.5 | 55.1 | 4.6 | 166.8 | 22.3 | 316.9 | 39.9 |
| 4s | 17.1 | 1.1 | 21.7 | 1.1 | 102.3 | 10.8 | 199.4 | 20.0 |
| 2p | 78.5 | 6.2 10 ¹ | 154.6 | 4.0 | 507.5 | 37.4 | 988.7 | 132.7 |
| Зp | 34.9 | 2.2 10 1 | 68.8 | 1.4 | 229.4 | 12.7 | 458.4 | 45.8 |
| 4p | 19.6 | 9.6 10 2 | 38.7 | 6.1 10 1 | 130.5 | 5.7 | 265.1 | 20.8 |
| 3d | 34.9 | $2.3 \ 10^{4}$ | 69.3 | $2.9 10^{-3}$ | 243.0 | 4.2 10 ¹ | 520.9 | 5.0 |
| 4d | 19.6 | 1.4 10 4 | 39.0 | $1.7 \ 10^{-3}$ | 136.6 | 2.5 10 1 | 292.6 | 2.9 |
| 4f | 19.6 | 1.0 10 ⁸ | 38.9 | 4.8 10 ⁷ | 136.5 | $3.2 \ 10^{4}$ | 293.7 | $1.7 \ 10^{2}$ |

Table I. Calculated binding energies and widths of kaon-¹¹B, -¹⁵N, -²⁷Aland -³⁹K systems with the phenom enological optical potential in units of MeV for kaonic nuclear states and in units of keV for kaonic atom states.

Table II. Calculated binding energies and widths of kaon-¹¹B, -¹⁵N, -²⁷Aland -³⁹K systems with the optical potential of the chiral unitary model, in units of MeV for kaonic nuclear states and in units of keV for kaonic atom states.

| N uclear State | ¹¹ B | | ¹⁵ N | | ²⁷ Al | | ³⁹ K | |
|----------------|-----------------|---------------------|-----------------|---------------------|------------------|-----------------|-----------------|----------------|
| (MeV) | В.Е. | | В£. | | В£. | | в£. | |
| 1s | 4.6 | 81.6 | 11.0 | 87.9 | 22.5 | 96.6 | 29.3 | 100.3 |
| 2p | - | _ | - | _ | -1.1 | 79.5 | 9.0 | 87.4 |
| Atom ic State | | | | | | | | |
| (keV) | | | | | | | | |
| 1s | 197.5 | 35.0 | 340.5 | 67.8 | 852.1 | 199.1 | 1422.7 | 337.3 |
| 2s | 61.3 | 62 | 112.3 | 13.2 | 320.4 | 47.6 | 584.3 | 92.5 |
| 3s | 29.5 | 2.1 | 55.3 | 1.6 | 167.2 | 18.2 | 318.3 | 37.8 |
| 4s | 17.3 | 9.3 10 ¹ | 21.7 | 1.1 | 102.5 | 8.8 | 200.1 | 19.0 |
| 2p | 78.4 | 6.3 10 1 | 154.4 | 3.1 | 507.3 | 34.8 | 986.8 | 109.3 |
| Зp | 34.8 | 2.2 10 1 | 68.7 | 1.1 | 229.3 | 11.8 | 457.7 | 37.7 |
| 4p | 19.6 | 9.7 10 2 | 38.7 | 4.8 10 ¹ | 130.5 | 5.3 | 264.8 | 17.1 |
| 3d | 34.9 | $1.6 \ 10^{4}$ | 69.3 | $2.6 \ 10^{-3}$ | 243.0 | 3.4 10 1 | 520.8 | 4.5 |
| 4d | 19.6 | 9.5 10 ⁵ | 39.0 | $1.6 \ 10^{-3}$ | 136.6 | $2.0 \ 10^{-1}$ | 292.6 | 2.6 |
| 4f | 19.6 | 1.0 10 ⁸ | 38.9 | $3.7 \ 10^{-7}$ | 136.5 | 2.8 10 4 | 293.7 | $1.5 \ 10^{2}$ |



Fig. 3. The kaonic bound state density distributions jr (r) f in coordinate space for ³⁹K obtained with the phenom enological optical potential. The solid and dotted curves indicate the distributions of the 1s and 2s states. The dashed curve represents the density of the 4s state and is regarded as a kaonic atom 1s state. The half-density radius of ³⁹K is also shown.

solution, due to the large widths of the nuclear states, as shown below. The binding energies B_{KG} and widths $_{KG}$ of the K lein-G ordon equation, which are tabulated in Tables I and II, are de ned as $E = (B_{KG}) \frac{i}{2}_{KG}$ by the eigenenergy E in Eq. (2.1). The non-relativistic binding energy B_S and width $_S$ of the Schrodinger equation are related to B_{KG} and $_{KG}$ as

$$B_{S} = B_{KG} - \frac{B_{KG}^{2}}{2} + \frac{B_{KG}^{2}}{8};$$
 (2.6)

$$_{\rm S} = _{\rm KG} \quad \frac{{\rm B}_{\rm KG}}{{\rm K}_{\rm G}} \quad (2.7)$$

Thus, in the case of the kaonic 2p nuclear state in 27 Al described in Table II, the non-relativistic binding energies and widths are B_S 0:5M eV and _{S KG}, which indicate that the state is bound. It should be noted that the asymptotic behavior of the wavefunction is determined by B_S.

x3. K aonic atom s and kaonic nuclei form ation in (K ;p) reactions

3.1. Form alism

W e adopt the theoretical model presented in Ref.24) to calculate the form ation cross sections of the kaonic atom s and kaonic nuclei in the (K ; p) reaction. In this model, the em itted proton energy spectra can be written as

$$\frac{d^2}{dE_p d_p} = \frac{d}{d} \int_{K_p! \, pK_p}^{lab} \frac{X}{E_p! \, pK_p} \frac{K}{E_p! \, pK_p} \frac{1}{E_p! \, pK_p} (3.1)$$

where the sum is over all (kaon-particle) (proton-hole) con gurations in the nal kaonic bound states. The di erential cross section for the elementary process of the reaction K + p ! p + K in the laboratory fram e, (d = d) $_{K}^{lab}_{p! pK}$ is evaluated using the K p total elastic cross section data in Ref. 30) by assuming a at angular

distribution in the center-of-m ass frame at each energy. The resonance peak energy is determined by E appearing in Eq. (3.1), which is dened as

$$E = T_{p} (T_{K} S_{p} (j_{p}^{1}) + B_{K}); \qquad (3.2)$$

where T_K is the incident kaon kinetic energy, T_p the emitted proton kinetic energy, and B_K the kaon binding energy in the nal state. The proton separation energy S_p from each single particle level listed in Table III is obtained from the data in Refs. 31), 32), 33), 34), 35), 36). We use the data in Ref. 37) for the separation energies of the proton-hole levels corresponding to the ground states of the daughter nuclei. The widths of the hole states $_p$ are also listed in Table III. These were obtained from the same data sets by assuming the widths of the ground states of the daughter nuclei to be zero because of their stabilities.

Table III. One proton separation energies S_p and widths $_p$ of the hole states of ^{12}C , ^{16}O , ^{28}Si and ^{40}C a deduced from the data given in Ref. 31) for ^{12}C , those given in Refs. 32) and 33) for ^{16}O , those given in Refs. 34) and 35) for ^{28}Si , and those given in Ref 36) for ^{40}C a. The separation energies corresponding to the ground states of the daughter nuclei are taken from Ref. 37). The widths $_p$ indicate FW HM of the Lorentz distribution for ^{16}O and of G aussian distributions for other nuclei. The widths of the ground states of the daughter nuclei are xed to zero, because of their stabilities. For the 1p states in ^{28}Si and ^{40}C a, two levels, $1p_{1=2}$ and $1p_{3=2}$, have not been observed separately, and therefore S_p and $_p$ are set to the same values for both levels.

| single particle | 12 | С | | ¹⁶ O | 28 | Si | 40 | Ca |
|-------------------|------|------|------|---------------------|------|------|------|------|
| states [M eV] | Sp | р | Sp | р | Sp | р | Sp | р |
| 1d ₃₌₂ | | | | | | | 8.3 | 0 |
| $2s_{1=2}$ | | | | | | | 11.5 | 7.7 |
| $1d_{5=2}$ | | | | | 11.6 | 0 | 16.3 | 3.7 |
| 1p ₁₌₂ | | | 12.1 | 0 | 27.5 | 17.0 | 33.2 | 21.6 |
| 1p ₃₌₂ | 16.0 | 0 | 18.4 | 3.1 10 ⁶ | 27.5 | 17.0 | 33,2 | 21.6 |
| $1s_{1=2}$ | 33.9 | 12.1 | 41.1 | 19.0 | 46.5 | 21.0 | 56.3 | 30.6 |

The e ective number N $_{\rm e}$ is de ned as

$$N_{e} = \frac{X_{f}^{2}}{\int M_{m_{s}}^{3}} d^{3}r_{f}(r) = 2m_{s} [l_{k}(r) j_{p}(r)]_{JM_{i}}(r)^{2} : (3.3)$$

The proton and the kaon wavefunctions are denoted by j_p and l_k . We adopt the harmonic oscillator wavefunction for j_p . The spin wave function is denoted by $1=2m_s$, and we take the spin average with respect to m_s , so as to take into account the possible spin directions of the protons in the target nucleus. The functions i and f are the initial and naldistorted waves of the projectile and ejectile, respectively. We use the eikonal approximation and replace f and i by employing the relation

$$_{f}(r)_{i}(r) = \exp(iq r)D(z;b);$$
 (34)

where q is the momentum transfer between the projectile and ejectile, and the distortion factor D (z;b) is de ned as

D (z;b) = exp
$$\frac{1}{2} K_{N} \int_{1}^{Z_{z}} dz^{0} A(z^{0};b) \frac{1}{2} V_{N} \int_{1}^{Z_{1}} dz^{0} A(z^{0};b) : (3.5)$$

Here, the kaon-nucleon and proton-nucleon total cross sections are denoted by $_{KN}$ and $_{pN}$. The functions $_{A}$ (z;b) and $_{A-1}$ (z;b) are the density distributions of the target and daughter nuclei in the beam direction coordinate z with in pact parameter b, respectively.

We calculated the kaonic bound state wavefunctions using the optical potentials obtained with the chiral unitary model¹³⁾ and the phenom enological t,²⁷⁾ as described in x2. In the chiral unitary model, the depth of the attractive potential is only approximately 50 M eV at the center of the nucleus, which is much weaker than the phenom enological potential used in Ref. 27). For the case of the phenom enological potential, there exist kaonic nuclear bound states with very large binding energies, for example, 100 { 200 M eV . For these bound states, the K N system cannot decay into , because of the threshold, and hence, the widths of these states are expected to be narrow er. On the other hand, for chiral unitary potential cases, we do not have narrow nuclear states, like those in Refs. 17) and 21), because the decay phase space for the K N system to the channel is su ciently large, ow ing to the sm aller binding energies. In order to include 'narrow ing e ects' for the widths due to the phase space suppression, we introduce a phase space factor f^{M FG}, de ned in Ref. 38) by M ares, Friedm an, and G al as,

$$f^{M FG}(E) = 0.8 f_1^{M FG}(E) + 0.2 f_2^{M FG}(E); \qquad (3.6)$$

where f_1^{MFG} and f_2^{MFG} are the phase space factors for K N ! decay and K N N ! N, respectively. These factors are de ned as

$$f_{1}^{MFG}(E) = \frac{M_{01}^{3}}{M_{1}^{3}} \frac{M_{1}^{2}(m + m)^{2}M_{1}^{2}(m - m)^{2}}{M_{01}^{2}(m + m)^{2}M_{01}^{2}(m - m)^{2}} (M_{1} - m);$$
(3.7)

and

$$\mathbf{f}_{2}^{M FG} (\mathbf{E}) = \frac{M_{02}^{3}}{M_{2}^{3}} \frac{M_{2}^{2} (\mathfrak{m}_{N} + \mathfrak{m}_{N})^{2} [M_{2}^{2} (\mathfrak{m}_{N} + \mathfrak{m}_{N})^{2}] [M_{02}^{2} (\mathfrak{m}_{N} + \mathfrak{m}_{N})^{2}] [M_{02}^{2} (\mathfrak{m}_{N} - \mathfrak{m}_{N})^{2}]} (M_{2} - \mathfrak{m}_{N} - \mathfrak{m}_{N})^{2}]$$
(3.8)

Here, the branching ratios of m esic decay and non-m esic decay are assumed to be 80% and 20%. The m assess are de ned as M₀₁ = m_K + m_N, M₁ = M₀₁ + E, M₀₂ = m_K + 2m_N, M₂ = M₀₂ + E, and E is the kaon energy de ned as E = T_K T_p S_p, using the same kinem atical variables as in Eq. (3.2). We multiply the energy independent kaonic widths _K by the phase space factor f^{M FG} in order to introduce the energy dependence due to the phase space suppression as

$$_{\rm K} ! _{\rm K} ({\rm E}) = _{\rm K} f^{\rm M FG} ({\rm E}):$$
 (3.9)

3.2. Num erical results

W e present the num erical results for the kaon bound state form ation spectra in this subsection. First, we consider the momentum transfer of the (K ;p) reactions as a function of the incident kaon energy, which is an important guide to determ ine suitable incident energies in order to obtain a large production rate of the bound states. Because we consider both atom ic and nuclear kaon states, we assume four di erent binding energies to calculate the momentum transfer. We consider the forward reactions and the momentum transfer in the laboratory frame, as shown in Fig. 4. As can be seen in the gure, the condition of zero recoil can be satis ed only for atom ic states with $T_{\rm K} = 10$ { 20 MeV. For deeply bound nuclear states, the reaction always requires a certain momentum transfer. However, the incident energy dependence of the momentum transfer is not strong for kaonic nuclear states, as shown in Fig. 4.



Fig. 4. M om entum transfers in the (K ;p) reactions with ⁴⁰C a for targets the form ation of kaon-³⁹K bound system s. The proton separation energy S_p is xed at 8.3 M eV, and the kaon binding energies are assumed to be 0 M eV (solid curve), 50 M eV (dotted curve), 100 M eV (dashed cureve), and 150 M eV (long-dashed curve).

We consider form ation of kaonic atom s and kaonic nuclei separately, because their properties, and hence, the optim alkinem atical conditions for their form ation are expected to be di erent. We extra consider the form ation of atom ic states. Because the binding energies of atom ic states are su ciently sm all, we can safely ignore the phase space e ect for the decay widths considered in Eqs. (3.6) { (3.9). For atom ic states, the obtained wavefunctions and energy spectra are alm ost identical for the chiral unitary and phenom enological potentials. For this reason, we show the results only for chiral unitary potential.

We rst study the energy dependence of the atom ic 1s state form ation rate in order to determ ine the optim al incident energy for the deepest atom ic 1s state observation in the (K ;p) reactions. For this purpose, we show in Fig.5 the energy dependence of the ratio of the calculated e ective num bers of the kaonic atom 1s and 2p states coupled to the $[d_{3=2}^{-1}]$ proton-hole state in ³⁹K. We in that the contribution of the 1s state is significantly larger than that of the 2p state at $T_K = 20 \text{ MeV}$, and we therefore hypothesize that the 1s state can be observed clearly without a large background due to the 2p kaonic state at this energy, where the minimum of the minimum state is state in the minimum state in the state is state in the state in the state can be observed clearly without a large background due to the 2p kaonic state at this energy, where the minimum states is state in the state is state in the state is state in the state in the state in the state in the state is state in the stat

transfer is reasonably sm all for atom ic state form ation. Next, we consider the energy dependence of the peak height of the 1s kaonic atom state coupled to the $[d_{3=2}^{1}]$ proton-hole state in 39 K to determ ine the suitable incident energy to have a large cross section. A swe can see in Fig. 6, the cross section is maximal som ewhere in the range $T_K = 30$ { 40 M eV, and we nd that the cross section has a local maximum value near $T_K = 400$ M eV. From these observations, we take $T_K = 20$ and 400 M eV as the incident kaon energies to calculate the energy spectrum of the em itted proton. We also consider $T_K = 100$ M eV as an energy between 20 and 400 M eV. We mention here that the eikonal approximation is known to be valid only for high energies. Thus, the results for low energies, i.e. $T_K = 100$ M eV, should be regarded as rough estimations.



Fig. 5. Ratio of the e ective numbers of 1s and 2p kaonic atom states form ed with the $[\! d_{3^{-2}}^{-1}]$ proton-hole state in ^{39}K plotted as a function of the incident kaon energy $T_{\rm K}$.

Fig. 6. Double di erential cross section at the resonance peak energy of the kaonic atom 1s state formation with the $[l_{3=2}^{1}]$ proton-hole state in ³⁹K at $_{p}^{Lab} = 0$ [degrees] plotted as a function of the incident kaon kinetic energy.

In Fig.7 we show the calculated spectra for a ^{40}C a target, including contributions from the kaonic atom states up to 4f for $[\text{Id}_{3=2}^{1}]$, $[\text{Id}_{5=2}^{1}]$ and $[2s_{1=2}^{1}]$ proton states in ^{39}K at $T_{\rm K}~=~20$ MeV. The spectra without the widths of the proton-hole states $_{\rm p}$ are represented by the dashed curve. We nd that the contributions coupled to di erent proton-hole states are localized in di erent energy regions and are well separated from each other. This feature of the spectra is di erent from that of the pionic atom form ation in the (d; ^3He) reaction, ^{24)} where the contributions from di erent neutron-hole states overlap. We nd the same features of the spectra for other incident energies. $^{39)}$ The realistic spectra including $_{\rm p}$ are plotted by the solid lines in the same gure. As can be seen in the gure, $_{\rm p}$ is too large to identify each kaonic bound state, except for the $[d_{3=2}^{1}]$ hole state, corresponding to the ground state of the naldaughter nucleus ^{39}K .

We show in Fig. 8 the detailed structure of the kaonic atom formation cross section coupled to the ground state of the daughter nucleus 39 K.Contributions from deeper proton-hole states provide the sm ooth background of the spectra in this



Fig. 7. Kaonic atom formation cross sections for each proton-hole state in the (K ;p) reactions plotted as functions of the em itted proton energy at the incident kaon energy $T_K = 20 \text{ MeV}$ and $\frac{\text{Lab}}{\text{p}} = 0$ [degrees]. The solid and dashed curves are results with and without the widths of the proton-hole states, respectively.

energy region because of their large widths. We nd that the contributions from deeply bound 1s and 2p kaonic atom states are well separated. At $T_K = 20 \text{ MeV}$, the peak due to the 1s state is signi cantly larger than that due to the 2p state, as expected from the result in Fig. 5. The peak height of the atom ic 1s contribution is approximately 7 [b/(sr M eV)] at $T_K = 20$ and 100 M eV. At $T_K = 100 \text{ MeV}$, the 2p peak is enhanced and is approximately 24 [b/(sr M eV)]. The overall shapes of the cross sections at 100 M eV and 400 M eV are similar, while the absolute strength at 400 M eV is approximately 1/3 { 1/4 of that at 100 M eV. We should mention here that the energy resolution of experiments must be good enough so as to observe the separate peak stracture in the spectrum for the atom ic states form ation.



Fig. 8. K aonic atom form ation cross sections in ${}^{40}Ca(K ;p)$ reactions coupled to the $[d_{3=2}^{1}]$ protonhole state in ${}^{39}K$ plotted as functions of the em itted proton energy at ${}^{Lab}_{p} = 0$ [degrees] at the incident kaon energies $T_{K} = 20 \text{ MeV}$, 100 M eV and 400 M eV, respectively.

W e next consider the form ation spectra of the kaonic nuclear states in the (K ;p) reactions. W e choose the incident kaon energy to be $T_K = 600 \text{ MeV}$, for which there exist experimental data.¹⁸⁾ W e show in Fig. 9 the form ation spectra of kaonic nuclei together with those of kaonic atoms as functions of the em itted proton kinetic energies. Here, the widths of the kaonic states are xed to the values listed in Tables I and II. The e ects of the proton-hole widths $_p$ are included. W e found that the spectra do not exhibit any peak-like structure due to the nuclear state form ation but

have only a smooth slope in both the chiral unitary and phenom enological optical potential cases. The contributions from the atom ic state form ation appear as two very narrow peaks around, the threshold energies for both potentials. Each large peak contains several smaller peaks, due to the form ation of several atom ic states, as in the spectrum shown in Fig. 8.



Fig. 9. K aonic nucleus form ation cross sections in 40 C a (K ;p) reactions plotted as functions of the em itted proton energies at ${}_{p}^{L\,ab} = 0$ [degrees] and $T_{K} = 600$ M eV for (left) the chiral unitary m odel and (right) the phenom enological K -nucleus optical potential. The vertical dashed lines indicate the threshold energies, and the sharp peaks around the threshold are due to the atom ic state form ations.

In order to include the phase space e ects on the decay widths for the nalkaon system, we multiply the widths of kaonic states $_{\rm K}$ by the phase space factor de ned in Eqs. (3.6) { (3.9) and calculate the (K ;p) spectra. We present the results in Fig. 10 for both potentials. We not that the spectrum for the chiral unitary potential is not a ected signi cantly by the phase space e ect. However, the (K ;p) spectrum shape for the phenom enological potential is distorted by including the phase space factor and is expected to possess a bum py structure, as reported in Ref. 18).



Fig. 10. K aonic nucleus form ation cross sections in 40 C a (K ;p) reactions plotted as functions of the em itted proton energies at ${}^{Lab}_{p} = 0$ [degrees] and $T_{K} = 600$ M eV for (left) the chiral unitary m odel and (right) the phenom enological K -nucleus optical potential. The vertical dashed lines indicate the threshold energies, and the sharp peaks around the threshold are due to the atom ic state form ations. The energy dependent decay widths for kaonic states are used. (See the main text in details.)

W e perform ed system atic calculations for other target nuclei, 12 C, 16 O and 28 Si

at $T_K = 600 \text{ M eV}$ and present the results in Fig.11. In Ref.18), experimental data for a ¹⁶O target are reported and data for ¹²C and ²⁸Si targets could be obtained in the future.⁴⁰⁾ As shown in Fig. 11, we have found that some bum py structures in the (K ;p) spectra m ay appear, due to the form ation of the kaon nucleus states, especially in the case of the ¹²C target, if the kaon-nucleus optical potential is as deep as 200 M eV, as reported in Ref. 27). On the other hand, if the depth of the optical potential is as shallow as 50 M eV, as predicted by the chiral unitary m odel, the spectrum m ay not possess any bum py structures, but only exhibit a sm ooth slope for all targets considered here.

F inally, we present energy integrated cross sections $\frac{d}{d}$ for kaonic nuclear 1s state form ation for ¹²C and ²⁸Si targets with the results obtained in Refs. 10) and 17). We found that our results with the phenom enological optical potential qualitatively agree with those in Ref. 10) and are smaller than those in Ref. 17). The present results with the chiral unitary potential are signi cantly larger than those with the phenom enological potential because of the smaller momentum transfer in the (K ;p) reactions due to the smaller binding energies of kaonic nuclear 1s states.

Table IV. Energy integrated cross sections for the form ation of kaonic 1s nuclear states in units of [b=sr]. The results in Refs.10) and 17) are also shown for comparison. Proton hole states are $[1p_{3=2}]^{1}$ and $[1d_{5=2}]^{1}$ for ¹²C and ²⁸Sitargets, respectively.

| | (d =d) (K p) [b=sr] | | | | | | |
|------------------|----------------------|----------------|---------|---------|--|--|--|
| Target nucleus | ChiralUnitary | Phenom enology | Ref.10) | Ref.17) | | | |
| ¹² C | 425 | 65 | 47 | 100{490 | | | |
| ²⁸ Si | 92.6 | 2.7 | 6.0 | 35{180 | | | |

x4. Conclusion

W e have studied the structure and form ation of kaonic atom s and kaonic nuclei in this paper. W e used two di erent kaon-nucleus optical potentials, which are obtained from the chiral unitary m odel and a phenom enological t of existing kaonic atom data. W e theoretically studied the structure of kaonic atom s and kaonic nuclei using these potentials and determ ined the di erences between the obtained level schem es of the kaonic nuclear states.

W e also studied the form ation cross sections of deeply bound kaonic atom s and kaonic nuclei which cannot be observed with standard X-ray spectroscopy. All the atom ic states are theoretically predicted to be quasi-stable. We investigate the (K ;p) reaction theoretically and evaluate the cross section of the ${}^{40}Ca(K ;p)$ reaction in detail. For deep atom ic state form ation, the cross sections are predicted to be approximately 7 [b/(sr M eV)] at $T_K = 20$ and 100 M eV and 2 [b/(sr M eV)] at 400 M eV for kaonic atom 1s state form ation. For the atom ic 2p state, the cross section is predicted to be approximately 24 [b/(sr M eV)] at $T_K = 100 \text{ M eV}$.

We also system atically studied the form ation cross sections of kaonic nuclear states in (K ; p) reactions for various targets. In order to take into account the phase space suppression e ects of the decay widths, we introduced a phase space factor to obtain the (K ; p) spectra. We found in our theoretically studies that in



Fig.11. Same as Fig. 10, except that here the target nuclei are (top) 12 C, (m iddle) 16 O, and (bottom) 28 Si.

the (K ;p) reactions, a certain bum py structure due to kaonic nucleus form ation can be seen only for the case of a deep (200 MeV) phenom enological kaon nucleus potential. Due to the phase space suppression, the decay widths of kaonic states become so narrow that we can see certain bum py structure in the reaction spectrum, which could be seen in experiments. For the case of the chiral unitary potential, the binding energies are too small to reduce the decay widths and to see the bum py structure in the spectra of the (K ;p) reactions. However, we should properly include the energy dependence of the chiral unitary potential in future studies of kaonic nuclear states to evaluate m ore realistic form ation spectra.

In order to obtain m ore conclusive theoretically results, we need to apply G reen function m ethods for states with large widths⁴¹⁾ and to consider the energy dependence of the optical potential properly. Furtherm ore, we should consider the changes and/or deform ations of the nucleus due to the existence of the kaon inside and solve the problem in a self-consistent m anner for kaonic nucleus states. How ever, we believe that the present theoretical e ort to evaluate the absolute cross sections for the kaonic bound state form ation are relevant for determ ining a suitable m ethod to observe them and helpful for developing the physics of kaon-nuclear bound system s and kaon behavior in nuclear m edium. Further investigations both theoretically and experimentally are needed to understand kaon behavior in nuclear m edium m ore precisely.

A cknow ledgem ents

W e acknowledge E.Oset and A.Ram os for stimulating discussions on kaon bound systems and the chiral unitary model. W e would like to thank M. Iwasaki and T.Kishim oto for stimulating discussions on the latest experimental data of kaonic nucleus formation. W e also would like to thank T.Yamazaki, Y.Akaishi, and A.D ote for useful discussions on theoretical aspects of kaonic nucleus systems. W e are grateful to H.Toki and E.H iyama for many suggestions and discussions regarding the kaon-nucleus systems. W e also thank A.G al for his careful reading of our preprint and useful comments. This work is partly supported by G rants-in-A id for scienti c research of M onbuK agakusho and Japan Society for the P rom otion of Science (No. 16540254).

References

- [1] M.Alberg, E.M. Henley and L.W ilets, Ann. of Phys. 96 (1976), 43.
- [2] R.Brockmann, W.Weise and L.Tauscher, Nucl. Phys. A 308 (1978), 365.
- [3] M.M izoguchi, S.H irenzaki and H.Toki, Nucl. Phys. A 567 (1994), 893.
- [4] C.J.Batty, Nucl. Phys. A 372 (1982), 418.
- [5] N.Kaiser, P.B.Siegel and W.Weise, Nucl. Phys. A 594 (1995), 325.
- [6] E.Oset and A.Ramos, Nucl. Phys. A 635 (1998), 99.
- [7] T.Waas, N.Kaiser and W.Weise, Phys.Lett. B 365 (1996), 12; B 379 (1996), 34.
 T.Waas and W.Weise, Nucl. Phys. A 625 (1997), 287.
- [8] M.Lutz, Phys.Lett.B 426 (1998), 12.
- [9] A.Ram os and E.O set, Nucl. Phys. A 671 (2000), 481.
- [10] A.Cieply et al, Nucl. Phys. A 696 (2001), 173.
- [11] L.Tolos et al, Nucl. Phys. A 690 (2001), 547.
- [12] L.Tolos et al, Phys.Rev.C 65 (2002), 054907.
- [13] S. Hirenzaki, Y. Okumura, H. Toki, E. Oset and A. Ramos, Phys. Rev. C 61 (2000), 055205.
- [14] E.Friedm an and A.Gal, Phys.Lett.B 459 (1999), 43.
- [15] E.Friedm an and A.Gal, Nucl. Phys. A 658 (1999), 345.
- [16] J.Nieves and E.Oset, Phys. Lett. B 282 (1992), 24.
- [17] T.Kishim oto, Phys. Rev. Lett. 83 (1999), 4701.
- [18] T.Kishim oto et al., Prog. Theor. Phys. Suppl. No.149 (2003), 264.
- [19] M. Iwasakiet al, nucl-ex/0310018.
- [20] T.Suzukiet al, Phys.Lett.B 597 (2004), 263.
- [21] Y.Akaishiand T.Yamazaki, Phys.Rev.C 65 (2002), 044005.
- [22] A.Dote, H.Horiuchi, Y.Akaishi and T.Yam azaki, Phys. Lett. B 590 (2004), 51.
- [23] H. Tokiand T. Yam azaki, Phys.Lett. B 213 (1988), 129.
 H. Toki, S. Hirenzaki, R. S. Hayano and T. Yam azaki, Nucl. Phys. A 501 (1989), 653.
- [24] S. Hirenzaki, H. Toki and T. Yam azaki, Phys. Rev. C 44 (1991), 2472.
- H.Toki, S.Hirenzakiand T.Yam azaki, Nucl. Phys. A 530 (1991), 679.
- [25] H.Gilgetal, Phys. Rev.C 62 (2000), 025201.
- [26] K.Itahashiet al, Phys.Rev.C 62 (2000), 025202.
- [27] C.J.Batty, E.Friedm an and A.Gal, Phys. Rep. 287 (1997), 385.
- [28] E.O set and L.L.Saledo, J.Com put. Phys. 57 (1985), 361.
- [29] J.Nieves, E.Oset, C.Garcia-Recio, Nucl. Phys. A 554 (1993), 509.
- [30] Review of Particle Physics, Particle Data G roup, Phys. Lett. B 592 (2004), 1.
- [31] S.L.Belostotskiiet al, Sov.J.Nucl.Phys.41 (1985), 6.
- [32] M.Yosoietal, Nucl. Phys. A 738 (2004), 451.
- [33] F.A jzenberg-Selove, Nucl. Phys. A 523 (1991), 1.

Form ation of Kaon-Nucleus Bound States by In-ight (K ;p) Reactions 17

- [34] J.M ougey et al., Nucl. Phys. A 262 (1976), 461.
- [35] U.Amaldiet al, Phys. Rev. Lett. 13 (1964), 341.
- [36] K.Nakamura et al, Phys.Rev.Lett.33 (1974),853.
- [37] R.B.Firestone et al., Table of Isotopes, 8th edition (John W iley & Sons, Inc., 1996)
- [38] J.M ares, E.Friedm an and A.G al, Phys.Lett.B 606 (2005), 295.
- [39] Y.Okumura, Master Thesis, Nara Women's University (2000).
- [40] T.Kishim oto, private communication.
- [41] O.Morim atsu and K.Yazaki, Nucl. Phys. A 435 (1985), 727; Nucl. Phys. A 483 (1988), 493.