# Quantum Chaos, Degeneracies and Exceptional Points

W D. Heiss and S. Radu

C entre for N onlinear Studies and D epartm ent of P hysics U niversity of the W itwatersmand, PO W its 2050, Johannesburg, South A frica

It is argued that, if a regular H am iltonian is perturbed by a term that produces chaos, the onset of chaos is shifted towards larger values of the perturbation parameter if the unperturbed spectrum is degenerate and the lifting of the degeneracy is of second order in this parameter. The argument is based on the behaviour of the exceptional points of the full problem .

PACS Nos.: 36.40.+ d, 05.45.+ b

## I. IN TRODUCTION

The study of single particle motion in deformed mean elds has attracted much attention recently because of its relevance to nuclear physics and for the description of metallic clusters [1,2]. For the case of harmonic oscillator potentials, deformations which go beyond a quadrupole deformation are nonintegrable and show chaotic behaviour in the classical and quantum mechanical treatment. However, closer scrutiny has revealed that the addition of an octupole term to a prolate quadrupole potential produces the typical signatures of chaos only for fairly large octupole strength, in fact, the problem appears to be close to integrability. Moreover, the quantum mechanical treatment produces new shell structures for nite octupole strength even though the classical problem is nonlinear [2,3]. In contrast, addition of an octupole term to an oblate quadrupole deform ed potential yields chaos with a positive Lyapunov exponent, and the corresponding quantum spectrum has the typical statistical properties ascribed to quantum chaos.

In this paper we address the question: what are the intrinsic properties of the quantum mechanical operators which give rise to the dimensional described above. This question is of interest for a possible characterisation of what is called quantum chaos. The example mentioned in the previous paragraph renders a good case for such studies as it refers to the orthodox situation where a classically chaotic system is treated quantum mechanically. Our aim is to unravel the universal operator properties that produce the typical patterns ascribed to quantum chaos without reference to an underlying classical system. There is progress in previous work towards this aim as it has been recognised that, for a problem of the form H<sup>0</sup> + H<sup>1</sup>, it is precisely the high density of the exceptional points [4,5], i.e. the singularities of the spectrum  $E_n$  (), which bring about the statistical properties of the spectrum associated with quantum chaos [6]. The new aspect in the present paper is the exceptional points are of chaos is suppressed and will occur only for larger values of . It is this pattern which explains the dimension between the prolate and oblate case as described above. The major point of the present paper is the universal validity of this main quantum the points around = 0 provides for the explanation.

In the following section we introduce brie y the concept of exceptional points and present the argument for the statement made above. Section three contains examples for illustration and section four refers to the particular physical example introduced above. A summary and discussion is given in the last section.

#### II. EFFECT OF DEGENERACIES ON EXCEPTIONAL POINTS

To make the paper selfcontained we brie y recapitulate the signi cance of exceptional points and their connection to avoided level crossings. This will facilitate the discussion about the e ect of degeneracies on exceptional points and hence the global structure of the spectrum.

There is essentially a one to one relationship between avoided level crossings and exceptional points [6]. If we have a quantum mechanical problem of the form  $H^0 + H^1$  with  $H^0$  and  $H^1$  given as N N matrices, then the spectrum  $E_n()$ ; n = 1; :::;N is determined by one analytic function evaluated on N Riemann sheets [5]. If there are no degeneracies, the N sheets are connected by N (N 1) branch points, the exceptional points. They occur in complex conjugate pairs and are the points in the complex -plane where any two pairs of energies coalesce. If this happens su ciently close to the real -axis, an avoided level crossing occurs since the pair of energies coalescing in the complex plane still assumes values near to each other on the neighbouring real -axis. In principle, the positions of the

exceptional points are determ ined by the resultant of the secular equation for the spectrum, i.e. by the simultaneous solution of the two ploynom ial equations

$$\det (E \quad H^0 \quad H^1) = 0$$

$$\frac{d}{dE} \det (E \quad H^0 \quad H^1) = 0$$
(1)

which leads by elimination of E to the resultant which is a polynom isolforder N (N 1) in  $\cdot$ . For large values of N, an explicit calculation of the exceptional points is prohibitive. However, it is possible to determ ine a distribution for the real parts of the exceptional points from the know ledge of H  $^0$  and H  $^1$  alone [6], and in this way the regions of real

-values with large scale avoided level crossings can be found. These are the regions where the spectrum shows the properties ascribed to quantum chaos. This method has been tested and applied to the particular physical situation of the hydrogen atom in a strong magnetic eld [7].

The e ect of removing degeneracies by small random perturbation was discussed previously [8]. In the present paper we address the particular problem of lifting degeneracies by the term  $H^{1}$ , i.e. we assume that  $H^{0}$  has certain system atic degeneracies (like for example the harm onic oscillator). Guided by the particular physical problem as discussed in the introduction and resumed in section four, we consider and compare two situations which turn out to be signi cantly di erent in the context of quantum chaotic behaviour. The two situations are characterised by the presence of a linear term in the expansion of  $E_n$  () in powers of around = 0.

It is obvious that the spectrum has a distinctly di erent behaviour in the vicinity of = 0 for the two cases considered. A non-vanishing linear term exhibits the typical behaviour of a lifting of degeneracies, when a perturbation is switched on, in that the levels fan out of the degenerate level when is turned on. W ith a zero linear term the levels stay closely together and separate only when the quadratic term becomes signi cant. From this picture one intuitively expects that, if the additional term H<sup>1</sup> produces chaos at all, the onset of chaos is delayed for increasing in the latter case when comparing with the form er case. In the following we con in this expectation using the

exceptional points for the argument, while the next section presents some illustrative examples.

Each m -fold degeneracy reduces the total num ber of exceptional points by the num berm (m 1) if the linear term s are present in the expansion of the energy levels. If the leading perturbing terms are quadratic, there is a further 1). This follows from the resultant which starts with m(m-1) as the lowest reduction by the same numberm (m order in the former case and with <sup>2m (m 1)</sup> in the latter case. This behaviour is valid for each m-fold degeneracy, in other words, if the levels have at = 0 the degeneracies  $m_1; m_2; :::$ , the low est order term s of the resultant start with  $m_1(m_1 \ 1)$ m<sub>2</sub> (m<sub>2</sub> 1) when the lifting of the degeneracy is linear and  $w \operatorname{ifth}^{m_1 (m_1 - 1)}$ 2m<sub>2</sub>(m<sub>2</sub> 1) a quadratic lifting. A particular m-fold degeneracy can be viewed as a con uent situation where m (m 1) com plex conjugate branch points have merged into the point = 0 thus cancelling the singularities altogether; likewise, in the quadratic case, 2m (m 1) singularities have m erged. The essential point of the argum ent lies in the com parison between the two cases: switching on linear terms leads to the emergence of  $m_i(m_i)$ 1) additional exceptional points. The distance from = 0 of the additional exceptional points depends on the magnitude of the linear term s. For typical values they are rather closer to than rem ote from the origin. For large m atrices the additional num ber of exceptional points can amount to many thousands. This causes avoided level crossings in the spectrum in a region where without the linear term the spectrum appears sm ooth.

### III. ILLUSTRATION OF SIM PLE EXAM PLES

The realistic example which initiated this work produces more than twenty nine thousand exceptional points for the matrix size used in our work [2]. We return to its treatment in the following section. Here we illustrate the behaviour of the exceptional points in a low dimensional example. The e ect on the spectrum, in particular the dimensional of quantum chaos for the two dimensional examples, is subsequently demonstrated in a generic matrix model.

We consider a nine dimensionalm odel where H<sup>0</sup> has the diagonal entries (1,1,1,2,2,2,3,3,3). The three-fold degeneracies are lifted by the term H<sup>1</sup> with

| (                   | )     | 0   | 0   | 0:5 | 2   | 1:5 | 2   | 1:5 | 0             |
|---------------------|-------|-----|-----|-----|-----|-----|-----|-----|---------------|
| H <sup>1</sup> () = | 30    | 0   | 0   | 1:5 | 2   | 3   | 0:5 | 1:5 | 0:5 C         |
|                     | 30    | 0   |     | 2   | 3   | 1   | 2   | 1:5 | 2 Č           |
|                     | 305   | 1:5 | 2   |     | 0   | 0   | 1:5 | 2:5 | οğ            |
|                     | 2     | 2   | 3   | 0   | 0   | 0   | 1   | 1:5 | 1 Č           |
|                     | 15    | 3   | 1   | 0   | 0   |     | 1   | 1:5 | 0 <b>:</b> 5C |
|                     | 2     | 0:5 | 2   | 1:5 | 1   | 1   |     | 0   | 0 8           |
|                     | 9 0:5 | 1:5 | 1:5 | 2:5 | 1:5 | 1:5 | 0   | 0   | 0 A           |
|                     | 0     | 0:5 | 2   | 0   | 1   | 0:5 | 0   | 0   |               |

(2)

for

For  $= 0, H^{1}$  has only zero entries in the degenerate subspaces of H<sup>0</sup>, therefore the nine levels of the full problem have a vanishing linear term when expanded at = 0.0 there ise the entries of H<sup>1</sup> have no signi cance, they have been generated random ly, any other choice (with similar orders of magnitude) yields the same equalitative result. A lso, a diagonal form of H<sup>1</sup> in the degenerate subspaces can always be achieved by orthogonal transform ations within the subspaces without changing the global spectrum. The speci c form chosen is convenient as it leaves out additional irrelevant parameters and it is in line with the physical situation discussed in the following section.

A nine dimensional model yields 36 exceptional points in each half plane, our particular case reduces this num ber to 18. In Fig.(1a) we display most of the exceptional points in the upper -plane. When the parameter is switched on, nine additional exceptional points emerge from = 0 in each half plane. This is illustrated in Fig.(1b). The basic di erence between the two gures lies in the additional singularities scattered around = 0. Note that the other exceptional points change only moderately under variation of . The corresponding spectra are displayed in Figs.(2). The growing distance from the origin of the avoided crossings with increasing value is clearly discernible in Figs.(2c) and (2d), where only part of the spectrum is shown for better illustration. These avoided crossings are due to the 'new' exceptional points. Note that in particular the levels which originate from the same degenerate energy are a ected. When this occurs on a large scale, the nearest neighbour distribution of the spectrum will typically assume the W igner sum ise for rather sm all values of while there is no resemblence to the W igner curve when the linear term s of the perturbation vanish. This is demonstrated in the follow ing study case.

E sentially we repeat the model used above. We consider N > 100 for the full dimension and the dimensions of the degenerate subspaces can be two, three, four and so on. The diagonal matrix H<sup>0</sup> contains the entries (1,1,...,2,2,...) and the entries of H<sup>1</sup> are led random ly with elements between 1 and + 1. The o -diagonal elements of the diagonal blocks of the degenerate subspaces are set equal to zero while the corresponding diagonals are led with the num bers ( $n_{deg}=2+1=2$ ;  $n_{deg}=2+1$ ;::; $n_{deg}=2$  1=2) with  $n_{deg}$  being the dimension of the degenerate subspace. Because of the huge number of exceptional points, we no longer focus our attention on their positions. Instead, we com pare nearest neighbour distributions for = 0 and  $\notin$  0 for sm all values of .

In Figs.(3) typical results for the two di erent cases are presented. For the same value of sample averages of ten samples of nearest neighbour distributions are illustrated for = 0 and = 1. There is of course a continuous transition from Fig.(3a) to Fig.(3b), and only if the order of magnitude of the diagonal elements of H $^1$  has reached that of the other matrix elements, that is for '1, the W igner distribution has fully developed. We can understand, in terms of the exceptional points, why for 0 < 1 the situation closely resembles that of = 0 even though a large number of exceptional points has already en erged from the origin of the -plane. When the exceptional points are still very close to the origin, that is very close to each other, there is a cancellation with regard to the e ect on the spectrum . In fact, two square root branch points connecting the sam e Riem ann sheets, are barely noticeable from a distance; the function  $z^{2} \overline{z^{2} + u^{2}}$  boks just like  $z^{2}$  for  $z^{2}$  . Only when the branch points have moved out su ciently far does the e ect on the spectrum become signi cant as clearly seen in Figs. (3). The results presented refer to  $n_{deg} = 8$  and N = 304, which yields 2128 exceptional points having emerged from the origin. When  $n_{deg}$ is increased, the window of -values, where this transition is strongly pronounced, becomes wider. Conversely, for smaller values of n<sub>deg</sub> the e ect is clearly seen only in a smaller range of -values. This follows from the higher density of exceptional points associated with larger values of ndeq. Also, the -values beyond which a plain Wigner distribution occurs even when = 0 m oves closer tow ands the origin with increasing  $n_{deq}$ . These two observations can qualitatively be understood from perturbative argum ents in that we expect the onset of a W igner distribution beyond the intersection point of two straight lines or parabolae from neighbouring levels for  $\frac{1}{2}$  0 or = 0, respectively; the curves intersect at intset which is proportional to  $1=n_{deg}$  in the form er and  $1=p_{n_{deg}}$  in the latter case. This argument indicates where perturbation breaks down; we recall that the present paper deals with a nonperturbative situation as it is just the exceptional points which break down perturbation. We note that in a typical physical situation, like the one discussed in the next section, one is usually faced with steadily increasing values of ndeg with increasing energy.

#### IV . A PHYSICAL EXAMPLE

Here we present the physical example [2] which actually initiated this work. Phenom enological mean elds which contain quadrupole and octupole deform ations have been investigated classically [9] and quantum mechanically [1,2] as this is of interest in nuclear physics and more recently also for the description of metallic clusters. As a detailed discussion is presented in the quoted papers we here focus our attention on the aspect of interest in the present context.

The single particle potential

$$V(\mathfrak{F};z) = \frac{m}{2}!^{2}(\mathfrak{F}^{2} + \frac{z^{2}}{b^{2}} + \frac{2z^{3}}{p}\frac{3z\mathfrak{F}^{2}}{\mathfrak{F}^{2} + z^{2}})$$
(3)

is, for b > 1 (b < 1), a quadrupole deform ed harm onic oscillator of prolate (oblate) shape with an additional octupole term; in fact the term multiplying is proportional to  $r^2P_3$  (cos) with  $P_3$  the third order Legendre polynom ial. We use cylindrical coordinates z and  $\$ = \frac{p}{x^2 + y^2}$ . For € 0 this is a two degrees of freedom system which is non-integrable. For b < 1 it turns out that the switching on of the octupole term very quickly gives rise to classically chaotic behaviour while the onset of chaos is barely discernable when b 2. The statistical analyses of the corresponding quantum spectra reveal the expected results in that for b < 1 the W igner sum ise is obtained for the nearest neighbour distribution while for b 2 a Poisson distribution is obtained, and for particular values of even a new shell structure emerges. The latter is understood in term s of corresponding classical periodic orbits [1,2].

In the spirit of the present paper the quantum mechanical ndings should be directly obtained from the matrix structure of the associated H am iltonian and the exceptional points related to it. The appropriate basis where H  $^0$  =  $p^2=2m + m!^2(k^2 + z^2=b^2)=2$  is diagonal is given by the occupation numbers  $n_2$  and  $n_z$ . Note that the z-component of the angular momentum is conserved and we consider here only  $l_z = 0$ . The arrangement of the quantum numbers in the matrix H  $\frac{1}{(n_z,n_z);(n_z^0,n_z^0)}$  is determined by the arithmetic ascending order of the unperturbed levels  $E_{n_z,n_z}^0$ . The arrangement will therefore depend on the value of b. The selection rules restrict entries in H<sup>1</sup> to  $n_{z}^{2} = n_{z}$ (2k + 1)and  $n_2^0 = n_2$ (2k) with  $k = 0;1;2;\ldots$  It turns out that for b 2 the entries vanish in the blocks of H<sup>1</sup> which refer to the subspaces in which the unperturbed energies  $E^{0}$  are degenerate. As a consequence, the perturbative expansion at = 0 of the eigenvalues starts with the quadratic term . From the discussion in the previous section it follows that a great number of exceptional points is trapped at = 0 and we therefore expect the onset of quantum chaos only for values of at an appreciable distance from zero. Using 544 544 m atrices we illustrate in Fig.(4a) the distribution of the real parts of the exceptional points for the matrix problem H<sup>0</sup> + H<sup>1</sup> with b = 2. The bulk of the exceptional points occur in fact at values of which fall outside the physical range (for = c 1 the potential no longer binds). In Fig.(4b) we diplay the corresponding distribution for b = 1=2. Now we obtain the maximum density of exceptional points as soon as the parameter is switched on. This is in accordance with the results of the previous section since now the linear term does occur in the expansion of each level around = 0 as there are non-zero entries in the blocks of H<sup>1</sup> which refer to the degenerate subspaces. The e ect of this di erence in the distribution of the exceptional points manifests itself in the di erent behaviour of the respective quantum spectra in that a W igner distribution is found for > 0 when b = 1=2 while for b = 2 a W igner distribution never develops for  $< _{c}$ .

## V.SUMMARY AND DISCUSSION

The previous section provides a ne example where the statistical properties of the quantum spectrum can be predicted from the properties of the individual matrices H<sup>0</sup> and H<sup>1</sup> alone. To obtain the distribution of the exceptional points as illustrated in Figs.(4) it is not necessary to solve the full problem H<sup>0</sup> + H<sup>1</sup> let alone to determ ine their positions (which would be in possible for more than 295,000 exceptional points). The distribution was found using the method developed in [6,7] and explained in the Appendix. There, only some simple properties of H<sup>0</sup> and H<sup>1</sup> are used. W hile this result alone yields just another commation of the method employed in [7], the new aspect of this paper lies in the prediction that, if the unperturbed levels are degenerate to second order, level statistics ascribed to quantum chaos are substantially suppressed initially and become manifest only for su ciently large values of the perturbation. The argument comes about rather naturally from the behaviour of the exceptional points which in turn determ ine the degree of quantum chaos [6]. W hile the previous section provides the physical relevance of our know ledge, this is the rst example where even ner details with regard to level statistics can be extracted from the distribution and the general behaviour of the exceptional points. W e believe that m ore re nem ent can eventually even predict new shell structure as the one discovered in the model discussed. W ork towards this aim is in progress.

#### APPENDIX A: DISTRIBUTION OF THE REAL PARTS OF THE EXCEPTIONAL POINTS

We use the method of the unperturbed curves [7], where the actual spectrum is approximated by simple algebraic curves. The intesection points of the approximate curves are then identified with the real parts of the exceptional points. Since the aim is to obtain the distribution rather than the excat positions, the approximate curves succe. They are obtained from the individual matrices H<sup>0</sup> and H<sup>1</sup> only. The requirement that the actual spectrum and the approximate curves coincide exactly for small and for large values of leads for the prolate case (b = 2) to the form

$$F_{i}() = p_{3} \frac{(1 + i)^{2}}{!_{i}^{3} + 3!_{i}^{2} + c_{i} + d_{i}} \text{ for } 0.9_{c}$$

where the " $_{i}$  and ! $_{i}$  are the eigenvalues of H  $^{0}$  and H  $^{1}$ , respectively, and

$$_{i} = \frac{X}{_{n \in i}} \frac{\mathcal{H}_{i,n}^{1} \mathcal{L}}{_{i,n}^{*} \mathbf{u}_{n}^{*}}; \qquad _{i} = (U H^{0} U^{-1})_{i;i}$$

with U being the orthogonal matrix that diagonalises H $^1$ . The coe cients  $c_i$  and  $d_i$  are determined so as to sm oothly m atch the two curves at = 0.9 c. We note that the second order perturbation is correct up to third order terms in

and is in fact very near to the actual spectrum for  $0.9_{c}$ , while the expression for  $0.9_{c}$  is correct up to term s of the order 1= .

For the oblate case (b = 1=2) the spectrum is, for the purpose considered, su ciently well approximated by the unperturbed lines

$$G_{i}() = "_{i} + !_{i}$$

Recall that the eigenvalues "i and !i are di erent from those in the prolate case.

- [1] R Balian and C. Bloch, Ann Phys. (N Y.) 69, 76 (1972); R Arvieu, F Brut, JC arbonell, and J.Touchard, PhysRev. A 35, 2389 (1987); W. A. Heer, Rev. M od. Phys. 65, 611 (1993); M. Brack, Rev. M od. Phys. 65, 677 (1993); JB locki, J.-J.Shi, and W. J.Swiatecki, Nucl. Phys. A 554, 387 (1993).
- [2] W D.Heiss, R.G.Nazm itdinov and S.Radu, Phys.Rev.Lett. 72, 2351 (1994); W D.Heiss, R.G.Nazm itdinov and S.Radu, Phys.Rev.B 51, 1874 (1995); W D.Heiss, R.G.Nazm itdinov and S.Radu, Phys.Scrpt., to appear
- [3] K Arita, Prog. Theor. Phys. 90, 747 (1993); K Arita and K M atsuyanagi, ibid. 91, 723 (1994).
- [4] C M .Bender and T T .W u, Phys. Rev. D 7, 1620 (1973); P E .Shanley, Ann. Phys. (NY) 186, 292 (1988); A G .U shveridze, J. Phys. A 21,955 (1988);
- [5] W D.Heiss and W.H Steeb, J.M ath. Phys. 32, 3003 (1991).
- [6] W D.Heiss and A.L.Sannino, JPhys A: Mathematical and General, 23, 1167 (1990)
- [7] A A.Kotze and W D.Heiss, J.Phys.A: Math Gen, 27, 3059 (1994)
- [8] W D. Heiss and M. Muller, Phys. Rev. A 48, 2558 (1993); W D. Heiss and J.C. H. Chiang, Phys. Rev. A 47, 2533 (1993)
- [9] W D.Heiss and R.G.Nazm itdinov, Phys.Rev.Lett. 73, 1235 (1994)

### Figure captions

Fig.1 Exceptional points of the 9-dimensional model in the complex -plane for = 0 (a) and = 1=2 (diamonds in (b)) and = 1 (crosses in (b)). Note the di erence of the scale in (a) and (b) and the motion away from the origin when increases.

Fig.2 Spectra of the 9-dimensional model for = 0 (a) and = 1 (b). For better illustration a partial view is given for = 1=2 (c) and = 1 (d). Note how the position of the avoided crossing not only moves outwards but also become s more pronounced with increasing value of .

F ig 3 Sample averages of nearest neighbour distributions of 304 levels for = 0 (a) with an eightfold degeneracy at = 0 and for = 1 (b); both distributions are for = 02.

F ig 4 D istribution of the real parts of the exceptional points for the prolate case (a) and the oblate case (b).