# The hadron-quark transition with a lattice of nonlocal con ning solitons

Charles W . Johnson and George Fail

Center for Nuclear Research, Department of Physics, Kent State University, Kent, OH 44242

M ichael R . Frank<sup>z</sup>

Institute for Nuclear Theory, University of W ashington, Seattle, W A 98195

#### Abstract

We use a lattice of nonlocal con ning solitons to describe nuclear matter in the W igner-Seitz approximation. The average density is varied by changing the size of the W igner-Seitz cell. At su ciently large density quark energy bands develop. The intersection of the led valence band with the next empty band at a few times standard nuclear density signals a transition from a color insulator to a color conductor and is identified with the critical density for quark decon nement.

Typeset using REVT<sub>E</sub>X

electronic m ail: johnson@ksuvxd.kent.edu

<sup>&</sup>lt;sup>y</sup>electronic m ail: fai@ ksuvxd.kent.edu

<sup>&</sup>lt;sup>z</sup>electronic m ail: frank@ phys.w ashington edu

The search for the hadron-quark transition in nuclear collisions has provided powerful m otivation for experimental and theoretical research in high-energy nuclear physics. Experimental results are now emerging from Brookhaven's Alternating Gradient Synchrotron (AGS) and CERN's Superconducting Proton Synchrotron (SPS), while the planned Relativistic Heavy Ion Collider (RHIC) at Brookhaven represents the rst major construction priority of the US nuclear physics community and is expected to be operational before the turn of the century. W ork on the theory side can be broadly categorized into studying the fundamental properties of the strong interaction and exploring phenomenological aspects such as the signatures and characterization of the guark-gluon plasm a (QGP).W hile phenomenological support for experiment is crucial for the success of the overall e ort, in the present paper we address the more fundamental question of modeling the hadron-quark transition.

M odeling the expected transition will remain in portant as long as a complete solution of quantum chromodynamics (QCD) in a reasonably large space-time volume is out of reach. Soliton matter was proposed as a model of dense nuclear matter by several authors [1{3}. In the following we focus attention on the behavior of soliton matter as a function of density at zero temperature. The simplest approach in this context is to think about nuclear matter as a crystal of MIT bags [4]. In more elaborate many-soliton models the rigid walls of the MIT bags are replaced by external elds which are supposed to represent the collective nonperturbative elects of QCD. [5] Several aspects of this picture were developed further for the case of a single soliton in the fram ework of the Color Dielectric M odel (CDM) [6].

The alternative approach taken here is based on the hadronization [7,8] of a model truncation of QCD, known as the G lobalColorM odel (GCM) [9]. The GCM is de ned by the action

$$S_{GCM}[\bar{q};q] = \int_{-\infty}^{Z} d^{4}x d^{4}y \ \bar{q}(x) \quad (\qquad 0 + m_{q}) \quad (x \quad y) q(y) + \frac{q^{2}}{2} j^{a}(x) D(x \quad y) j^{a}(y) ; \quad (1)$$

where  $j^{a}(x) = \overline{q}(x)$  is the color quark current. QCD phenom enology such as con ne-

m ent is incorporated into the quark-quark interaction, D<sup>1</sup> Thism odel has proven successful in the description of low-energy QCD through the reproduction of chiral perturbation theory [9,10], m eson form factors [11] and spectra [12,13], and the soliton [14] and Fadeev [7] descriptions of the nucleon. The relationship of this model to other models of QCD has recently been summarized [15].

The form ation of solitons is achieved by m inim izing the energy functional for the system at the mean eld level with the baryon number constrained by a chemical potential. For sim plicity, as with the previous treatment of this model [14], we restrict our attention here to a scalar-isoscalar mean eld. The self-consistent solution of the mean-eld equations entails iterating to convergence the D irac equation for the vacuum -dressed quarks and a nonlinear K lein-G ordon equation for the scalar eld.

There are two novel features of this model that are particularly relevant to the present discussion. First, the meson elds and their self interactions are generated through the hadronization of Eq. (1). This implies in particular that the \m acroscopic" behavior is determined entirely from the underlying quark-quark interaction and parameters which enter at the quark level. Second, the con nem ent mechanism that arises here is due to vacuum repulsion as supplied by the quark self-energy dressing. This point is demonstrated as follows.

The quark-quark interaction, D in Eq. (1), can be chosen to produce a quark selfenergy, (p) = i  $p[A(\vec{p}) \ 1] + B(p^2)$ , which prohibits the propagation of quarks in the vacuum. Here we take for example in momentum space  $g^2D(q) = 3^{-4-2-(4)}(q)$ , which in the rainbow approximation [16] to the D yson-Schwinger equation gives the result

$$A (p^{2}) = \begin{cases} 8 \\ \gtrless \\ 2 \\ \frac{1}{2} \left[1 + \left(1 + \frac{2^{2}}{p^{2}}\right)^{\frac{1}{2}}\right] \end{cases} ; B (p^{2}) = \begin{cases} 8 \\ \gtrless \\ 2 \\ 4 \\ 2 \\ \frac{1}{2} \left[1 + \left(2 + \frac{2^{2}}{p^{2}}\right)^{\frac{1}{2}}\right] \end{cases} ; B (p^{2}) = \begin{cases} 8 \\ \gtrless \\ 4 \\ \frac{1}{2} \left[1 + \frac{2^{2}}{p^{2}}\right]^{\frac{1}{2}} \\ \frac{$$

<sup>1</sup>For convenience a gauge is chosen here for which the e ective gluon two-point function is diagonal in Lorentz indices, D (x y) = D (x y), and the Euclidean metric is employed. That the form s (2) produce a model of connement as described above can be seen by the absence of a solution to the equation  $p^2 + M^2(p^2) = 0$ , with M = B = A (no on-mass-shell point).

The mean eld produces a cavity in the vacuum in which the quarks can propagate. For example, in the case of quarks coupled to a constant scalar mean eld by the scalar self-energy function<sup>2</sup> B, the quark inverse G reen's function obtains the following form :  $G^{-1}(p) = i (pA(p^2) + B(p^2)(1 + ), where 0$  2 characterizes the strength of the mean eld. In this case a continuous single-particle energy spectrum,  $E^{-2}(p^2) = p^2 + M_c^2$ , is obtained because of the constant potential, with the constituent m ass given by

$$M_{\rm C}^{2} = \frac{2}{4} \frac{(1+)^{2}}{1(1+)^{2}} :$$
 (3)

From (3) it is evident that the increase of the constituent m ass as the strength, , of the m ean eld decreases toward its vacuum value (= 0), is due to the repulsive interaction with the vacuum. For the case of the constant m ean eld, the quarks are allowed to propagate throughout space. However, due to the energy stored in the m ean eld, the self-consistent (m inim um -energy) solution acquires a nite range. In that case, as a quark in the system is separated from the others, the in uence of the m ean eld on the quark dim inishes and the m ass rises as in (3). The quarks are thus con ned to the region of nonzero m ean eld by virtue of their interaction with the vacuum.

In the description of matter, as the density increases the space between isolated solitons decreases, and them ean eld can acquire nonzero values throughout space. One can conclude based on Eq.(3) that as the vacuum is led by the nonzero mean eld the quarks can sample a larger region of space. Finally, in su ciently dense matter, localized clusters of quarks (nucleons) cease to be energetically favored, and a conducting phase is established.

To discuss the many-soliton problem in more detail and to identify the transition density, we now consider a lattice of static solitons as described above. The relative motion of the

<sup>&</sup>lt;sup>2</sup>This coupling actually arises from chiral symmetry considerations [9].

nucleons is thus neglected in the present treatment and in similar studies. Here we are not interested in reproducing the saturation properties of nuclear matter. Rather, we focus on qualitative changes in the behavior of the system (as a function of the lattice spacing) that can be identified with the hadron-quark transition in the fram ework of the model. A more complete treatment should include the kinetic energy of moving solitons.

As opposed to the single-soliton situation, where the boundary conditions require that the self-consistent solutions of the D irac and K lein-G ordon equations vanish as r ! 1, we have periodic boundary conditions on the lattice, (r) = (r + a) for the meson eld, and similarly for the quark spinors, where a is the lattice vector. As is well known from solid state physics [17], for su ciently sm all values of the lattice constant a = jaj, the sharp energy eigenvalues of the isolated soliton are replaced by sets of closely spaced eigenvalues (energy bands), each soliton contributing one energy level to each band. As the lattice constant goes to in nity, the bands converge to the single-soliton solutions.

Following earlier calculations for soliton matter [1,5], we address the problem in the W igner-Seitz approximation [18]. Introducing a spherical W igner-Seitz cell of radius R permits the decomposition of the quark spinors as

$$u_{n} (\mathbf{r}) = \begin{cases} 0 & 1 \\ B & g_{n} (\mathbf{r}) Y_{J}^{m,J} (\mathbf{\hat{r}}) \\ C & A \\ f_{n} (\mathbf{r}) Y_{J}^{m,J} (\mathbf{\hat{r}}) \end{cases}$$
(4)

where the Y -s are vector spherical harm onics, and prescribes the following boundary conditions at the edge of the W igner-Seitz cell:

$$\frac{\mathrm{d}}{\mathrm{d}\mathbf{r}} (\mathbf{r})_{\mathbf{r}=R} = 0$$
(5)

for the sigma eld, and

$$\frac{d}{dr}g_{n}(r) = f_{n}(r) = 0$$
 (6)

for the upper and low er components of the quark spinors corresponding to the low est-energy state in each energy band [1,5].

To solve the coupled equations for the many-soliton problem, it is advantageous to work with the Fourier expansions of the radial quark wave functions  $g_n$  (r) and  $f_n$  (r). The D irac equation then reduces to a matrix equation in momentum space and the boundary conditions can be easily incorporated. We start by solving for the quark energy eigenvalues in an arbitrary sign a eld. The quark wave functions of the lowest-energy state are then generated, which (after Fourier transform ation) provide the source of the sign a eld as input to the nonlinear K lein-G ordon equation in coordinate space. The D irac equation is solved again in the new sigma eld and the quark wave functions are compared (in quadrature) to the ones obtained in the previous iteration. Dependent on the radius of the W igner-Seitz cell, R, and on the required tolerance, the procedure typically converges in 3-6 iterations. Once the nalsign a eld has been obtained at a given R, the D irac equation in this eld is solved for the energies of the higher excited states. Each D irac spinor carries two di erent orbital quantum numbers (' and  $^{\circ}$ ) and only the total angular momentum j is a good quantum number. In particular, the lowest-energy state has = 0;  $^{0} = 1$ ; j = 1=2, and the next state is characterized by = 1; = 2; j = 3=2. [19] W e will refer to these states as the 1s1=2 and 1p3=2 states, respectively.

The bottom and top energies of the lowest energy bands as a function of the radius of the W igner-Seitz cell for the parameter value = 1:35 GeV in (2) are shown in the F igure. The dienent symbols represent the calculated energies of the bottom of the three lowest energy bands. On one point we indicate a typical uncertainty we associate with our calculation. This uncertainty mainly arises from the freedom in prescribed tolerances at dierent stages of the calculation. The calculated points are tted to smooth curves to guide the eye and to approximately determine the intersection point between the top of the lowest energy band (1s1=2) and the bottom of the next two bands (1p3=2 and 2s1=2). For the top of the energy bands top as a function of the cell radius R we use the simple approximation [1]

$$_{top} = ({}^{2}_{bot} + ({}^{2}_{2R})^{2})^{1=2}$$
; (7)

where  $_{\rm bot}$  is the bottom of the energy band obtained with the boundary conditions (6).

M ore elaborate prescriptions for the top of the bands are also used in the literature [5], but the above simple estimate is consistent with the qualitative nature of the present study. For large R, top ! bot, and the eigenvalues approach those of the isolated soliton. As R is decreased, we rst observe a decrease in the energy of the lowest band. This attraction is a consequence of the boundary condition on the large component of the quark wave function (6), which allows less curvature and therefore less quark kinetic energy than in the case of the isolated soliton. For higher densities, when the solitons and the quark wave functions start to signi cantly overlap, the resulting repulsion overcom es the attraction, and a m inimum develops. Since we have a mean- eld m odel lacking the details of the nucleonnucleon interaction, we do not expect the m inimum to be at the saturation density of nuclear m atter. U pon further increase of the increase of the sigma eld at the cell boundary, which m akes it advantageous for the system to arrange itself in a con guration with nonzero orbital angularm on entum and with quark density peaked away from the origin.

The intersection of di erent bands can be interpreted to signal a transition to a qualitatively di erent high-energy phase. For moderate values of R no bands intersect, and m atter is an insulator. However, as R decreases, the lowest, occupied band intersects the 1p3=2 and 2s1=2 bands. When the energy of an unled level falls below that of the highest occupied state, quarks with this energy become free to migrate throughout the crystal as electrons in a metal, and color conductivity sets in. We follow Ref. [1] and identify the onset of color conductivity with the decon nem ent transition. The value of the critical density  $_{\rm c}$  in the model depends on the assumptions on band lling. [5] Taking the simple approximation of uniform ly led bands and considering the 1s1=2 | 1p3=2 band crossing, with = 1:35 G eV we get  $_{\rm c} = (2:6 \quad 0.2)_{\rm 0}$ , where  $_{\rm 0} = 0:17$  fm  $^{1}$  is the standard nuclear matter density. A ssum ing that the lowest energy band is partially led will increase the critical density obtained in the model.

In the Table we show variations corresponding to changes in the value of the parameter in Eq. (2). The root m can square radius of the soliton for large R (isolated nucleon), the

7

W igner-Seitz radius and density belonging to the m in im um energy, the critical W igner-Seitz radius, and the critical density are displayed for selected values of . The present m odel does not include any m eson dressing, and thus care should be taken when the calculated  $hr^2 i^{1=2}$  is compared to the experimental value of the root m ean square charge radius of the proton,  $hr^2 i_{exp}^{1=2} = 0.83$  fm . W e feel that in this schematic m odel the range selected represents a reasonable interval of variation for the single free parameter .

In summary, we discussed the properties of an in nite system of nonlocal, con ning solitons in the W igner-Seitz approximation. Due to the internally-generated meson elds and the specie c con nement mechanism associated with the model, our study represents a step forward in the description of nuclear matter as a system of solitons. The quark energy bands that develop in the system with decreasing cell size were calculated. We identied the insulator-conductor transition in the crystal with the decon nement transition expected in dense hadronic matter. In studies of the nuclear equation of state it is custom any to use a hadronic equation of state at low energy densities and a bag-model equation of state at high energy densities; the critical parameters are than determined by extrapolating both descriptions to the phase transition region and solving the G ibbs conditions for phase coexistence [20]. The present model is much more satisfactory as it uses the same basic degrees of freedom in both phases and across the phase transition. In the future, the model can also be used to examine in-medium properties such as the dependence of the nucleon mass on the density of the medium.

#### ACKNOW LEDGMENTS

Two of us (C W J. and G F.) are grateful for the partial support and hospitality of the Institute for Nuclear Theory at the University of W ashington. This work was supported in part by the D epartm ent of Energy under G rant Nos. DOE/DE-FG02-86ER-40251 and DE-FG06-90ER-40561.

8

#### REFERENCES

- [1] B. Banerjee, N.K. Glendenning, and V. Soni, Phys. Lett. 155B (1985)213; N.K. Glendenning and B. Banerjee, Phys. Rev. C 34 (1986)1072.
- [2] H. Reinhardt, B.V. Dang, and H. Schulz, Phys. Lett. 159B (1985)161.
- [3] J. Achtzehnter, W. Scheid, and L. Wilets, Phys. Rev. D 32 (1985)2414.
- [4] Q. Zhang, C. Derreth, A. Schafer, and W. Greiner, J. Phys. G: Nucl. Phys. 12 (1986) L19.
- [5] M C. Birse, J.J. Rehr, and L. W ilets, Phys. Rev. C 38 (1988)359.
- [6] G.Fai, R. Perry, and L. W ilets, Phys. Lett. 208B (1988)1.
- [7] R.J. Cahill, Nucl. Phys. A 543 (1992)63.
- [8] M. R. Frank in Few-Body Problems in Physics, Ed., Franz Gross (AIP Conference Proceedings 334) pp. 15-30.
- [9] R.T. Cahill and C.D. Roberts, Phys. Rev. D 32 (1985)2419; C.D. Roberts, R.T. Cahill and J. Praschifka, Ann. Phys. (N.Y.) 188 (1988)20.
- [10] C D. Roberts, R.T. Cahill, M E. Sevior, and N. Iannella, Phys. Rev. D 49 (1994)125;M R. Frank and T. Meissner, Phys. Rev. C 53 (1996)2410.
- [11] M R.Frank and P.C.Tandy, Phys.Rev.C 49 (1994)478; C D.Roberts, E lectrom agnetic Pion Form Factor and Neutral Pion Decay W idth, ANL preprint ANL-PHY-7842-TH-94 (1994); M R, Frank, K L.M itchell, C D.Roberts, and P.C.Tandy, Phys.Lett.B 359 (1995)17; C.J.Burden, C.D.Roberts and M.J.Thom son, E lectrom agnetic Form Factors of Charged and Neutral K aons, ANL preprint ANL-PHY-8240-TH-95 (1995).
- [12] M R.Frank, Phys. Rev. C 51 (1995) 987.
- [13] M R. Frank and C D. Roberts, Physical Review C 53 (1996) 390.
- [14] M R. Frank, P.C. Tandy, and G. Fai, Phys. Rev. C 43 (1991)2808; P.C. Tandy and

M R. Frank, Aust. J. Phys. 44 (1991)181; M R. Frank and P.C. Tandy, Phys. Rev. C 46 (1992)338.

- [15] R.T.Cahill and SM.Gunner, The global colour model of QCD and its relationship to the NJL model, chiral perturbation theory and other models, hep-ph/9601319 (1996).
- [16] C D. Roberts and A G. W illiams, Dyson-Schwinger equations and their application to hadronic physics, in Progress in Particle and Nuclear Physics, Ed., A. Faessler (Pergam on Press, Oxford 1994); and references therein.
- [17] C.K ittel, Introduction to Solid State Physics, Sixth Edition, W iley, 1986.
- [18] E.W igner and F. Seitz, Phys. Rev. 43 (1933)804; 46 (1934)509.
- [19] R K. Bhaduri, M odels of the Nucleon, Addison-W esley, 1988.
- [20] V K.Mishra, G.Fai, LP.C semai, and E.O snes, Phys. Rev.C 47 (1993)1519.

## Figure caption

The bottom and top energies of the lowest energy bands of the soliton lattice as a function of the radius of the W igner-Seitz cell for = 1.35 GeV. The sym bols represent the calculated energies of the bottom of the three lowest energy bands. For the top of the energy bands the approximation (7) is used.

### Table caption

Variations of the root m ean square radius of a single soliton, the W igner-Seitz radius and density belonging to the m inimum energy, the critical W igner-Seitz radius, and the critical density with the single m odel parameter  $\cdot$ .

(GeV)	1.25		1.35		1.45	
$hr^2i^{1=2}$ (fm )	.71		<b>.</b> 67		<b>.</b> 64	
R <sub>min</sub> (ffm)	1.33	.02	126	.02	1.17	.02
min = 0	<b>.</b> 60	.03	.70	.04	.88	.05
R <sub>c</sub> (ffm)	<b>.</b> 86	.02	<b>.</b> 82	.02	.77	.02
c= 0	22	2	2.6	2	3.1	З

