

Strong decays of nonstrange q^3 baryons

R. Bijker^a, F. Tachello^b and A. Leviatan^c

^aInstituto de Ciencias Nucleares, U N A M ., A P . 70-543, 04510 M exico, D F ., M exico

^bCenter for Theoretical Physics, Sloane Laboratory, Yale University,
New Haven, CT 06520-8120, U S A .

^cRacah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

Abstract

We study strong decays of nonstrange baryons by making use of the algebraic approach to hadron structure. Within this framework we derive closed expressions for decay widths in an elementary-meson emission model and use these to analyze the experimental data for $N \rightarrow N + \pi$, $N \rightarrow N + \eta$, $N \rightarrow N + \omega$, $N \rightarrow N + \rho$, $N \rightarrow N + \sigma$ and $N \rightarrow N + \delta$ decays.

PACS numbers: 13.30.Eg, 11.30.Na, 14.20.Gk

I. INTRODUCTION

A large amount of experimental data was accumulated in the 1960's and 1970's on the spectroscopy of light hadrons. These data led first to the introduction of $SU_f(3)$ by Gell'Mann [1] and Ne'eman [2] (later enlarged to $SU_{sf}(6)$ by Gursey and Radicati [3]), and subsequently to $SU_c(3)$ color symmetry as a gauge symmetry of strong interactions. The construction of dedicated facilities (e.g. CEBAF, MAMI, ELSA) that promise to produce new and more accurate data on the same subject, have stimulated us to revisit baryon spectroscopy with the intent to reexamine whether or not the new data can shed some new light on the structure of hadrons. In this reanalysis we have introduced, in addition to the basic spin-flavor-color symmetry, $SU_{sf}(6) \times SU_c(3)$, a new ingredient, namely a space symmetry, G , which we have taken to be $G = U(7)$ for baryons [4]. The introduction of the space symmetry allows us to examine, in a straightforward way, several limiting situations (e.g. harmonic oscillator and collective dynamics), and to produce transparent results that can be used to analyze the experimental data. This approach has been used [4,5] to analyze the mass spectrum and electromagnetic couplings of nonstrange baryon resonances. It presents an alternative for the use of nonrelativistic or relativized Schrodinger equations. In addition to electromagnetic couplings, strong decays of baryons provide an important, complementary, tool to study the structure of baryons. These strong couplings are needed to analyze the new upcoming experimental data, especially $(e;e^0)$ and $(e;e^0)$. Furthermore, we want to understand whether or not unusual features appear in the data, which may point out to 'new' physics, new meaning here unconventional configurations of quarks and gluons, such as hybrid states, $q^3 g$, or multiquark states, $q^3 qq$. For example, the observed large width of the $N(1535)S_{11}$ resonance has led to considerable discussion [6-10] about the nature of this resonance.

The strong decays have been analyzed previously in the nonrelativistic [11] and relativized quark models [12]. These models emphasize single-particle aspects of quark dynamics in which only a few low-lying configurations in the confining potential contribute significantly to the baryon wave function. In the framework of the earlier mentioned algebraic approach, it is possible to study also other, more collective, types of dynamics. In this article, we analyze in detail the strong decays in a collective model of baryon structure. The article is organized as follows: in Section II we briefly discuss the method of calculation, in Section III we present the results, which are compared with the existing data in Section IV. Finally, in Section V we present our conclusions and point out some open problems.

II. METHOD OF CALCULATION

We consider in this article strong decays of baryons by the emission of a pseudoscalar meson

$$B \rightarrow B^0 + M : \quad (1)$$

In order to calculate these decays we need two ingredients: (i) the wave function of the initial and final states and (ii) the form of the transition operator. We write both the wave functions and the transition operators in algebraic form. We take the wave functions as representations of $U(7) \supset SU_{sf}(6) \supset SU_c(3)$, as discussed in [4]. The algebraic formulation allows us to study a large class of models, all with the same spin-avor-color structure, but different types of quark dynamics (e.g. single-particle and collective). Each scenario corresponds to different ways in which the $U(7)$ spatial symmetry is broken down to the angular momentum group $SO(3)$. Among the models in this class is the familiar harmonic oscillator quark model characterized by the breaking

$$U(7) \supset U(6) \supset U(3) \supset U(3) \supset SO(3) \supset SO(3) \supset SO(3) \supset SO(2); \quad (2)$$

where the index α and β refers to the two relative Jacobi coordinates

$$\begin{aligned} \vec{r}_\alpha &= \frac{1}{\sqrt{2}} (\vec{r}_1 - \vec{r}_2); \\ \vec{r}_\beta &= \frac{1}{\sqrt{6}} (\vec{r}_1 + \vec{r}_2 - 2\vec{r}_3); \end{aligned} \quad (3)$$

The nonrelativistic and relativized quark potential models [13,14] are formulated in this harmonic oscillator basis.

In a collective model the baryon resonances are interpreted in terms of rotations and vibrations of the string configuration in Fig. 1, which is characterized by the two relative Jacobi coordinates of Eq. (3). The three constituent parts carry the global quantum numbers: avor = triplet = u, d, s ; spin = doublet = $1/2$; and color = triplet (we do not consider here heavy quark avors). Different types of collective models are specified by a distribution of the charge, magnetic moment, etc., over the entire volume. For the present analysis of strong decays we use the (normalized) distribution

$$g(r) = \frac{4}{\pi^{3/2}} e^{-r^2/a^2} = 2a^3; \quad (4)$$

where r is a radial coordinate and a is a scale parameter. We have shown in [5] that the above distribution appears to describe the available data on electromagnetic form factors and helicity amplitudes up to $Q^2 \approx 10 - 20 \text{ (GeV/c)}^2$.

The spatial part of the baryon wave functions in the 'collective' model is characterized by the following labels: $(v_1; v_2); K; L_t^P$ [4], where $(v_1; v_2)$ denotes the vibrations (stretching and bending) of the configuration of Fig. 1; K denotes the projection of the rotational angular momentum L on the body-fixed axis; P the parity and t the symmetry type of the state under the point group D_3 . The classification under D_3 and parity is equivalent to that of the point group D_{3h} , which describes the discrete symmetry of the object of Fig. 1. Instead of the D_3 labels ($t = A_1, A_2, E$) one can use the labels of S_3 ($t = S, A, M$), the group of permutations of the three constituent parts (S_3 and D_3 are isomorphic). The permutation symmetry t of the spatial part of the wave function must be the same as that of the spin-avor part in order to have total wave functions that are antisymmetric (the color part is a

color-singlet, i.e. antisymmetric). Therefore one can also use the dimension of the SU_{sf} (6) representations to label the states: $A_1 \sim S \sim 56$, $A_2 \sim A \sim 20$ and $E \sim M \sim 70$. We adhere to the latter notation and label the states by

$$[\dim fSU_{sf}(6)g; L^P]_{(v_1, v_2), K}^E : \quad (5)$$

When the spin-flavor quantum numbers are explicitly added, we obtain the total baryon wave function

$$^{2S+1} \dim fSU_f(3) g_J [\dim fSU_{sf}(6)g; L^P]_{(v_1, v_2), K}^E ; \quad (6)$$

where S and J are the spin and total angular momentum $J = L + S$. For example, in this notation the nucleon and delta wave functions are given by

$$^2 8_{1=2} [56; 0^+]_{(0,0),0}^E \quad \text{and} \quad ^4 10_{3=2} [56; 0^+]_{(0,0),0}^E ; \quad (7)$$

respectively.

The second ingredient in the calculation is the form of the operator inducing the strong transition. Several forms have been suggested [15]. We use here the simple form [11]

$$H = \frac{1}{(2)^{3=2} (2k_0)^{1=2}} X_j^M \sum_{j=1}^h 2g(s_j \cdot \hat{k}) e^{i\hat{k} \cdot \mathbf{r}_j} + h s_j \cdot (\mathbf{p}_j e^{i\hat{k} \cdot \mathbf{r}_j} + e^{i\hat{k} \cdot \mathbf{r}_j} \mathbf{p}_j) ; \quad (8)$$

where \mathbf{r}_j , \mathbf{p}_j and \mathbf{s}_j are the coordinate, momentum and spin of the j -th constituent, respectively; $k_0 = E_M = E_B - E_{B^0}$ is the meson energy, and $\hat{k} = \hat{P}_M = \hat{P} - \hat{P}^0 = k\hat{z}$ denotes the momentum carried by the outgoing meson. Here $\hat{P} = P_z \hat{z}$ and $\hat{P}^0 (= P_z^0 \hat{z})$ are the momenta of the initial and final baryon. The coefficients g and h denote the strength of the two terms in the transition operator of Eq. (8). The flavor operator X_j^M (to be discussed below) corresponds to the emission of an elementary meson (M) by the j -th constituent: $q_j \rightarrow q_j^0 + M$ (see Figure 2).

Using the symmetry of the wave functions for nonstrange baryons (the only case we discuss here), transforming to Jacobi coordinates and integrating over the baryon center of mass coordinate, we find

$$H = \frac{1}{(2)^{3=2} (2k_0)^{1=2}} 6X_3^M \left[g k s_{3,z} \hat{U} + h s_{3,z} (\hat{T}_z - \frac{1}{6} (P_z + P_z^0) \hat{U}) - \frac{1}{2} h (s_{3,+} \hat{T}_+ + s_{3,-} \hat{T}_-) \right] ; \quad (9)$$

with

$$\begin{aligned} \hat{U} &= e^{ik \frac{P_z}{3} z} ; \\ \hat{T}_m &= \frac{1}{2} \left(\frac{2}{3} p_m e^{ik \frac{P_z}{3} z} + e^{ik \frac{P_z}{3} z} \frac{2}{3} p_m \right) A : \end{aligned} \quad (10)$$

By using momentum conservation $P_z + P_z^0$ can be written as $2P_z - k$. The dependence on the overall center of mass momentum is spurious and must be eliminated. (In the rest frame

of B , $P_z = 0$ and the spurious term is automatically absent.) By making the replacement, $\mathbf{p} \rightarrow \mathbf{p}_q$ [16], and the mapping onto the algebraic operators, $2=3_m$! $\hat{D}_{3m} = X_D$ [4,5], we can write Eq. (10) as

$$\begin{aligned}\hat{U} &= e^{ik \cdot \hat{D}_{3m} / z = X_D} ; \\ \hat{T}_m &= \frac{im_q k_0}{2X_D} \hat{D}_{3m} e^{ik \cdot \hat{D}_{3m} / z = X_D} + e^{ik \cdot \hat{D}_{3m} / z = X_D} \hat{D}_{3m} : \end{aligned} \quad (11)$$

The dipole operator \hat{D}_{3m} is a generator of $U(7)$ and X_D is its normalization, as discussed in [4,5]. The calculation of the matrix elements of H can be done in configuration space (\mathbf{r}, \mathbf{r}) or in momentum space (\mathbf{p}, \mathbf{p}) . The mapping onto the algebraic space of $U(7)$ is a convenient way to carry out the calculations, much in the same way as the mapping of coordinates and momenta onto creation and annihilation operators in the harmonic oscillator space.

In the collective model the matrix elements of H are obtained by folding with the distribution function $g(\mathbf{r})$ of Eq. (4). These collective matrix elements can be expressed in terms of helicity amplitudes. For decays in which the initial baryon has angular momentum $J = L + S$ and in which the final baryon is either the nucleon or the delta with $L^0 = 0$ and thus $J^0 = S^0$, the helicity amplitudes are

$$\begin{aligned}A(\mathbf{k}) &= \int d\mathbf{r} g(\mathbf{r}) h^0; L^0 = 0; S^0; J^0 = S^0; M_J^0 = -j; H; j; L; S; J; M_J = -i \\ &= \frac{1}{(2\pi)^{3/2} (2k_0)^{1/2}} hL; 0; S; -j; i_0 Z_0(\mathbf{k}) + \frac{1}{2} hL; 1; S; -1; j; i_+ Z_+(\mathbf{k}) \\ &\quad + \frac{1}{2} hL; 1; S; +1; j; i_- Z_-(\mathbf{k}) : \end{aligned} \quad (12)$$

Here i_\pm denotes the labels that, in addition to L, S, J and \mathbf{r} , are needed to specify the baryon wave function (see Eq. (6)). The coefficients m are the spin-flavor matrix elements of $X_3^M s_{3m}$, to be discussed below, and $Z_m(\mathbf{k})$ ($m = 0; \pm$) are the radial matrix elements

$$\begin{aligned}Z_0(\mathbf{k}) &= 6 \int d\mathbf{r} g(\mathbf{r}) h^0; L^0 = M_L^0 = 0; j; k; \hat{U} = h(\hat{T}_z - \frac{1}{6}(P_z + P_z^0)\hat{U}) j; L; M_L = 0; i \\ &= 6 [jk + h \frac{1}{6}(P_z + P_z^0)] F(\mathbf{k}) - 6h G_z(\mathbf{k}) ; \\ Z_+(\mathbf{k}) &= 6h \int d\mathbf{r} g(\mathbf{r}) h^0; L^0 = M_L^0 = 0; j; \hat{T} = j; L; M_L = -1; i \\ &= 6h G_-(\mathbf{k}) : \end{aligned} \quad (13)$$

The calculation of the radial matrix elements is identical to that already reported for electromagnetic couplings [4,5]. Therefore, we do not repeat it here, but only show in Table I the matrix elements $F(\mathbf{k})$ and $G_m(\mathbf{k})$ of \hat{U} and \hat{T}_m , respectively, for the 'collective' model with distribution given by Eq. (4). Note that Table I presents the results for an emission process, whereas the corresponding table for the 'collective' model in [4,5] shows the results for an absorption process. For any other model of baryons with the same spin-flavor structure, the

corresponding results can be obtained by replacing Table I with the appropriate table (for example, by using harmonic oscillator wave functions as discussed in [4]).

Contrary to the case of electromagnetic couplings where the spin- flavor part is relatively simple, the calculation of the spin- flavor part for strong decays is somewhat more involved. The spin- flavor matrix elements factorize into a spin matrix element of $S_{3\mu}$ between spin wave functions and a flavor matrix element of X_3^M between flavor wave functions (λ and λ'). The calculation of the spin part is straightforward. For the flavor part we take the flavor operators of the form

$$\begin{aligned} X^+ &= \frac{1}{2} (\lambda_1 - i\lambda_2) ; \\ X^0 &= \lambda_3 ; \\ X^- &= \frac{1}{2} (\lambda_1 + i\lambda_2) ; \\ X^8 &= \frac{1}{\sqrt{3}} \lambda_8 ; \\ X^1 &= \frac{2}{3} I ; \end{aligned} \quad (14)$$

where λ_i are the Gell-Mann matrices [17] and I denotes the unit operator in flavor space. For the pseudoscalar mesons we introduce a mixing angle θ_P between the octet and singlet mesons [17]

$$\begin{aligned} \phi &= \lambda_8 \cos \theta_P - \lambda_1 \sin \theta_P ; \\ \omega &= \lambda_8 \sin \theta_P + \lambda_1 \cos \theta_P ; \end{aligned} \quad (15)$$

and similarly for the corresponding flavor operators

$$\begin{aligned} X^+ &= X^8 \cos \theta_P - X^1 \sin \theta_P ; \\ X^0 &= X^8 \sin \theta_P + X^1 \cos \theta_P ; \end{aligned} \quad (16)$$

The flavor operators X_3^M appearing in Eq. (9) only act on the 3-rd constituent. The corresponding matrix elements can either be evaluated explicitly for each channel separately, or more conveniently, by using the Wigner-Eckart theorem and isoscalar factors of SU_3 [18]. For the decay process $B \rightarrow B^0 + M$ we have

$$\begin{aligned} \langle \lambda' M' | X_3^M | \lambda M \rangle &= \langle (\lambda_2; q_2); I_2; M_2; Y_2 | T^{(\lambda_2 q_2) I_2 Y_2} | (\lambda_1; q_1); I_1; M_1; Y_1 \rangle + \\ &= \langle \lambda_1; M_1; I_1; M_1 | T^{(\lambda_1 q_1) I_1 Y_1} | \lambda; M; I; M \rangle \langle (\lambda_2; q_2); I_2; Y_2 | T^{(\lambda_2 q_2) I_2 Y_2} | (\lambda_1; q_1); I_1; Y_1 \rangle \\ &= \langle (\lambda_2; q_2); I_2; Y_2 | T^{(\lambda_2 q_2) I_2 Y_2} | (\lambda_1; q_1); I_1; Y_1 \rangle \langle \lambda; M; I; M | T^{(\lambda q) I Y} | \lambda; M; I; M \rangle ; \end{aligned} \quad (17)$$

In this notation $(\lambda_i; q_i) = (1; 1)$ or $(3; 0)$ for the baryon flavor octet and decuplet, respectively, and $(p; q) = (1; 1)$ or $(0; 0)$ for the meson flavor octet and singlet, respectively (see Eq. (14)). The sum over i is over different multiplicities. The flavor states are labeled by the quantum

numbers $(p; q; I; M_I; Y)$ corresponding to the reduction $SU_f(3) \rightarrow SU_I(2) \times U_Y(1)$. The right hand side contains the sum of products of an isospin Clebsch-Gordan coefficient, which contains the dependence on the charge channel, a $SU(3)$ isoscalar factor, which depends on the isospin channel, and a $SU(3)$ reduced matrix element, which depends on the coupling of the Λ baryon multiplets $(p; q)$, but not on isospin $I; M_I$ and hypercharge Y . In Table II we give the expressions for the $SU_f(3)$ reduced matrix elements.

Using Eqs. (14)–(17) and the matrix elements of the spin operator, we can compute all spin- Λ baryon matrix elements for a given isospin channel. In Tables III and IV we present the results for the decay into Λ and Λ^0 , respectively. Results for a specific charge channel can be obtained by multiplying with the appropriate isospin Clebsch-Gordan coefficient.

The helicity amplitudes $A_l(k)$ of Eq. (12) can be converted to partial wave amplitudes $a_l(k)$ by [19]

$$\begin{aligned} A_l(k) &= \sum_{l=L-1}^L \sum_p \frac{1}{2l+1} \langle l; 0; J^0; -j; i | A_l(k) \rangle; \\ a_l(k) &= \frac{1}{2J+1} \sum_p \frac{1}{2l+1} \langle l; 0; J^0; -j; i | A_l(k) \rangle; \end{aligned} \quad (18)$$

Here l is the relative orbital angular momentum between the Λ baryon and the emitted meson. It takes the values $l = L-1$ (the value $l = L$ is not allowed because of parity conservation). With the definition of the transition operator in Eq. (8) and the helicity amplitudes and partial wave amplitudes, the decay widths for a specific isospin channel are given by [15]

$$\begin{aligned} \Gamma(B \rightarrow B^0 + M) &= 2 \int_{\Omega} \frac{2}{2J+1} \sum_{l=L-1}^L |A_l(k)|^2 \\ &= 2 \int_{\Omega} \sum_{l=L-1}^L |a_l(k)|^2; \end{aligned} \quad (19)$$

where \int_{Ω} is a phase space factor.

III. RESULTS

For all resonances with the same value of $(v_1; v_2); L^P$ the expression for the decay widths of Eq. (19) can be rewritten in a more transparent form in terms of only two elementary partial wave amplitudes $W_l(k)$,

$$\Gamma(B \rightarrow B^0 + M) = 2 \int_{\Omega} \sum_{l=L-1}^L |a_l(k)|^2 = 2 \int_{\Omega} \frac{1}{(2\pi)^3 2k_0} \sum_{l=L-1}^L c_l^2 |W_l(k)|^2; \quad (20)$$

For this set of resonances, the k dependence of the partial wave amplitudes $a_l(k)$ is contained in the amplitudes $W_l(k)$, while the dependence on the individual baryon resonance is contained in the coefficients c_l . In the algebraic method, the elementary partial wave amplitudes $W_l(k)$ can be obtained in closed form.

In Table V we present the values of c_l for the negative parity resonances with $(v_1; v_2); L^P = (0; 0); 1^-$. In the 'collective' model with distribution given by Eq. (4) the corresponding S and D elementary partial wave amplitudes are

$$\begin{aligned} W_0(k) &= i \left[gk - \frac{1}{6}hk \right] \frac{ka}{(1 + k^2 a^2)^2} + h m_3 k_0 a \frac{3 k^2 a^2}{(1 + k^2 a^2)^3} ; \\ W_2(k) &= i \left[gk - \frac{1}{6}hk \right] \frac{ka}{(1 + k^2 a^2)^2} - h m_3 k_0 a \frac{4k^2 a^2}{(1 + k^2 a^2)^3} : \end{aligned} \quad (21)$$

Similarly, in Table VI we present the c_l coefficients for the positive parity resonances with $(v_1; v_2); L^P = (0; 0); 2^+$. The corresponding P and F elementary partial wave amplitudes are

$$\begin{aligned} W_1(k) &= \left[gk - \frac{1}{6}hk \right] \left(\frac{1}{(1 + k^2 a^2)^2} + \frac{3}{2k^3 a^3} H(ka) \right) \\ &\quad + h m_3 k_0 a \frac{1}{ka} \frac{4k^2 a^2}{(1 + k^2 a^2)^3} ; \\ W_3(k) &= \left[gk - \frac{1}{6}hk \right] \left(\frac{1}{(1 + k^2 a^2)^2} + \frac{3}{2k^3 a^3} H(ka) \right) \\ &\quad + h m_3 k_0 a \frac{1}{ka} \left(\frac{5 + 9k^2 a^2}{(1 + k^2 a^2)^3} - \frac{15}{2k^3 a^3} H(ka) \right) ; \end{aligned} \quad (22)$$

with $H(ka) = \arctan ka$ $ka = (1 + k^2 a^2)^{1/2}$.

Partial widths for other models of the nucleon and its resonances can be obtained by introducing the corresponding expressions for the elementary amplitudes $W_l(k)$. For example, the relevant expressions in the harmonic oscillator quark model are

$$\begin{aligned} W_0(k) &= \frac{i}{3} \left[gk - \frac{1}{6}hk \right] k + h m_3 k_0 \left(3 - \frac{k^2 a^2}{3} \right) e^{-k^2 a^2/6} ; \\ W_2(k) &= \frac{i}{3} \left[gk - \frac{1}{6}hk \right] k - \frac{1}{3} h m_3 k_0 k^2 a^2 e^{-k^2 a^2/6} ; \end{aligned} \quad (23)$$

and

$$\begin{aligned} W_1(k) &= \frac{p}{3} \frac{\sqrt{2}}{15} k \left[gk - \frac{1}{6}hk \right] k + h m_3 k_0 \left(5 - \frac{k^2 a^2}{3} \right) e^{-k^2 a^2/6} ; \\ W_3(k) &= \frac{p}{3} \frac{\sqrt{2}}{15} k \left[gk - \frac{1}{6}hk \right] k - \frac{1}{3} h m_3 k_0 k^2 a^2 e^{-k^2 a^2/6} : \end{aligned} \quad (24)$$

IV. ANALYSIS OF EXPERIMENTAL DATA

Use of Eqs. (20)-(22) allows us to do a straightforward and systematic analysis of the experimental data. Here we adopt the procedure of [15], in which calculations are performed in the rest frame of the decaying resonance, and in which the relativistic expression for the

phase space factor ϕ_f as well as for the momentum k of the emitted meson are retained. The expressions for k and ϕ_f are

$$k^2 = \frac{m_B^2}{4} + \frac{(m_B^2 - m_{B^0}^2 + m_M^2)^2}{4m_B^2};$$

$$\phi_f = \frac{dP^0 dP_M}{m_B} (m_B - E_{B^0}(P_M) - E_M(P_M)) (P^0 + P_M)$$

$$= 4 \frac{E_{B^0}(k) E_M(k) k}{m_B} \quad (25)$$

with $E_{B^0}(k) = \sqrt{m_{B^0}^2 + k^2}$ and $E_M(k) = \sqrt{m_M^2 + k^2}$.

We consider here decays with emission of π and η . The experimental data, extracted from the compilation by the Particle Data Group [17] are shown in Tables V II, V III and IX, where they are compared with the results of our calculation. The calculated values depend on the two parameters g and h in the transition operator of Eq. (8), and on the scale parameter a of Eq. (4). We keep g , h and a equal for all resonances and all decay channels (N^* , N^* , Σ^* , Λ^*). In comparing with previous calculations, it should be noted that in the calculation in the nonrelativistic quark model of [11] the decay widths are parametrized by four reduced partial wave amplitudes instead of the two elementary amplitudes g and h . Furthermore, the momentum dependence of these reduced amplitudes are represented by constants. The calculation in the relativized quark model of [12] was done using a pair-creation model for the decay and involved a different assumption on the phase space factor. Both the nonrelativistic and relativized quark model calculations include the effects of mixing induced by the hyperfine interaction, which in the present calculation are not taken into account.

In the present analysis we determine the values of g , h and a from a least square fit to the N^* partial widths (which are relatively well known) with the exclusion of the S_{11} resonances. For the latter the situation is not clear due to possible mixing of $N(1535)S_{11}$ and $N(1650)S_{11}$ and the possible existence of a third S_{11} resonance [9]. As a result we find $g = 1.164 \text{ GeV}^{-1}$ and $h = 0.094 \text{ GeV}^{-1}$. The relative sign is consistent with a previous analysis of the strong decay of mesons [20] and with a derivation from the axial-vector coupling (see e.g. [15]). The scale parameter, $a = 0.232 \text{ fm}$, extracted in the present fit is found to be equal to the value extracted in the calculation of electromagnetic couplings [5]. Just as in our study of the electromagnetic couplings [5] we present the strong decays of the resonances, that in the 'collective' model are assigned as vibrational excitations of the configuration of Fig. 1, in terms of two coefficients, $\alpha_1 = (1 - R^2)/R = \frac{p}{N}$ and $\alpha_2 = \frac{1}{1 + R^2} = \frac{p}{N}$, one for each fundamental vibration (here N determines the size of the model space and R^2 is a size parameter, as discussed in [4]).

The calculation of decay widths into the N^* channel, as shown for the 3 and 4 star resonances in Table V II, is in fair agreement with experiment. This is emphasized in Figs. 3 and 4. The results are to a large extent a consequence of spin-flavor symmetry. The use of 'collective' form factors improves somewhat the results when compared with older (harmonic oscillator) calculations. This is shown in Table X where the decay of a Regge trajectory

into N is analyzed and compared with the calculations of [15], which are based on the harmonic oscillator model discussed in [21]. We also include the results of more recent calculations in the nonrelativistic quark model [11] and in the relativized quark model [12]. There does not seem to be anything unusual in the decays into π and our analysis confirms the results of previous analyses. In Table XI we show the N decays of the calculated resonances below 2 GeV which have not been observed (missing resonances). The resonances with $L^P = 1^+$ and $L^P = 2^-$ are decoupled because of the spin-flavor symmetry. Most of other resonances have small decay widths, with the exception of the P_{13} and F_{15} states, which have widths comparable to those of the well-established P_{13} and F_{15} states of Table VII. This behaviour can be understood by inspection of Table VI. The coefficients for the P_{13} states are $c_1 = 25/12$ and $c_1 = 2$. This leads to comparable N widths of 31 MeV and 56 MeV, respectively. The difference is due to the k values of these resonances that enter in the elementary partial wave amplitude $W_1(k)$ of Eq. (22). A similar situation holds for the F_{15} states. The relative magnitude of the c_1 coefficients in Table VI explain the smaller decay widths for the other missing resonances of Table XI.

Contrary to the decays into π , the decay widths into η have some unusual properties. The calculation gives systematically small values for these widths. This is due to a combination of phase space factors and the structure of the transition operator. Both depend on the momentum transfer k . The values of k in Tables VII, VIII and IX show that, due to the difference between the π and η mass, the momentum carried by the η is smaller than that carried by the π . Therefore, the η decay widths are suppressed relative to the π decays. The spin-flavor part is approximately the same for N and N , since π and η are in the same $SU_f(3)$ multiplet. We emphasize here, that the transition operator was determined by fitting the coefficients g and h to the N decays of the 3 and 4 star resonances. Hence the η decays are calculated without introducing any further parameters.

The experimental situation is unclear. The 1992 PDG compilation [22] gave systematically small widths (< 1 MeV) for all resonances except $N(1535)S_{11}$. The 1994 PDG compilation [23] deleted all widths with the exception of $N(1535)S_{11}$. This situation persists in the latest PDG compilation [17], where $N(1650)S_{11}$ is now assigned a small but non-zero width. The results of our analysis suggest that the large width for the $N(1535)S_{11}$ is not due to a conventional q^3 state. One possible explanation is the presence of another state in the same mass region, e.g. a quasi-bound meson-baryon S wave resonance just below or above threshold, for example N , K or K [8]. Another possibility is an exotic configuration of four quarks and one antiquark ($q^4\bar{q}$).

For possible use in the analysis of new experimental data, we give in Table XI the strong decay widths of the so-called missing resonances with a calculated mass below 2 GeV which up to now have not been observed experimentally.

V. CONCLUSIONS

We have presented a calculation of the strong decay widths $N \rightarrow N + \pi$, $N \rightarrow N + \eta$, $N \rightarrow N + \rho$, $\Lambda \rightarrow N + \pi$, $\Lambda \rightarrow N + \eta$ and $\Lambda \rightarrow N + \rho$ in a collective model of baryons. By exploiting the symmetry of the problem, both in its spin-color part, $SU_{sf}(6) \supset SU_c(3)$, and in its space part, $U(7)$, we have been able to write the results in a transparent analytic way (Section III). The analysis of experimental data shows that, while the decays into π follow the expected pattern, the decays into η have some unusual features. Our calculations do not show any indication for a large width, as is observed for the $N(1535)S_{11}$ resonance. The observed large width indicates the presence of another configuration, which is outside the present model space. This suggests, that in order to elucidate this point, particular attention be paid at CEBAF to the $N \rightarrow N + \eta$ channel.

Our calculations can be easily extended to include other decay channels, such as K and \bar{K} . These calculations are currently underway and are part of the extension of the model to include strange resonances as well.

ACKNOWLEDGEMENTS

We wish to thank T.-S.H. Lee and D. Kurath for interesting discussions. This work is supported in part by CONACYT, Mexico under project 400340-5-3401E, DGAPA-UNAM under project IN105194 (R.B.), by D.O.E. Grant DE-FG 02-91ER 40608 (F.I.) and by grant No. 94-00059 from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel (A.L.).

REFERENCES

- [1] M . Gell-M ann, Phys. Rev. 125, 1067 (1962).
- [2] Y . Ne'eman, Nucl. Phys. 26, 222 (1961).
- [3] F . G ursey and L A . R adicati, Phys. Rev. Lett. 13, 173 (1964).
- [4] R . B iker, F . Iachello and A . Leviatan, Ann. Phys. (N .Y .) 236, 69 (1994).
- [5] R . B iker, F . Iachello and A . Leviatan, preprint nucl-th/9510001, Phys. Rev. C , in press.
- [6] M . Benm errouche and N C . M ukhopadhyay, Phys. Rev. Lett. 67, 1070 (1991); M . Benm errouche, N C . M ukhopadhyay and J F . Zhang, Phys. Rev. D 51, 3237 (1995); N C . M ukhopadhyay, J F . Zhang and M . Benm errouche, Phys. Lett. B 364, 1 (1995).
- [7] Z . Li, Phys. Rev. D 52, 4961 (1995).
- [8] N . K aiser, P B . S iegel and W . W eise, Phys. Lett. B 362, 23 (1995).
- [9] Z . Li and R . W orkman, Phys. Rev. C 53, R 549 (1996).
- [10] L . Ya . G lozman and D O . R iska, Phys. Lett. B 366, 305 (1996).
- [11] R . K oniuk and N . Isgur, Phys. Rev. Lett. 44, 845 (1980); Phys. Rev. D 21, 1868 (1980).
- [12] S . Capstick and W . Roberts, Phys. Rev. D 47, 1994 (1993); *ibid.* D 49, 4570 (1994).
- [13] N . Isgur and G . Karl, Phys. Rev. D 18, 4187 (1978); D 19, 2653 (1979); D 20, 1191 (1979).
- [14] S . Capstick and N . Isgur, Phys. Rev. D 34, 2809 (1986).
- [15] A . Le Yaouanc, L . O liver, O . Pene and J .C . Raynal, Hadron transitions in the quark model', G ordon and B reach (1988).
- [16] R . M cClary and N . Byers, Phys. Rev. D 28, 1692 (1983).
- [17] Particle D ata G roup, Phys. Rev. D 54, 1 (1996).
- [18] J .J . de Swart, Rev. M od. Phys. 35, 916 (1963).
- [19] W . P . Petersen and J .L . Rosner, Phys. Rev. D 6, 820 (1972).
- [20] C . G obbi, F . Iachello and D . K usnezov, Phys. Rev. D 50, 2048 (1994).
- [21] R . H . D alitz, in Quarks and Hadronic Structure', Ed. M . M ompurgo, Plenum (1977).
- [22] Particle D ata G roup, Phys. Rev. D 45, S1 (1992).
- [23] Particle D ata G roup, Phys. Rev. D 50, 1173 (1994).

TABLES

TABLE I. Matrix elements $F(k)$ and $G(k)$ in the collective model for $N = 1$ (large model space). The final state is $[56; 0^+]_{(0;0);0}$. The matrix elements for the vibrational excitations are given in terms of $\eta_1 = (1 - R^2)/R = N$ and $\eta_2 = \frac{1}{1 + R^2} = \frac{1}{N}$ [4,5]. $H(x) = \arctan x$, $x = (1 + x^2)^{-1/2}$.

Initial state	$F(k)$	$G_z(k) = m_3 k_0 a$	$G(k) = m_3 k_0 a$
$[56; 0^+]_{(0;0);0}$	$\frac{1}{(1 + k^2 a^2)^2}$	$\frac{4ka}{(1 + k^2 a^2)^3}$	0
$[20; 1^+]_{(0;0);0}$	0	0	0
$[70; 1^-]_{(0;0);1}$	$i \frac{1}{2} \frac{3ka}{(1 + k^2 a^2)^2}$	$i \frac{1}{2} \frac{3k^2 a^2}{(1 + k^2 a^2)^3}$	$i \frac{1}{2} \frac{1}{(1 + k^2 a^2)^2}$
$[56; 2^+]_{(0;0);0}$	$\frac{1}{2} \frac{1}{(1 + k^2 a^2)^2} + \frac{3}{2k^3 a^3} H(ka)$	$\frac{1}{2} \frac{3 + 7k^2 a^2}{ka(1 + k^2 a^2)^3} + \frac{9}{2k^4 a^4} H(ka)$	$\frac{15}{2} \frac{1}{ka(1 + k^2 a^2)^2} + \frac{3}{2k^4 a^4} H(ka)$
$[70; 2^-]_{(0;0);1}$	0	0	0
$[70; 2^+]_{(0;0);2}$	$\frac{1}{2} \frac{1}{(1 + k^2 a^2)^2} + \frac{3}{2k^3 a^3} H(ka)$	$\frac{1}{2} \frac{3 + 7k^2 a^2}{ka(1 + k^2 a^2)^3} + \frac{9}{2k^4 a^4} H(ka)$	$\frac{3}{2} \frac{1}{ka(1 + k^2 a^2)^2} + \frac{3}{2k^4 a^4} H(ka)$
$[56; 0^+]_{(1;0);0}$	$\frac{2k^2 a^2}{(1 + k^2 a^2)^3}$	$\frac{4ka(1 - 2k^2 a^2)}{(1 + k^2 a^2)^4}$	0
$[70; 1^-]_{(1;0);1}$	$i \frac{1}{2} \frac{1}{3} \frac{ka(1 - 3k^2 a^2)}{(1 + k^2 a^2)^3}$	$i \frac{1}{2} \frac{1}{3} \frac{1 - 14k^2 a^2 + 9k^4 a^4}{(1 + k^2 a^2)^4}$	$i \frac{1}{2} \frac{1}{3} \frac{1 - 3k^2 a^2}{(1 + k^2 a^2)^3}$
$[70; 0^+]_{(0;1);0}$	$\frac{2k^2 a^2}{(1 + k^2 a^2)^3}$	$\frac{4ka(1 - 2k^2 a^2)}{(1 + k^2 a^2)^4}$	0
$[70; 1^-]_{(0;1);1}$	$i \frac{1}{2} \frac{1}{3} \frac{ka(1 - k^2 a^2)}{(1 + k^2 a^2)^3}$	$i \frac{1}{2} \frac{1}{3} \frac{1 - 8k^2 a^2 + 3k^4 a^4}{(1 + k^2 a^2)^4}$	$i \frac{1}{2} \frac{1}{3} \frac{k^2 a^2}{(1 + k^2 a^2)^3}$
$[56; 1^-]_{(0;1);1}$	$i \frac{1}{2} \frac{1}{6} \frac{k^3 a^3}{(1 + k^2 a^2)^3}$	$i \frac{1}{2} \frac{1}{6} \frac{3k^2 a^2(1 - k^2 a^2)}{(1 + k^2 a^2)^4}$	$i \frac{1}{2} \frac{1}{3} \frac{k^2 a^2}{(1 + k^2 a^2)^3}$

TABLE II. $SU_f(3)$ reduced matrix elements.

$(p_2; q_2)$	$\langle (p_2; q_2) T^{(1,1)} (p_1; q_1) \rangle$		
	$(p_1; q_1)$	$= 1$	$= 2$
$(1; 1)$	$(1; 1)$	$\frac{p-5}{p-3}$	$\frac{1}{p-3}$
$(1; 1)$	$(1; 1)$	$\frac{p-5}{p-3}$	$p-3$
$(3; 0)$	$(3; 0)$	$\frac{2}{p-3} \frac{p-2}{3}$	
$(3; 0)$	$(1; 1)$	$\frac{2}{p-3} \frac{p-2}{3}$	
$(3; 0)$	$(1; 1)$	0	
$(1; 1)$	$(3; 0)$	$\frac{p-10}{p-3}$	
$(1; 1)$	$(3; 0)$	0	

TABLE III. Spin-avormatrix elements m ($m = 0; \pm 1$) of Eq. (12) for strong decay of baryons $B \rightarrow B^0 + M$ with $M = \pi$. The final state is $^2 8_{1=2} [56; 0^+]$ for the nucleon ($B^0 = N$) and $^4 10_{3=2} [56; 0^+]$ for the delta ($B^0 = \Delta$).

Initial State	$A_{1=2}(N)$			$A_{1=2}(\Delta)$			$A_{3=2}(\Delta)$		
	0	± 1	± 2	0	± 1	± 2	0	± 1	± 2
$^2 8_J [56; L^P]$	$\frac{5}{6} \frac{p}{3}$	$\frac{5}{3} \frac{p}{3}$	0	$\frac{2}{3} \frac{p}{3}$	$\frac{2}{3} \frac{p}{3}$	0	0	$\frac{2}{3} \frac{p}{3}$	0
$^2 8_J [70; L^P]$	$\frac{p}{3} \frac{2}{3}$	$\frac{2}{3} \frac{p}{3}$	0	$\frac{2}{3} \frac{p}{3}$	$\frac{2}{3} \frac{p}{3}$	0	0	$\frac{2}{3}$	0
$^4 8_J [70; L^P]$	$\frac{1}{3} \frac{p}{6}$	$\frac{1}{3} \frac{p}{6}$	$\frac{1}{3} \frac{p}{2}$	$\frac{1}{3} \frac{p}{3}$	$\frac{4}{3} \frac{p}{3}$	$\frac{2}{3}$	$\frac{1}{3} \frac{p}{3}$	$\frac{2}{3}$	0
$^2 8_J [20; L^P]$	0	0	0	0	0	0	0	0	0
$^4 10_J [56; L^P]$	$\frac{2}{3} \frac{p}{3}$	$\frac{2}{3} \frac{p}{3}$	$\frac{2}{3}$	$\frac{p}{6} \frac{5}{3}$	$\frac{2}{3} \frac{p}{3} \frac{5}{3}$	$\frac{p}{3} \frac{5}{3}$	$\frac{p}{2} \frac{5}{3}$	$\frac{p}{3} \frac{5}{3}$	0
$^2 10_J [70; L^P]$	$\frac{1}{6} \frac{p}{3}$	$\frac{1}{3} \frac{p}{3}$	0	$\frac{p}{3} \frac{5}{3}$	$\frac{p}{3} \frac{5}{3}$	0	0	$\frac{p}{3} \frac{5}{3}$	0

TABLE IV. Same as Table III, but with $M = \frac{1}{2}$ and $M = 0$ corresponding to $\chi = (\cos \varphi - \frac{1}{2} \sin \varphi) = \frac{1}{3}$ and $\chi = (\sin \varphi + \frac{1}{2} \cos \varphi) = \frac{1}{3}$, respectively.

Initial State	$A_{1=2} (N) =$			$A_{1=2} (\chi) =$			$A_{3=2} (\chi) =$		
	0	+		0	+		0	+	
$^2 8_J [56; L^P]$	$\frac{1}{6}$	$\frac{1}{3}$	0	0	0	0	0	0	0
$^2 8_J [70; L^P]$	$\frac{1}{3} \frac{1}{2}$	$\frac{1}{3} \frac{1}{2}$	0	0	0	0	0	0	0
$^4 8_J [70; L^P]$	$\frac{1}{3} \frac{1}{2}$	$\frac{1}{3} \frac{1}{2}$	$\frac{1}{6}$	0	0	0	0	0	0
$^2 8_J [20; L^P]$	0	0	0	0	0	0	0	0	0
$^4 10_J [56; L^P]$	0	0	0	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0
$^2 10_J [70; L^P]$	0	0	0	$\frac{1}{3}$	$\frac{1}{3}$	0	0	$\frac{1}{3}$	0

TABLE V. Coefficients c_i of Eq. (20) for the strong decay widths $B \rightarrow B^0 + M$ of the negative parity resonances with $(v_1; v_2); L^P = (0; 0); 1^-$. The final state is $^2 8_{1=2} [56; 0^+]_{(0;0);0}$ for the nucleon ($B^0 = N$) and $^4 10_{3=2} [56; 0^+]_{(0;0);0}$ for the delta ($B^0 = \Delta$). α is defined in the caption of Table IV.

State	N		N		Δ		Δ	
	c_0	c_2	c_0	c_2	c_0	c_2	c_0	c_2
$S_{11} \quad ^2 8_{1=2} [70; 1^-]_{(0;0);1}$	$\frac{8}{3}$		2^{-2}			$\frac{16}{3}$		
$D_{13} \quad ^2 8_{3=2} [70; 1^-]_{(0;0);1}$		$\frac{8}{3}$		2^{-2}	$\frac{8}{3}$	$\frac{8}{3}$		
$S_{11} \quad ^4 8_{1=2} [70; 1^-]_{(0;0);1}$	$\frac{2}{3}$		2^{-2}			$\frac{4}{3}$		
$D_{13} \quad ^4 8_{3=2} [70; 1^-]_{(0;0);1}$		$\frac{1}{15}$		$\frac{1}{5} 2^{-2}$	$\frac{20}{3}$	$\frac{64}{15}$		
$D_{15} \quad ^4 8_{5=2} [70; 1^-]_{(0;0);1}$		$\frac{2}{5}$		$\frac{6}{5} 2^{-2}$		$\frac{28}{5}$		
$S_{31} \quad ^2 10_{1=2} [70; 1^-]_{(0;0);1}$	$\frac{1}{3}$					$\frac{20}{3}$		4^{-2}
$D_{33} \quad ^2 10_{3=2} [70; 1^-]_{(0;0);1}$		$\frac{1}{3}$			$\frac{10}{3}$	$\frac{10}{3}$	2^{-2}	2^{-2}

TABLE VI. Coefficients c_1 of Eq. (20) for the strong decay widths of the positive parity resonances with $(v_1; v_2); L^P = (0; 0); 2^+$. For notation see Table V.

State	N		N		N		N	
	c_1	c_3	c_1	c_3	c_1	c_3	c_1	c_3
$P_{13} \quad {}^2 8_{3=2} [56; 2^+]_{(0;0);0}$	$\frac{25}{12}$		$\frac{1}{4} \quad {}^2$		$\frac{4}{15}$	$\frac{12}{5}$		
$F_{15} \quad {}^2 8_{5=2} [56; 2^+]_{(0;0);0}$		$\frac{25}{12}$		$\frac{1}{4} \quad {}^2$	$\frac{8}{5}$	$\frac{16}{15}$		
$P_{13} \quad {}^2 8_{3=2} [70; 2^+]_{(0;0);2}$	2		$\frac{3}{2} \quad {}^2$		$\frac{2}{5}$	$\frac{18}{5}$		
$F_{15} \quad {}^2 8_{5=2} [70; 2^+]_{(0;0);2}$		2		$\frac{3}{2} \quad {}^2$	$\frac{12}{5}$	$\frac{8}{5}$		
$P_{11} \quad {}^4 8_{1=2} [70; 2^+]_{(0;0);2}$	$\frac{1}{2}$		$\frac{3}{2} \quad {}^2$		1			
$P_{13} \quad {}^4 8_{3=2} [70; 2^+]_{(0;0);2}$	$\frac{1}{4}$		$\frac{3}{4} \quad {}^2$		$\frac{16}{5}$	$\frac{9}{5}$		
$F_{15} \quad {}^4 8_{5=2} [70; 2^+]_{(0;0);2}$		$\frac{1}{14}$		$\frac{3}{14} \quad {}^2$	$\frac{21}{5}$	$\frac{128}{35}$		
$F_{17} \quad {}^4 8_{7=2} [70; 2^+]_{(0;0);2}$		$\frac{9}{28}$		$\frac{27}{28} \quad {}^2$		$\frac{27}{7}$		
$P_{31} \quad {}^4 10_{1=2} [56; 2^+]_{(0;0);0}$	$\frac{4}{3}$				$\frac{5}{12}$		$\frac{1}{4} \quad {}^2$	
$P_{33} \quad {}^4 10_{3=2} [56; 2^+]_{(0;0);0}$	$\frac{2}{3}$				$\frac{4}{3}$	$\frac{3}{4}$	$\frac{4}{5} \quad {}^2$	$\frac{9}{20} \quad {}^2$
$F_{35} \quad {}^4 10_{5=2} [56; 2^+]_{(0;0);0}$		$\frac{4}{21}$			$\frac{7}{4}$	$\frac{32}{21}$	$\frac{21}{20} \quad {}^2$	$\frac{32}{35} \quad {}^2$
$F_{37} \quad {}^4 10_{7=2} [56; 2^+]_{(0;0);0}$		$\frac{6}{7}$				$\frac{45}{28}$		$\frac{27}{28} \quad {}^2$
$P_{33} \quad {}^2 10_{3=2} [70; 2^+]_{(0;0);2}$	$\frac{1}{4}$				$\frac{1}{2}$	$\frac{9}{2}$	$\frac{3}{10} \quad {}^2$	$\frac{27}{10} \quad {}^2$
$F_{35} \quad {}^2 10_{5=2} [70; 2^+]_{(0;0);2}$		$\frac{1}{4}$			3	2	$\frac{9}{5} \quad {}^2$	$\frac{6}{5} \quad {}^2$

TABLE VII. N decay widths of (3 and 4 star) nucleon and delta resonances in MeV. The experimental values are taken from [17]. The mixing angle for the mesons is $\theta_p = 23^\circ$ [20]. θ_1 and θ_2 are defined in the caption of Table I.

State	Mass	Resonance	k (MeV)	(th)	(exp)
S_{11}	N (1535)	$^2 8_{1=2} [70; 1]_{(0;0);1}$	467	85	79 38
S_{11}	N (1650)	$^4 8_{1=2} [70; 1]_{(0;0);1}$	547	35	130 27
P_{13}	N (1720)	$^2 8_{3=2} [56; 2^+]_{(0;0);0}$	594	31	22 11
D_{13}	N (1520)	$^2 8_{3=2} [70; 1]_{(0;0);1}$	456	115	67 9
D_{13}	N (1700)	$^4 8_{3=2} [70; 1]_{(0;0);1}$	580	5	10 7
D_{15}	N (1675)	$^4 8_{5=2} [70; 1]_{(0;0);1}$	564	31	72 12
F_{15}	N (1680)	$^2 8_{5=2} [56; 2^+]_{(0;0);0}$	567	41	84 9
G_{17}	N (2190)	$^2 8_{7=2} [70; 3]_{(0;0);1}$	888	34	67 27
G_{19}	N (2250)	$^4 8_{9=2} [70; 3]_{(0;0);1}$	923	7	38 21
H_{19}	N (2220)	$^2 8_{9=2} [56; 4^+]_{(0;0);0}$	905	15	65 28
$I_{1;11}$	N (2600)	$^2 8_{11=2} [70; 5]_{(0;0);1}$	1126	9	49 20
P_{11}	N (1440)	$^2 8_{1=2} [56; 0^+]_{(1;0);0}$	398	$108 \frac{2}{1}$	227 67
P_{11}	N (1710)	$^2 8_{1=2} [70; 0^+]_{(0;1);0}$	587	$173 \frac{2}{2}$	22 17
S_{31}	(1620)	$^2 10_{1=2} [70; 1]_{(0;0);1}$	526	16	37 11
P_{31}	(1910)	$^4 10_{1=2} [56; 2^+]_{(0;0);0}$	716	42	52 19
P_{33}	(1232)	$^4 10_{3=2} [56; 0^+]_{(0;0);0}$	229	116	119 5
P_{33}	(1920)	$^4 10_{3=2} [56; 2^+]_{(0;0);0}$	723	22	28 19
D_{33}	(1700)	$^2 10_{3=2} [70; 1]_{(0;0);1}$	580	27	45 21
D_{35}	(1930)	$^2 10_{5=2} [70; 2]_{(0;0);1}$	729	0	52 23
F_{35}	(1905)	$^4 10_{5=2} [56; 2^+]_{(0;0);0}$	713	9	36 20
F_{37}	(1950)	$^4 10_{7=2} [56; 2^+]_{(0;0);2}$	741	45	120 14
$H_{3;11}$	(2420)	$^4 10_{11=2} [56; 4^+]_{(0;0);0}$	1023	12	40 22
S_{31}	(1900)	$^2 10_{1=2} [70; 1]_{(1;0);1}$	710	$2 \frac{2}{1}$	38 21
P_{33}	(1600)	$^4 10_{3=2} [56; 0^+]_{(1;0);0}$	513	$108 \frac{2}{1}$	61 32

TABLE V III. decay widths of (3 and 4 star) nucleon and delta resonances in MeV . For classification of the resonances see Table V II. The experimental values are taken from [17].

State	Mass	k (MeV)	l	(th)	(exp)	l	(th)	(exp)
S ₁₁	N (1535)	244				D	23	1 1
S ₁₁	N (1650)	345				D	24	7 5
P ₁₃	N (1720)	402	P	1	102 89	F	10	
D ₁₃	N (1520)	230	S	3	10 4	D	9	15 3
D ₁₃	N (1700)	386	S	111	2 4	D	114	14 26
D ₁₅	N (1675)	366				D	123	88 14
F ₁₅	N (1680)	370	P	2	13 5	F	3	1 1
G ₁₇	N (2190)	740	D	13		G	12	
G ₁₉	N (2250)	780				G	40	
H ₁₉	N (2220)	760	F	3		H	3	
I _{1,11}	N (2600)	1003	G	4		I	3	
P ₁₁	N (1440)	147				P	0:1 $\frac{2}{1}$	87 30
P ₁₁	N (1710)	394				P	70 $\frac{2}{2}$	41 33
S ₃₁	(1620)	320				D	89	67 26
P ₃₁	(1910)	546	P	4				
P ₃₃	(1920)	553	P	15		F	14	
D ₃₃	(1700)	386	S	55	112 53	D	89	12 10
D ₃₅	(1930)	560	P	0		F	0	
F ₃₅	(1905)	542	P	18		F	27	
F ₃₇	(1950)	575				F	36	77 20
H _{3,11}	(2420)	890				H	11	
S ₃₁	(1900)	539				D	$\frac{2}{1}$	
P ₃₃	(1600)	303	P	25 $\frac{2}{1}$	180 143	F	0	16 25

TABLE IX. $N \rightarrow N$ and $\Delta \rightarrow N$ decay widths of (3 and 4 star) nucleon and delta resonances in MeV. For classification of the resonances see Table V II. The experimental values are taken from [17]. The mixing angle for the mesons is $\theta_P = 23^\circ$ [20]. α_1 and α_2 are defined in the caption of Table I.

State	Mass	Γ (MeV)	(th)	(exp)
S_{11}	N (1535)	182	0.1	74-39
S_{11}	N (1650)	346	8	11-6
P_{13}	N (1720)	420	0.2	
D_{13}	N (1520)	150	0.6	
D_{13}	N (1700)	400	4	
D_{15}	N (1675)	374	17	
F_{15}	N (1680)	379	0.5	
G_{17}	N (2190)	791	11	
G_{19}	N (2250)	831	9	
H_{19}	N (2220)	811	0.7	
$I_{1,11}$	N (2600)	1054	3	
P_{11}	N (1710)	410	$17 \frac{2}{2}$	
P_{31}	(1910)	322	0.0	
P_{33}	(1920)	335	0.5	
D_{35}	(1930)	348	0	
F_{35}	(1905)	316	1	
F_{37}	(1950)	372	2	
$H_{3,11}$	(2420)	786	2	
S_{31}	(1900)	309	$3 \frac{2}{1}$	

TABLE X . Strong decay widths for $\Lambda \rightarrow N + \pi$ and $N \rightarrow N + \pi$ in MeV . Experimental values are from [17].

Resonance	L	(th)				(exp)	
		Ref. [15]	Ref. [11]	Ref. [12]	Present		
(1232) P_{33}	0	70	121	108	116	119	5
(1950) F_{37}	2	27	56	50	45	120	14
(2420) $H_{3,11}$	4	4		8	12	40	22
(2950) $K_{3,15}$	6	1		3	5	13	8
$N(1520)D_{13}$	1		85	74	115	67	9
$N(2190)G_{17}$	3			48	34	67	27
$N(2600)I_{1,11}$	5			11	9	49	20

TABLE XI. Strong decay widths for the missing nucleon and delta resonances below 2 GeV. The mixing angle for the mesons is $\theta_P = 23^\circ$ [20].

State	Mass	Resonance	(MeV)		
			N	Δ	N
$P_{1;2J}$	N (1720)	$^2 8_J [20; 1^+]_{(0;0);0}$	0	0	0
$D_{1;2J}$	N (1875)	$^2 8_J [70; 2^-]_{(0;0);1}$	0	0	0
P_{13}	N (1875)	$^2 8_{3=2} [70; 2^+]_{(0;0);2}$	56	56	9
F_{15}	N (1875)	$^2 8_{5=2} [70; 2^+]_{(0;0);2}$	85	43	19
S_{11}	N (1972)	$^4 8_{1=2} [70; 2^-]_{(0;0);1}$	0	0	0
D_{13}	N (1972)	$^4 8_{3=2} [70; 2^-]_{(0;0);1}$	0	0	0
D_{15}	N (1972)	$^4 8_{5=2} [70; 2^-]_{(0;0);1}$	0	0	0
G_{17}	N (1972)	$^4 8_{7=2} [70; 2^-]_{(0;0);1}$	0	0	0
P_{11}	N (1972)	$^4 8_{1=2} [70; 2^+]_{(0;0);2}$	19	15	17
P_{13}	N (1972)	$^4 8_{3=2} [70; 2^+]_{(0;0);2}$	9	94	9
F_{15}	N (1972)	$^4 8_{5=2} [70; 2^+]_{(0;0);2}$	4	156	4
F_{17}	N (1972)	$^4 8_{7=2} [70; 2^+]_{(0;0);2}$	18	96	20
S_{11}	N (1909)	$^2 8_{1=2} [70; 1^-]_{(1;0);1}$	$22 \frac{2}{1}$	$0.7 \frac{2}{1}$	$0.5 \frac{2}{1}$
D_{13}	N (1909)	$^2 8_{3=2} [70; 1^-]_{(1;0);1}$	$28 \frac{2}{1}$	$0.6 \frac{2}{1}$	$\frac{2}{1}$
P_{13}	N (1815)	$^4 8_{3=2} [70; 0^+]_{(0;1);0}$	$30 \frac{2}{2}$	$211 \frac{2}{2}$	$24 \frac{2}{2}$
S_{11}	N (1866)	$^2 8_{1=2} [56; 1^-]_{(0;1);1}$	$211 \frac{2}{2}$	$84 \frac{2}{2}$	$4 \frac{2}{2}$
D_{13}	N (1866)	$^2 8_{3=2} [56; 1^-]_{(0;1);1}$	$271 \frac{2}{2}$	$71 \frac{2}{2}$	$7 \frac{2}{2}$
$P_{1;2J}$	N (1997)	$^2 8_J [70; 1^+]_{(0;1);0}$	0	0	0
S_{11}	N (1997)	$^2 8_{1=2} [70; 1^-]_{(0;1);1}$	$3 \frac{2}{2}$	$34 \frac{2}{2}$	$5 \frac{2}{2}$
D_{13}	N (1997)	$^2 8_{3=2} [70; 1^-]_{(0;1);1}$	$3 \frac{2}{2}$	$30 \frac{2}{2}$	$7 \frac{2}{2}$
D_{33}	(1945)	$^2 10_{3=2} [70; 2^-]_{(0;0);1}$	0	0	0
P_{33}	(1945)	$^2 10_{3=2} [70; 2^+]_{(0;0);2}$	9	105	4
F_{35}	(1945)	$^2 10_{5=2} [70; 2^+]_{(0;0);2}$	13	83	2
D_{33}	(1977)	$^2 10_{3=2} [70; 1^-]_{(1;0);1}$	$5 \frac{2}{1}$	$7 \frac{2}{1}$	$3 \frac{2}{1}$
P_{31}	(1786)	$^2 10_{1=2} [70; 0^+]_{(0;1);0}$	$27 \frac{2}{2}$	$172 \frac{2}{2}$	0.0

FIGURES

FIG .1. Collective model of baryons.

FIG .2. Elementary meson emission.

FIG .3. Strong decay widths for $N \rightarrow N + \pi$ decays of negative parity resonances with $L^P = 1^-$. The theoretical values are in parenthesis. All values in MeV.

FIG .4. Strong decay widths for $\Delta \rightarrow N + \pi$ and $\Delta \rightarrow \pi + \pi$ decays of positive parity resonances with $L^P = 2^+$ and negative parity resonances with $L^P = 1^-$. Notation as in Fig. 3.







