System atic analysis of group identi cation in stock m arkets

Dong-Hee Kim and Hawoong Jeong^y

Department of Physics, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Korea

(D ated: January 24, 2022)

W e propose in proved m ethods to identify stock groups using the correlation m atrix of stock price changes. By ltering out the marketwide e ect and the random noise, we construct the correlation m atrix of stock groups in which nontrivial high correlations between stocks are found. Using the

Itered correlation matrix, we successfully identify the multiple stock groups without any extra know ledge of the stocks by the optim ization of the matrix representation and the percolation approach to the correlation-based network of stocks. These m ethods drastically reduce the am biquities while nding stock groups using the eigenvectors of the correlation matrix.

PACS num bers: 89.65.G h, 05.40.C a, 89.75 Fb, 89.75.-k

I. IN TRODUCTION

The study of correlations in stock markets has attracted much interest of physicists because of its challenging complexity as a complex system and its possible future applications to the realmarkets [1]. In the early years, a correlation-based taxonom y of stocks and stock market indices was studied by the method of the hierarchical tree [2, 3]. Recently, the minimum spanning tree technique was introduced to study the structure and dynam ics of the stock network [4, 5, 6], the random matrix theory was applied to nd out the di erence between the random and nonrandom property of the correlations [7, 8, 9, 10, 11], and the maximum likelihood clustering method was developed and applied to identify cluster structures in stock markets [12]. A lso, these studies have been extended to the applications to the portfolio optim ization in realm arket [5, 9].

C om m only, the correlation between stocks is expressed by the Pearson correlation coe cient of log-returns,

$$G_{i}(t) \ln S_{i}(t+t) \ln S_{i}(t);$$
 (1)

where S_i (t) is the price of stock i at time t. From real time series data of N stock prices, we can calculate the elem ent of N N correlation matrix C as following

$$C_{ij} = \frac{h(G_{i}(t) \quad hG_{i}i) (G_{j}(t) \quad hG_{j}i)i}{(hG_{i}^{2}i \quad hG_{i}i^{2}) (hG_{j}^{2}i \quad hG_{j}i^{2})}; \qquad (2)$$

where h time series. By de nition, $C_{ii} = 1$ and C_{ij} has a value in [1;1].

Labux et al. [7] and Plerou et al. [8, 9] studied the statistical properties of an empirical correlation matrix between stock price changes de ned in Eq. 2) for real markets. In comparison with the prediction of the random matrix theory, they found that the statistics of the

bulk eigenvalues are in remarkable agreem ents with the universal properties of the random correlation matrix. For example, the bulk part of the eigenvalue spectrum of the empirical correlation matrix for N stocks over L price data has the form of the spectrum of the random correlation m atrix [13] which is given by

$$() = \frac{Q}{2} \frac{p_{(max)}(min)}{(min)};$$
 (3)

2 [m in ; m ax] in the lim it of N;L $[p \ 1 \ w$ ith Q L = N, where $m \text{ ax} = (1 + 1 = p \ 2)^2$ and = $(1 \ 1 = p \ 2)^2$. M oreover, the level spacing statisfor xed Q m in = (1)tics of eigenvalues exhibits good agreem ent with the results from the Gaussian orthogonal ensemble of random m atrices [8, 9].

On the other hand, the nonrandom properties of the correlation matrix have also been studied with the empirical correlation matrix [8, 9, 10]. From the empirical data for the New York Stock Exchange, it was found that each eigenvector corresponding to the few largest eigenvalues larger than the upper bound of the bulk eigenvalue spectrum, is localized, in a sense that only a few com ponents contribute to the eigenvector mostly, and the stocks corresponding to those dom inant com ponents of the eigenvector are found to belong to a comm on industry sector. Very recently, Utsugiet al. con med and improved those results through the similar analysis for the Tokyo Stock Exchange [11].

In order to con m the localization property of eigenvectors, we perform the sim ilar analysis to the previous i indicates tim e averages over the period of the studies [8, 9, 10] on eigenvectors of the correlation m atrix using our own dataset of stock prices. We analyze the daily prices of N = 135 stocks belonging to the New York Stock Exchange (NYSE) for the 20-year period 1983 2003 (L ' 5000 trading days) which is publicly available from the web-site (http:// nance.yahoo.com) [14]. Indeed, if we put stocks in the order of their industrial sectors, we observe that the eigenvector com ponents corresponding to stocks which belong to speci c industrial sectors give high contributions to each of the eigenvectors for the few largest eigenvalues (see Fig. 1). For instance, the stocks belonging to the energy, technology, transportation, and utilities sectors highly contribute to

Em ailaddress: dhkim @ stat.kaist.ac.kr

^yEm ailaddress: hjeong@ kaist.ac.kr




FIG. 1: The normalized eigenvector components $u_i()$ of stock i corresponding to the second to fith largest eigenvalues $_1 - _4$ of the correlation matrix. The stocks are sorted by industrial sectors, A : basic materials, B : capital goods, C : conglom erates, D : consum er (cyclical), E : consum er (noncyclical), F : energy, G : nancial, H : healthcare, I : services, J : technology, K : transportation, and L : utilities, which are separated by dashed lines.

the eigenvector for the second largest eigenvalue; the energy sector constitutes the big part of the eigenvector for the third largest eigenvalue; the fourth largest eigenvalue gives the eigenvector localized on the basic m aterials, consum er (noncyclical), healthcare, and utilities sectors; the eigenvector for the fth largest eigenvalue is also localized on several speci c industrial sectors.

However, it is not straightforward to nd out speci c stock groups, such as the industrial sectors, inversely. If each of the eigenvectors had well-de ned dom inant com ponents and the corresponding set of stocks were independent of the sets from other eigenvectors, it would becom e easy to identify the stock groups. Unfortunately, in our study, it turns out that not only the set of eigenvector components with dominant contribution can be hardly de ned in the eigenvector but also such a set is likely to overlap with the sets from other eigenvectors unless we pick a very small number of stocks with few highest ranks of their contributions to the eigenvectors; Figure 1 indicates that each of the eigenvectors is localized on a multiple number of industrial sectors and the corresponding stocks severely overlap with those from the other eigenvectors. Therefore it is very am biguous to identify the stock groups for practical purposes. The aim of this study is to get rid of these am biguities and

nally nd out relevant stock groups without any aid of the table of industrial sectors.

In this paper, we introduce the improved method to identify stock groups which drastically reduce the ambiquities in nding multiple groups using eigenvectors of the correlation matrix. We rst lter out the random noise and the marketwide e ect from the correlation matrix. With the ltered correlation matrix, we apply optim ization and percolation approaches to nd the stock groups. Through the optim ization of the stock sequences representing the matrix indices, the litered correlation matrix is transformed into the block diagonal matrix in which all stocks in a block are found to belong to the sam e group. By constructing a network of stocks using the percolation approach on the ltered correlation matrix, we also successfully identify the stock groups which appear in the form of isolated clusters in the resulting network.

This paper is organized as follows. In Sec. II, the detailed litering method to construct the group correlation matrix is given. For the litering, the largest eigenvalue and the corresponding eigenvector are required and they are calculated from the rst-order perturbation theory. In Sec. III, detailed stock group noting methods using the optimization and the percolation are given and the resulting stock groups are specified. In Sec.IV, a sum – mary and conclusions are presented.

II. GROUP CORRELATION MATRIX

A. Filtering

The group of stocks is de ned as a set of highly intercorrelated stocks in their price changes. In the empirical correlation m atrix, because several types of noises are expected to coexist with the intragroup correlations, it is essential to liter out such noises to isolate the intragroup correlations which we are interested in. With the com plete set of eigenvalues and eigenvectors, the correlation m atrix in Eq. (2) can be expanded as

$$C = \int_{0}^{1} j \, \mathrm{i}h \, j \qquad (4)$$

where is the eigenvalue sorted in descending order and j i is the corresponding eigenvector. Because only the eigenvectors corresponding to the few largest eigenvalues are believed to contain the inform ation on signi cant stock groups, we can identify a litered correlation m atrix for stock groups by choosing a partial sum of j ih j relevant to stock groups, which we will call the group correlation m atrix, C^g.

In order to extract C^g from the correlation matrix, taking the previous results of P lerou et al. [8, 9, 10]for granted, we posit that the eigenvalue spectrum of the correlation matrix is organized by the marketwide part




FIG.2: (a) The eigenvalues > 1.0 of the correlation matrix C and (b) the distribution of bulk eigenvalues P () (solid line). The dashed-dot line marks our boundary between the random noise part and the group correlation part. (c) The matrix element distribution for the group correlation matrix C^g and the residual parts corresponding to the bulk eigenvalues C^r and the largest eigenvalue C^m.

of the largest eigenvalue, the group part of interm ediate discrete eigenvalues, and the random part of sm all bulk eigenvalues. Then, we can separate the correlation m atrix into three parts as

$$C = C^{m} + C^{g} + C^{r}$$

$$= {}_{0}j0ih0j + {}_{j}ihj + {}_{N_{g}+1}$$

$$M_{X}^{1} jihj + {}_{N_{g}+1} jihj (5)$$

where C^{m} , C^{g} , and C^{r} indicate the marketwide effect, the group correlation matrix, and the random noise term s, respectively.

W hile the determ ination of C^m is straightforw ard, it is not so clear to determ ine N $_{\rm g}$ for separating C $^{\rm g}$ and C $^{\rm r}$. If there were no correlation between stock prices, the bulk eigenvalues have to follow Eq. (3), and thus the upper bound of the bulk eigenvalues can be clearly determ ined from Q. However, in empirical correlation matrix, the bulk eigenvalue spectrum deviates from Eq. (3) due to the coupling with underlying structured correlations, such as the group correlation embedded in C^g [15]. Therefore we use a graphical estim ation to determ ine N_g; in the eigenvalue spectrum as shown in Fig. 2 (b) we choose the cut $N_g = 9$ in the vicinity of the blurred tail of the bulk part of the spectrum . Nevertheless, in spite of the rough estimation of N $_{\rm q}$, we note that our results in this work do not alter from a small change of N $_{\rm q}$, 1. This can be justi ed by the following arguments. In the group correlation m atrix, the corresponding component of the eigenvalues close to the bulk part of the spectrum is con-

ned to only a very small portion of the whole matrix; because the elements of the correlation matrix com ponent j ih jmust be smaller than the eigenvalue ,




FIG.3: The Q L=N dependence of the matrix element distribution for C^g (thick solid line), C^r (solid line), and C^m (dashed line). W ith xed N = 135, various time periods are tested for (a) L ' 2600 (1993 2003), (b) L ' 950 (2000 2003), and (c) L ' 240 (2003).

large discrete eigenvalues dom inantly contribute to the group correlation m atrix. In addition, even if we count one less eigenvalue near the boundary of bulk part of the spectrum in constructing the group correlation m atrix, a possible inform ation loss of groups is not likely serious because the pure eigenvectors of the groups generally turn out to be m ixed all together in the eigenvectors of the correlation m atrix (see Fig. 1). Therefore the in uence from the error in the determ ination of N_g is insigni cant so that it does not change the clustering result.

This decomposition of the correlation matrix gives nontrivial characteristics to the distribution of the group correlation matrix elements C_{ij}^{g} . In Fig. 2(c), it turns out that the distribution of C_{ij}^{g} shows positive heavy tail. This indicates that C^g contains a non-negligible num ber of strongly correlated stock pairs, which is expected to come from the correlation between the stocks belonging to the same group. On the other hand, C^{r} shows the Gaussian distribution consistent with the prediction of the random matrix theory [9]. W hile this Gaussian-like distribution is also observed partially in the distribution of C^g_{ii} due to the coupling between group correlations and random noises, it turns out that this rem aining noise does not seriously a ect the identication of stock groups. The distribution of C_{ij}^m shows that C^m also contains highly correlated stock pairs, but we nd that C_{ij}^m is not relevant to the group correlation and thus have to be ltered out for the clear identic cation of the stock groups, which is discussed in Sec. IIB.

Since the quality of the correlation m atrix can depend on the period of empirical data or generally Q L=N, our decom position of the correlation m atrix can also depend on Q. Here we simply check how the determ ination of N $_{\rm q}$ and the resulting matrix element distribution of the decomposed matrices are changed depending on Q (see Fig. 3). For Q = 19:4 (1993 2003) and Q = 7:0(1999 2003), C^{g} and C^{r} are separated at $N_{g} = 10$, which is not very di erent from $N_{\alpha} = 9$ of the larger dataset we use throughout this paper, and in addition, the distribution of the matrix element shows the sim ilar degree of the heavy tail in C_{ij}^{g} . However, decreasing Q much smaller, the bulk eigenvalue spectrum becomes wider so that m ore eigenvalues relevant to the group correlation can be buried in the bulk spectrum, which leads to smaller N_g that turns to be 7 for Q = 1.7 (2003). Even in this case of Q = 1:7, the positive heavy tail is still found in C $_{ij}^{g}$ but very weaker than higher Q 's. T hese im ply that we need a large enough Q for the stock group identi cation.

B. Largest eigenvalue and corresponding eigenvector

Our Itering is based on the following interpretations of the previous studies: the bulk part of the eigenvalues and their eigenvectors are expected to show the universal properties of the random matrix theory and the largest eigenvalue and its eigenvector are considered as a collective response of the entire market [8, 9, 10]. While the random characteristics of the bulk eigenvalues have been studied intensively, only the empirical tests have been done for the largest eigenvalue and its eigenvector so far [9, 10]. Thus, to understand the more accurate meaning, we calculate the largest eigenvalue and its eigenvector of the correlation matrix by using perturbation theory.

In stock markets, it has been understood that there exist three kinds of uctuations in stock price changes: a marketwide uctuation, synchronized uctuations of stock groups, and a random uctuation §, 9, 10]. For simplicity, we consider a situation in which a system with only the marketwide uctuation is perturbed by other

uctuations. Let us assume that the price changes of all the stocks in the market nd a synchronized background uctuation with zero mean and variance g as a marketwide e ect. Then, we can write down the N N unperturbed correlation matrix as

$$C^{0} = B^{B}_{B} C_{0} 1 : C^{C}_{C} C^{0}_{C} C^{0}_$$

which has the largest eigenvalue $_{0}^{(0)} = c_{0} (N - 1) + 1$ and its eigenvector components $u_{i}^{(0)} = h(\text{stock})$ if $D^{(0)}i = 1 = \frac{N}{N}$.

W hen a sm all perturbation is turned on, the total cor-




FIG. 4: The comparison of the eigenvector of the largest eigenvalue obtained by the exact diagonalization u_i and the dom inant term ($w_i + c_0$)=(c_0 N³⁻²) in Eq. (8). The dashed line has the slope 1:0. (Inset: the values of corresponding eigenvector components.)

relation m atrix becom es

$$C = C^{0} + ;$$
 (7)

where $_{ii} = 0$ and $_{ij} = _{ji}$. Applying the perturbation theory up to the rst order, the largest eigenvalue and the corresponding eigenvector components are easily calculated as

$$u_{i} = c_{0} (N = 1) + 1 + \frac{1}{N} X_{i;j};$$

$$u_{i} = \frac{1}{c_{0}N^{3=2}} @w_{i} + c_{0} = \frac{1}{N} X_{j;k}^{X}; (8)$$

where $w_i = \frac{P}{j \in i} C_{ij}$.

We check the validity of Eqs. (8) by comparing with the largest eigenvector obtained from the num erical diagonalization of the empirical correlation matrix. For the comparison, we make the distribution of C_{ij} in Eq. (7) to be close to the em pirical C_{ij} distribution by assuming ij follows the bell-shaped distribution with zero that m ean and letting c_0 to the m ean value of the empirical C_{ij} . Because the assumption not only reproduces the distribution of pem pirical Cij, but also allows us to neij term in Eqs. (8), we can directly glect the 1=N com pare the perturbation theory with the num erical result. Figure 4 displays the eigenvector components of the largest eigenvalue obtained from the empirical correlation m atrix and the dom inant term s of Eqs. (8), which show remarkable agreem ent with each other.

Equation (8) indicates that the eigenvector of the largest eigenvalue is contributed by not only the global

uctuation but also the unknown perturbations from including random noises. Thus, by $\exists tering out the C^m$

term, we can decrease the e ect of unnecessary perturbations in constructing the group correlation m atrix. Indeed, as seen in Fig.2(c), because the heavy tailpart of C^g, the highly correlated elements, are buried in C^m, the clustering of stocks would be seriously disturbed unless C^m is litered out.

In addition, Eqs. (8) also enable us to interpret more detailed m eaning of the eigenvector than the m arketw ide e ect. Because the ith eigenvector component u is mostly determ ined by w_i , the sum of the correlation over all the other stocks, it can be regarded as the in uencing power of the com pany in the entire stock m arket. In real data, the top four stocks with highest w_i are found to be GeneralE lectric (GE), American Express (AXP), Merrill Lynch (MER), and Emerson Electric (EMR), mostly conglom erates or huge nancial com panies, which convinces us that ui is indeed representing the in uencing power of stock i. However, these high in uencing companies prevent clear clustering of stocks because of their nonnegligible correlations with entire stocks in the market. This is easily comprehensible by considering an analogous situation in a network where the big hub, a node with a large number of links, can make indispensable connections between groups of nodes to cause di culties in distinguishing the groups [16]. Therefore it is very in portant to lter out C^m in order to identify the groups of stocks e ciently.

III. IDENTIFICATION OF STOCK GROUPS

In the group m odel for stock price correlation proposed by N oh [17], the correlation matrix C takes the form of C = C^g + C^r, where C^g and C^r are the correlation matrix of stock groups and random correlation matrix, respectively. The model assumes the ideal situation with C^g_{ij} = ____i; , where __i indicates the group to which the stock i belongs. Thus C^g is the block diagonal matrix,

$$C^{g} = \begin{bmatrix} 0 & 1_{0} & 0 & 1_{0} \\ B & 0 & 1_{1} & \vdots & C \\ B & 0 & 1_{1} & \vdots & C \\ C & \vdots & \ddots & 0 & A \\ 0 & 0_{n} & 1 \end{bmatrix}$$
(9)

where 1_i is the N_i N_im atrix (N_i is the num ber of stocks in the ith group) of which all elements are 1.

Here we use this group model to nd the groups of stocks. If the correlation matrix in the real market can be represented by the block diagonal matrix as in the model, it would be very easy to identify the groups of stocks. However, there exist in nitely many possible representations of the matrix depending on indexing of rows and columns even if we have a matrix equivalent to the block diagonal matrix. For instance, if we exchange the indices of the matrix (e.g., fi; j;kg ! fk; i; jg) them atrix may not be block-diagonal anymore. Therefore the problem in identifying the groups in stock correlation matrix




FIG.5: (Color online) The visualization of the group correlation m atrix with the optimized stock sequence fl_ig .

requires one to nd out the optim ized sequence of stocks to transform the matrix into the well-organized block diagonalmatrix [18].

To optim ize the sequence of stocks for clear block diagonalization, we consider the correlation between two stocks as an attraction force between them. For the ideal group correlation matrix in the group model, the block diagonal form is evidently the most stable form if the attractive force between stocks is proportional to their correlation within the group. To deal with the real correlation matrix, we de ne the total energy for a stock sequence as

$$E_{tot} = \begin{array}{c} X \\ C_{ij}^{g} \mathbf{j}_{i} \quad \mathbf{l}_{j} \mathbf{j} \quad (C_{ij}^{g} \quad \mathbf{c}_{c}); \quad (10) \\ \\ \mathbf{i} < \mathbf{j} \end{array}$$

where l_i is the location of the stock i in the new index sequence and the cuto $c_c = 0:1$ is introduced to get rid of the random noise part which still remains in C^g in spite of the litering 20].

We obtain the optim ized sequence of stocks to minimize the total energy de ned in Eq. 10) by using the simulated annealing technique [21] in M onte C arlo simulation. The following description of our problem is very similar to the well-known traveling salesman problem,

nding an optim ized sequence of visiting cities which m inim izes total traveling distance [22]:

- 1. C on guration. The stocks are numbered i = 0;:::;N 1. A con guration, a sequence of stocks fl_ig, is a permutation of the numbers 0;:::;N 1.
- 2. Rearrangements. A random ly chosen stock in the sequence is removed and inserted at the random position of the sequence.
- 3.0 bjective function. W e use E_{tot} in Eq. (10) as an objective function to be m inim ized after rearrangements.

TABLE I: The full list of the optim ized sequence of stocks. The footnotes correspond to the identi ed stock groups represented by the same footnotes in Fig. 5.

pi	T icker	Sector	pi	T icker	Sector	pi	T icker	Sector
0	XNR	Services	45	G	Consumernoncyclical ⁵	90	AM R	Transportation ⁷
1	WМВ	Utilities	46	AVP	Consum er noncyclical⁵	91	F	Consum er cyclical ⁷
2	VLO	Energy ¹	47	МCD	Services	92	GΜ	Consum er cyclical ⁷
3	NBL	Energy ¹	48	ΙFF	Basic materials	93	ΗΡC	Basic materials ⁸
4	APA	Energy ¹	49	WМТ	Services	94	D D	Basic materials ⁸
5	КМG	Energy ¹	50	FNM	Financial	95	CAT	Capitalgoods ⁸
6	HAL	Energy ¹	51	ΕC	Consum er cyclical	96	DOW	Basic materials ⁸
7	SLB	Energy ¹	52	KR	Services	97	WΥ	Basic materials ⁸
8	ΒP	Energy ¹	53	ΗΕΤ	Services	98	IP	Basic materials ⁸
9	COP	Energy ¹	54	ΤΧΙ	Capitalgoods	99	GΡ	Basic materials ⁸
10	CVX	Energy ¹	55	FO	C onglom erates	100	BCC	Basic materials ⁸
11	ОХҮ	Energy ¹	56	SK Y	Capitalgoods	101	AA	Basic materials ⁸
12	RD	Energy ¹	57	FLE	C apital goods	102	ΡD	Basic materials ⁸
13	MRO	Energy ¹	58	R SH	Services	103	LPX	Basic materials ⁸
14	ХОМ	Energy ¹	59	ΕK	Consum er cyclical	104	Ν	Basic materials ⁸
15	ΡGL	U tilities ²	60	EM R	C onglom erates	105	DE	Capitalgoods
16	CNP	U tilities ²	61	ТОҮ	Services	106	PBI	Technology
17	ETR	U tilities ²	62	ΤΕΝ	Consum er cyclical	107	BDK	Consum er cyclical
18	DTE	U tilities ²	63	ROK	Technology	108	UNP	Transportation ⁹
19	ЕXС	U tilities ²	64	ΗΟΝ	Capital goods	109	N SC	T ransportation ⁹
20	AEP	U tilities ²	65	ΑXΡ	Financial	110	CSX	T ransportation ⁹
21	PEG	U tilities ²	66	GRA	Basic materials	111	BNI	T ransportation ⁹
22	SO	U tilities ²	67	VVI	Services	112	CNF	T ransportation ⁹
23	ΕD	U tilities ²	68	CSC	Technology ⁶	113	MAT	Consum er cyclical
24	PCG	U tilities ²	69	DBD	Technology ⁶	114	С	Financial
25	ΕIX	U tilities ²	70	HRS	Technology ⁶	115	VIA	Services
26	LM T	Capital goods ³	71	STK	Technology ⁶	116	МММ	C onglom erates
27	NOC	$Capital goods^3$	72	ZL	Technology ⁶	117	D IS	Services
28	RTN	Conglom erates ³	73	ΤΕΚ	Technology ⁶	118	BC	Consum er cyclical
29	GD	Capital goods ³	74	AVT	Technology ⁶	119	CBE	Technology
30	ВA	Capital goods ³	75	GLW	Technology ⁶	120	ТНС	H ealthcare ¹⁰
31	BOL	H ealthcare ⁴	76	N SM	$Technology^6$	121	HUM	Financial ¹⁰
32	МDТ	Healthcare ⁴	77	ΤΧΝ	Technology ⁶	122	AET	Financia 1^{10}
33	BAX	Healthcare ⁴	78	МОТ	Technology ⁶	123	CI	Financial ¹⁰
34	WYE	Healthcare ⁴	79	НРQ	Technology ⁶	124	JC P	Services
35	ВМ Ү	H ealthcare ⁴	80	ΝT	Technology ⁶	125	MEE	Energy
36	LLY	H ealthcare ⁴	81	ШM	Technology ⁶	126	GΕ	Conglom erates
37	MRK	Healthcare ⁴	82	U IS	Technology ⁶	127	UTX	C onglom erates
38	PFE	H ealthcare ⁴	83	XRX	Technology ⁶	128	R	Services
39	JN J	H ealthcare ⁴	84	Т	Services	129	NVO	H ealthcare
40	PEP	Consumernoncyclical ⁵	85	H IT	C apital goods	130	GΤ	Consum er cyclical
41	КO	Consumernoncyclical ⁵	86	MER	Financial	131	S	Services
42	ΡG	Consumernoncyclical ⁵	87	FDX	Transportation ⁷	132	NAV	Consum er cyclical
43	MO	Consum er noncyclical ⁵	88	LUV	Transportation ⁷	133	CEN	Technology
44	CL	Consumernoncyclical ⁵	89	DAL	Transportation ⁷	134	FL	Services

Figure 5 visualizes the correlation matrix elements $C_{l_i l_j}^{g}$ with the most optim ized sequence fl_ig and Table I lists the optim ized sequence of stocks. The multiple independent blocks of highly correlated correlations in the matrix are clearly visible without any a priori know ledge of stocks, i.e., the stocks in dimensional event blocks are believed to belong to dimensional event groups. We succeed to identify about 70% of the entire 135 stocks from the blocks, which are listed in Table I and it turns out that most of the stocks

in a block are represented by a single industry sector or a detailed industrial classi cation such as aerospace and defense, airline transport, railroad, and insurance (see Fig. 5). There still remain a small number of ungrouped stocks, which arises from the fact that the correlations between them are too weak to be distinguished from the random noise that still exists in the group correlation matrix.

As an alternative method, we also perform a network-




FIG.6: The dependence of the number of isolated clusters, m, in the stock network on the threshold p in constructing the network from (a) the group correlation matrix and (b) the full correlation matrix.

based approach to nd the groups of stocks. In principle, the correlation matrix can be treated as an adjacency matrix of the weighted network of stocks, in which the weights indicate how closely correlated the stocks are in their price changes [23]. How ever, for the sim plicity and the clear de nition of groups in the network, we consider the binary network of stocks which perm its only two possible states of a stock pair, connected or disconnected.

To construct the binary network of stocks, we use the percolation approach because of its usefulness of nding groups. The method is very simple: for each pair of stocks, we connect them if the group correlation coe cient C_{ij}^{g} is larger than a preassigned threshold value p. If the heavy tail in the distribution of C_{ij}^{g} in Fig. 2 mostly comes from the correlation between the stocks in the same group, an appropriate choice of $p = p_{c}$ will give severalm eaningful isolated clusters, m, in the network which are expected to be identieed as dimensional entry stocks groups.

We determ ine p_c by observing the change of the network structure as p decreases. Figure 6(a) displays the number of isolated clusters in the network as a function of threshold p. A swe decrease p, the number of isolated clusters in the network increases slow ly and stays near the maximum value up to p = 0.1, and then it abruptly decreases to 1, which indicates there exists on ly one isolated cluster. Therefore we choose $p_c = 0.1$ to construct the m ost clustered but stable stock network [24].

We nd that the constructed network consists of separable groups of stocks which correspond to the industrial sectors of stocks (see Fig. 7). At $p_c = 0.1$, the network has 92 nodes and 357 links. The identication of stock group is very clear because the clusters in the network, which we consider to be equivalent to stock groups, are fully connected networks or very dense networks in which most of the nodes in the cluster are directly connected.

However, although most of the stock groups are represented by a single industrial sector, it is found that the stocks which belong to two di erent industrial sectors coexist in a cluster. For instance, the stocks in the healthcare sector and the noncyclical consum er sector cannot be separable in this network. Indeed, in Fig. 5, one can observe non-negligible correlation between the healthcare and the noncyclical consum er, which indicates the presence of an intergroup correlation. In the real market, this presence of such an intersector correlation can be expected and our clustering results shown in Figs. 5 and 7 present both of intergroup and intragroup correlations that exist in the real stock market.

The group identication based on the eigenvector analysis of the stock price correlation matrix has been studied by several research groups [9, 10, 11]. In spite of their pioneering achievements to reveal the localization properties of eigenvectors, the classication of stocks into groups was not so clear, and it only covered about 10% of their stocks because they used only the few highest contributions of eigenvector components due to the am biguity explained in Sec. I. In this work, we not only introduce a more rened and systematicm ethod to identify the stock groups, but also successfully cluster about 70% of stocks into groups although direct comparison of the success ratio might be inappropriate because our data set is die erent from theirs.

On the other hand, Onnela et al. [6] introduced the percolation approach to construct the stock network in which the links are added between stocks one by one in descending order from the highest element of the full correlation m atrix. In their work, though highly correlated groups of stocks were found, the threshold value of the correlation to settle the network structure was hardly determ ined; the num ber of isolated clusters according to the threshold did not show the clear cut. W e believe that this is attributed to the fact that they used the full correlation m atrix carrying m arketwide and random uctuation.We would also fail to determ ine the critical threshold value of correlation if we use the full correlation m atrix instead of the ltered one [see Fig.6 (b)]. This indicates that the Itering is crucial for the stock group identi cation.

Finally, we note that M arsili et al. introduced a different m ethod to liter noises from the time series of stock price log-returns for stock group identic cation. In their work, it was assumed that the norm alized log-return could be expressed by the linear combination of the noise at individual stock level and the noise at the level of the groups, which tted to the real data to determ ine the weights of two noises and the constituents of the groups. However, we found that the elect of the inhom ogeneous marketwide uctuation is quite signic cant that the marketwide elect needs to be considered seriously to describe the correlation between stock correctly. Indeed, it is found that the litering out of the corresponding C^m im proves the clustering result.




FIG.7: The stock network with the threshold $p_c = 0.1$. The thickness of links indicates the strength of the correlation in the group correlation m atrix.

IV. CONCLUSION

In conclusion, we successfully identify the multiple group of stocks from the empirical correlation matrix of stock price changes in the New York Stock Exchange. We propose re ned methods to nd stock groups which dramatically reduce ambiguities as compared to identifying stock groups from the localization in a single eigenvector of the correlation matrix [9, 10, 11]. From the analysis of the characteristics of eigenvectors, we construct the group correlation matrix of the stock groups excluding the marketwide e ect and random noise. By optimizing the representation of the group correlation matrix, we nd that the group correlation matrix is represented by the block diagonal matrix where the stocks in each block belong to the same group. This coincides with the theoretical model of N oh [17]. Equally good stock group identication is also achieved by the percolation approach on the group correlation matrix to construct the network of stocks.

A cknow ledgm ents

W e thank J.D. Noh for helpfuldiscussions. This work was supported by grant No. R14-2002-059-01002-0 from KOSEF-ABRL program.

- [1] R. N. Mantegna and H. E. Stanley, An Introduction to Econophysics: Correlation and Complexity in Finance (Cambridge University Press, Cambridge, England, 2000); J. P. Bouchaud and M. Potters, Theory of FinancialRick (Cambridge University Press, Cambridge, England, 2000).
- [2] R.N.Mantegna, Eur.Phys.J.B 11, 193 (1999).
- [3] G.Bonanno, N.Vandewalle, and R.N.M antegna, Phys. Rev.E 62, R7615 (2000).
- [4] G.Bonanno, G.Caldarelli, F.Lilb, and R.N.M antegna, Phys. Rev. E 68, 046130 (2003).
- [5] J.-P.Onnela, A. Chakraborti, K. Kaski, J. Kertesz, and A. Kanto, Phys. Rev. E 68 056110 (2003).
- [6] J.-P. Onnela, K. Kaski, and J. Kertesz, Eur. Phys. J. B 38, 353 (2004).
- [7] L. Laloux, P. Cizeau, J.-P. Bouchaud, and M. Potters, Phys. Rev. Lett. 83, 1467 (1999)
- [8] V. Plerou, P. Gopikrishnan, B. Rosenow, L. A. Nunes Am aral, and H. E. Stanley, Phys. Rev. Lett. 83, 1471 (1999).
- [9] V. Plerou, P. Gopikrishnan, B. Rosenow, L. A. Nunes

Am aral, T.Guhr, and H.E.Stanley, Phys.Rev.E 65, 066126 (2002).

- [10] P.G opikrishnan, B.R osenow, V.P lerou, and H.E.Stanley, Phys. Rev. E 64, 035106 (R) (2001).
- [11] A. Utsugi, K. Ino, and M. Oshikawa, Phys. Rev. E 70, 026110 (2004).
- [12] L.G iada and M.M arsili, Phys.Rev.E 63,061101 (2001);
 M.M arsili, Quant.Finance 2,297 (2002); L.G iada and
 M.M arsili, Physica A 315,650 (2002).
- [13] A.M. Sengupta and P.P.M itra, Phys. Rev. E. 60, 3389 (1999).
- [14] Our selection of stocks covers most available stock price data for the period 1983 2003 in the database. We use the adjusted daily stock prices and skip 1 days of the data when splits and the dividends of any stock occur in Eqs. (1) and (2) to avoid the possible artifact caused by the abrupt stock price changes [see http://help.yahoo.com/help/us/ n/quote/index.htm 1].
- [15] Z.Burda and J.Jurkiewicz, Physica A 344, 67 (2004).
- [16] P.Holme, M.Huss, and H.Jeong, Bioinformatics 19, 532 (2003).

- [17] J.D.Noh, Phys.Rev.E 61, 5981 (2000).
- [18] The sim ilar matrix element rearranging technique was used in Ref. [19].
- [19] S.Bilke, e-print physics/0006050.
- [20] The visualization of C $^{\rm g}$ is not very sensitive to the cuto value $c_{\rm c}$.
- [21] S.Kirkpatrick, C.D.Gerlatt, and M.P.Vecchi, Science 220, 671 (1983).
- [22] W .H.Press, S.A.Teukolsky, W .T.Vetterling, and B.P. Flannery, Numerical Recipes in C (C am bridge University Press, C am bridge, England, 1992).
- [23] S.-W. Son, H. Jeong, and J. D. Noh, e-print cond-m at/0502672.
- [24] Sm all change of $p_{\rm c}\,,~$ 0.01, does not seriously change the clustering result.