M onte Carlo simulation of melting transition on DNA nanocompartment

Song Chang^{1;2}, Youdong M ao¹, Zhengwei X ie² Chunxiong Luo¹, Q iO uyang^{1;2;3}

¹Laboratory of B iotechnology, School of P hysics, P eking U niversity, Beijing 100871, C hina

²C enter for T heoretical B iology, P eking U niversity, B eijing 100871, C hina

³To whom correspondence should be addressed.

E-m ail: qi@ pku.edu.cn

A bstract

DNA nanocom partment is a typical DNA -based machine whose function is dependent of m olecular collective e ect. Fundam ental properties of the device have been addressed via electrochem ical analysis, uorescent m icroscopy, and atom ic force m icroscopy. Interesting and novelphenom ena em erged during the switching of the device. W e have found that DNAs in this system exhibit a much steep melting transition compared to ones in bulk solution or conventional DNA array. To achieve an understanding to this discrepancy, we introduced DNA-DNA interaction potential to the conventional Ising-like Z im m -B ragg theory and Peyrard-Bishop model of DNA melting. To avoid unrealistic num erical calculation caused by m odi cation of the Peyrard-B ishop nonlinear Ham iltonian with the DNA-DNA interaction, we established coarse-gained M onte C arlo recursion relations by elucidation of ve components of energy change during melting transition. The result suggests that DNA-DNA interaction potential accounts for the observed steep transition.

1.Introduction

Studies on the physical chemistry of DNA denaturation have been lasted for almost forty years [1-3]. In 1964, Lifson proposed that a phase transition exists in one-dimensional polymer structure. He introduced several pivotal concepts, like sequence partition function, sequence generating function, etc., and established a systematic method to calculate the partition function [1]. These allow us to derive important therm odynamic quantities of the system. In 1966, Poland and Scherage applied Lifson's method to conduct research on am ino acid and nucleic acid chains. They built Poland-Scherage (PS) model for calculating the sequence partition function and discussing the behavior of polymers in melting transitions.

A nother excellent progress would be the building of Peyrard-B ishop (PB) model [4,5] for DNA chains. In PB model, the H am iltonian of a single DNA chain, which is constructed by phonon calculations, is given so that we can obtain the system properties through statistical physics method. The PB model has introduced mathematical form ula of stacking energy, as well as the kinetic energy and potential energy of each base pair. By theoretical calculation, one can show the entropy-driven transition that leads DNA to shift from ordered state to disorder one [6,7].

However, all these works have not involved the DNA-DNA interactions because the subject investigated is DNAs in bulk solution, and the interaction between them has ever been neglected. The main idea of this paper is to inspect the in uence of collective e ect on the DNA melting process, primarily motivated by the experiment results of DNA nanocompartment [8,9]. Under the enlightenment of Poland-Scherage model and Z im m-B ragg model [10], we simplify Peyrard-B ishop model to meet a reasonable M onte C arlo simulation by the elucidation of ve components of energy changes during melting transition. The result shows that the melting temperature and transition duration depend on whether we take into account the DNA-DNA interactions among columnar assemblies of DNA.

2.Experim ent

Recently, we found that specially designed DNA array can form a molecular cage on surfaces [8,9]. This molecular cage is switchable due to allosteric transform ation driven by the collective hybridization of DNA. We named it "active DNA nanocom partment (ADNC)". TypicalDNA motif designed to fabricate ADNC comprises two contiguous elements (inset to gure 1a): a double-stranded DNA (dsDNA) whose array is responsible for a compact membrane (gure 1a, right), and a single-stranded DNA (ssDNA) serving as skeleton supporting the dsDNA membrane, which is term inated on its 5 end by a surface linker such as an alkanethiol group that can be tethered to gold surface with a sulphur-gold bond [9] or an am ino group that can be tethered to SiD₂ substrate with speci c surface attachment desDNA, a compartment with designable e ective height (he , 5 50nm, commensurate with the length of ssDNA skeleton) can form between the dsDNA membrane and substrate surface.

Since ADNC is reversibly switchable, it is able to encage molecules with suitable size. We name this phenomenon molecular encaging e ect. Both electrochem ical methods [12] and uorescent microscopy are used to substantiate the molecular encaging e ect and the reversibility of switching. O noe the closed ADNC entraps som e chem ical reporters, the surface concentration ($_{nc}$) of the encaged reporters can be determined by cyclic voltam metry or uorescent microscopy. Figure 1b shows the isotherms of the molecular encaging e ect for uorescein ($C_{20}H_{10}Na_2O_5$). Figure 1c presents the melting curves of ADNC. U sing the encaged molecules as indicator greatly sharpens the melting proles for the perfectly complementary targets, and

attens denaturation pro les for the strands with a wobble m ism atch. The observation shows that single-base m ism atched strands are incapable of closing ADNC on surfaces. The result is highly consistent to our observation by electrochem ical analysis [12]. These observations bring up an intriguing question: why the melting curves exhibit so steep transition compared to the case of DNA in bulk solutions or on a loosely packed m icroarray? W e try to address this question in this paper.

Worthy of mention is that the steepness of melting transition is useful when the ADNC is applied to DNA detection [8,9]. First, it greatly enhances the discrepancy of perfect targets and single m ism atches. This provides much enhanced speci city in DNA recognition, 100:1 105:1 of our system versus 2:7:1 of conventional system. Second, more sensitivity is obtained with optim ally decreased ambiguity. Therefore, the claric cation of the origin of the steep shape should help us to further extend the experience to related elds or generate new techniques.

3 M odeling

Taking into account the directional speci city of the hydrogen bonds, the Ham iltonian of a single DNA chain is obtained as following form according to PB model [4-6],

$$H_{y} = \sum_{n}^{X} \frac{1}{2} m y_{n}^{2} + w (y_{n}; y_{n-1}) + V (y_{n})$$
(1)

where the y_n is the component of the relative displacement of bases along the direction of hydrogen bond. The stacking energy w (y_n ; y_{n-1}) corresponds to the interaction between neighboring base pair in one DNA chain

$$w (y_n; y_{n-1}) = \frac{k}{2} 1 + e^{(y_n + y_{n-1})} (y_n - y_{n-1})^2$$
(2)

The M orse potential describes the potential for the hydrogen bonds

$$V = D (e^{-y} - 1)^2$$
 (3)

However, in this study, the Ham iltonian in equation (1) is not su cient; it neglects the structure of close-packing of DNA in ADNC. In our system, one should take into account the interactions between the nearest neighboring m olecules [13,14]. To m odel the interaction, one envisions the m olecules




Figure 1: (a) Schematic drawing of a dual-state ADNC.By adding or rem oving 'fuel' strands (shorter segment), the ADNC can be switched between on (right) and o (left) state. Inset, a typical sequence used to fabricate ADNC. The 'fuel' strands is a segment of hum an p53 gene containing one site of m ost frequent m utation. (b) Isotherm s of for uorescein encaged in a closed ADNC ($h_{eff} = 15bp$). The isotherm ts well to the Langm uir model: $x = nc = (1 = nc_{max})x + (1 = K nc_{max})$, where x is the concentration of the reporter and K is the association constant per site, nc the surface concentration of encaged molecules. (c) Melting curves using the encaged uorescein molecules as indicators. Filled circles or squares are corresponding to perfect com plem entary strands, and hollow circles or squares to single-base m ism atched strands. The unit of relative intensity of uorescein is de ned as the light intensity of 5 110nM uoresœin on a spot size with 5m m diameter. Inset shows the length of the ssD NA skeleton of nanocom partm ent and the com plan entary type. The background noise is within 1 unit.

as rigid cylinders, carrying helical and continuous line charges on their surfaces. Each DNA duplex carries the negative charge of phosphates plus a compensating positive charge from the adsorbed counterions. Let 0 < < 1 be the degree of charge compensation, f_1 , f_2 and f_3 the fractions of condensed counterions in the m inor and m a pr grooves ($f_1 + f_2 + f_3 = 1$). The mobile counterions in solution screen the C oulom b interactions between the two m olecules, causing at large separations an exponential decay of the latter with the D ebye screening length 1. The solvent is accounted for by its dielectric constant ". The structural parameters of B-DNA are half azim uthal width of the m inor groove $\tilde{s} = 0.4$, pitch H = 34A (g = 2 = H), and hard-core radius a = 9A. We take the following form for the pair interaction potential [15–18]:

$$u(R;) = u_{0} \overset{X^{1}}{f_{1}} + (1)^{n} f_{2} (1 f_{3}) \cos(n^{2})^{2}$$

$$\frac{(1)^{n} \cos(ng z) K_{0}(nR)}{(n^{2})^{n} (n^{2})^{2}} (1 f_{3}) \cos(n^{2})^{2}}$$
(4)

where R (> 2a) is the distance between the two parallel DNA molecules, z a vertical displacement, equivalent to a "spin angle" = gz. Here, $\mu_0 = 8 \quad {}^2=$ " ² (about 2:9k_B T=A at physiological ionic strength), and n = $\frac{1}{2} + n^2 g^2$. ngm (x;y) is given by


$$m_{m}(x;y) = \int_{j=1}^{X} K_{n,j}(x) K_{j,m}(y) \frac{I_{j}^{0}(y)}{K_{j}^{0}(y)}$$
(5)

with the modi ed Bessel functions $K_n(x)$ and $I_j(y)$. The primes denote derivatives. The sum rapidly converges, and it can be truncated after jnj = 2. Since $_nR > 3$ and g , each of the terms in the sum decreases exponentially at increasing R with the decay length $_n^1$ / 1=n.

Figure 2 present a scheme of interaction between two neighboring colum – nar DNA molecules charged with counterions on its surface. The distance between two DNA columns in our simulation is about 30A and the helical pitch of DNA molecule is about 36A. For brevity, we take the mean-eld approximation that the pair interactions mainly exist between charges in the same height.

$4\ \mbox{M}$ onte Carlo Sim ulation


Let t be the dimensionless variable to m ark the time series of simulation (t = 0;1;2:::) and T the environmental temperature. Assuming that M N DNAs are on the ADNC, the position of each DNA can be represented by its coordinates (x;y), where x;y 2 N, and 0 < x < M; 0 < y < N. AllDNA molecules in ADNC have identical sequence with P base pairs. Therefore there is the collection of M N P base pairs. The degree of freedom of the



F igure 2: The pair interaction between two parallelB-DNA double helixes. The black balls with positive signs in the center represent the absorbed positive counterions, while the little grey balls represent the phosphate carrying negative charges. Each DNA duplex carries the negative charge of phosphates with area density of 16:8 C = cm² plus a compensating positive charge com ing from the adsorbed counterions. We take the assumption that, and the distance between them are approximately 30A.

system is also M N P. The position of each base pair is thus represented by coordinates (x;y;i), where i 2 N; 0 < i < P. We take that the indices of base pairs is assigned from the bottom to the top of the DNA.

At the time t_0 , the state for an arbitrary base pair at $(x_0; y_0; i)$ with well-form ed hydrogen bonds is represented as $(x_0; y_0; i; t_0) = 1$. Contrarily, the state of a base pair with decoupled hydrogen bonds is denoted as $(x_0; y_0; i; t_0) = 0$ [10, 19]. (x; y; i; t) is a function of the time and the position of the base pair. Therefore, the state of each DNA molecules in ADNC can be represented by a sequence of digits. The num ber of all possible states is $2^{M N P}$.



F igure 3: A schem atic graph for a double-stranded DNA molecule associated with boolquantities to represent well-form ed hydrogen bonds (denoted as 1) and decoupled bonds (denoted as 0).

The simulation begins at t = 0, T = 0 C. At each step, t increases by 1, and state of base pair at $(x_0; y_0; i)$ is inverted, i.e. $(x_0; y_0; i; t + 1) = 1$ $(x_0; y_0; i; t)$. We assume that by changing the state of the system for M N P Z times, the system will approximate the equilibrium state in nitely. The change will be applied to each base pair for average Z times. Z is determined by experience and should be reasonable. We increase T by T during the simulation. Therefore we have the relation $T = \frac{t}{M + N + 2} T$.

W hether the state inversion is perm itted depends on the energy change ($E_{\rm t})$ in each step. The possibility of the state change at each step is

$$P((t)!(t+1)) = \begin{cases} 8 \\ \ge & 1 & \text{for } E_t & 0 \\ \vdots & e^{E_t = k_B T} & \text{for } E_t > 0 \end{cases}$$
(6)

If the current state change is perm itted, we keep up changing the system state at t+1. If the state change is forbidden by the possibility, the system state remains unchanged at t and waits for another change at t+1.

To achieve a relatively precise simulation, the change of the total energy at time t + 1 relative to that at the time t is analyzed by vecom ponents.

The recursion relation of energy change in each step is written as:

$$E_{t} = E(t+1)$$
 $E(t) = \begin{bmatrix} X^{5} \\ E_{1} \end{bmatrix}$ (7)

where E (t+1) and E (t) are the system energy for the instant t+1 and t respectively, and E_1 (l= 1;2;3;4;5) is the variation of the lth component.

The energy change depends on both the recursion relation of the base pair at $(x_0; y_0; i)$ and the states of its nearest neighbors. The global energy variation is determined by the local states around the base pair $(x_0; y_0; i)$. Following analysis presents the recursion relation of energy changes.

4.1. The hydrogen-binding energy (E_1)

This component of the energy consists of the M orse potential (equation (3)) and kinetic energy along the orientation of the hydrogen bonds. The binding energy is independent of the states of its neighboring base pairs.

$$\begin{array}{c} & 8 \\ \geq & J & \text{if} & (x_0; y_0; i; t) = 0 \\ & E_1 = \\ \geq & & \\ & J & \text{if} & (x_0; y_0; i; t) = 1 \end{array}$$
(8)

where J (J < 0) is the binding energy for each base pair that is in '1' state, while the binding energy for the '0' state is zero to be reference.

4.2. The stacking energy (E_2)

To simplify the calculation of the stacking energy shown in equation (2), we take into account the states of base pairs at $(x_0; y_0; i = 1; t)$ and $(x_0; y_0; i + 1; t)$. Their states remain unchanged during the interval from t to t+ 1. We employ the periodic boundary condition (PBC) listed below.

$$(x + M ;y;i;t) = (x;y;i;t) (x;y + N ;i;t) = (x;y;i;t) (x;y;i + P;t) = (x;y;i;t)$$
(9)

Therefore $(x_0; y_0; i 1; t)$ and $(x_0; y_0; i + 1; t)$ are both well de ned. The stacking energy re ects the interaction between nearest neighboring base pairs in same DNA, and it exists only when two nearest neighbors are in '1' state at the same time. We use the symbol $(x; y; i_1; i_2; \dots; i_n; t) = b_1; b_2; \dots; b_n$ to denote states in the same DNA for convenience, which means

 $(x;y;i_1;t) = b_1, (x;y;i_2;t) = b_2, \dots, (x;y;i_n;t) = b_n.$

$$E_{2} = \begin{pmatrix} 0 & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f000g} \\ 0 & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f010g} \\ w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f011g} \\ w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f011g} \\ w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f010g} \\ w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f110g} \\ w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f101g} \\ 2w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f101g} \\ 2w & \text{if} & (x_{0}; y_{0}; \text{i} & 1; \text{i}; \text{i} + 1; \text{t}) = \text{f101g} \\ \end{pmatrix}$$
(10)

where w is the stacking energy stored in two nearest neighboring base pairs in '1' state.

4.3. M orse potential away from equilibrium point ($\rm E_{3})$

W e set $E_1 = 0$ for the uncoupled hydrogen bond at base pairs. However, for a '0' state is next near to a '1' state in the sam eDNA strand, the distance between two base pairs is so close that the M orse potential should be taken into account. W e assigned energy E to every two nearest neighboring base pairs that are in di erent states in the sam eDNA.

$$E_{3} = \begin{pmatrix} 2E & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f000g \\ 2E & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f010g \\ 0 & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f001g \\ 0 & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f011g \\ 0 & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f100g \\ 0 & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f100g \\ 2E & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f101g \\ 2E & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f101g \\ 2E & \text{if} & (x_{0};y_{0}; 1 1; 1; 1 + 1; t) = f111g \end{pmatrix}$$
(11)

4.4. The e ectof excluded volume (E $_4$)

Q


8

The e ect of excluded volum e in the nature of DNA phase transition is discussed in Fisher's work [20]. The excluded volum e e ect is connected to the system entropy variation. The e ect is prone to separate two com plem entary strands in a double helix. We use F to represent the energy change corresponding to this e ect. One should notice $\frac{@F}{@T} < 0$. We then have

$$E_{4} = \begin{cases} F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f000g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f010g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f011g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f011g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f100g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f110g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f101g} \\ F & \text{if} & (x_{0}; y_{0}; i \quad 1; i; i + \; 1; t) = \text{f111g} \end{cases}$$

$$(12)$$

The energy changes discussed above are sum marized in the gure 4 below, which does not take into account the DNA-DNA interactions so far.



F igure 4: Schem e for the energy changes. We calculate the energy change under every case when the state of a base pair takes inversion. The value overhanging the double-arrow represents the energy variation when the state changes from left to right = $E_1 + E_2 + E_3 + E_4$.

4.5. DNA-DNA interaction potential (E_5)


We have introduced DNA-DNA interaction in previous section. For each base pair, we denote the state of its m nearest neighbors with $_{i}$, (i = 1;2;:::;m; $_{i}$ = 0;1). E₅ can be written as

$$E_{5} = \begin{cases} P_{m} & \text{if } (x_{0}; y_{0}; i; t) = 0 \\ P_{5} = P_{m} & \text{if } (x_{0}; y_{0}; i; t) = 1 \end{cases}$$
(13)

where G is the interaction energy between each pair of ions. Adding E 5 to , we will get the energy variation including the DNA-DNA interaction.

5.Results and Discussions

Following the equation (5) - (12), we could achieve a coarse-gained simulation of them elting curves of ADNC as well as that of DNA in bulk solutions. To perform the task, we choose suitable scale parameters to carry out the simulation: M = 100, N = 100, P = 20. The values of M and N chosen are much smaller than ones of the actual situation, which is up to 10^4 in the experiment. Since we take the periodic boundary condition, the values of M and N used do not change our result. The starting temperature is 0 C, and the naltemperature is 100 C, with increment of 0:01 C for each step. To guarantee the system reaches equilibrium state, we take state changes under a speci c tem perature. Each base pair has average 5 times to be changed. At each step, we count the number of DNAs that is still hybridized and calculate the percentage for dsDNA in ADNC. The simulation result shown in gure 5 shows a steep melting transition (hollow circles), consistent to the experimental observations. The simulated result without considering DNA-DNA interaction show in lled circles in gure 5 also agrees with the DNA melting curves in bulk solution. Comparison between the two cases suggests that the DNA-DNA interaction greatly increases the melting point of dsDNA chains.



F igure 5: Simulation results of melting curves for collective DNA molecules. The led circle represents the phase transition curve without considering the DNA-DNA interaction, while the hollow circle is the counterpart that takes into account the interaction. Parameters used in the simulations are: $J = 1900k_B$, $w = 250k_B$, $E = 850k_B$, $F = 1650k_B$, $\frac{GF}{GT} = 10k_B$, $G = 125k_B$.

In conclusion, we have established a simple coarse-gained model to sim – ulate the melting transition of DNA in ADNC. The result provides a reasonable explanation for our experimental observations. A lthough the simulation method discretizes the Morse potential and stacking energy proposed in Peyrard-Bishop model, the result still present a comparable approximation to experimental data due to our ne treatment of energy changes during melting transition. However, this work is only the beginning of insightful theoretical investigation for the rationality of ADNC. In future work, we will establish a more precise model to employ an extensive investigation of phase transition occurring in ADNC as well as its derived DNA machines.

A cknow ledgem ents

This work was partly supported by the grants from Chinese Natural Science Foundation, M inistry of Science and Technology of China and nancial support from Peking University.

Reference

- [1] Lifson S 1964 J. Chem. Phys. 40 3705
- [2] Poland D and Scheraga H A 1966 J. Chem. Phys. 45 1456
- [3] Poland D and Scheraga A H 1966 J. Chem. Phys. 45 1464.
- [4] Zhang Y L, Zheng W M, Liu J X, and Chen Y Z 1997 Phys. Rev. E 56 7100
- [5] Theodorakopoulos N, Dauxois T and Peyard M 2000 Phys. Rev. Lett. 85 6
- [6] Dauxois T and Peyard M 1995 Phys. Rev. E 51 4027
- [7] Dauxois T and Peyard M 1993 Phys. Rev. E 47 R44
- [8] Mao Y D, Luo C X, Deng W, Jin G.Y, Yu X M, Zhang Z H, Ouyang Q, Chen R S and Yu D P 2004 Nucleic Acids Res. 32 e144
- [9] Mao Y D, Luo C X and Ouyang Q 2003 Nucleic Acids Res. 31 e108
- [10] Zimm B H and Bragg J K 1959 J. Chem. Phys. 28 1246
- [11] Chrisey L A, Lee G U and O 'Ferrall C E 1996 Nucleic Acids Res. 24 3031
- [12] Bard A J and Fulkner L R 1980 Electrochem ical m ethods W iley, New York
- [13] Harreis H M, Korryshev A A, Likos C N, Lowen H, and Sutmann G 2002 Phys. Rev. Lett. 89 018303
- [14] Harreis H M, Likos C N, and Lowen H 2003 Biophys. J. 84 3607
- [15] Komyshev A A and Leikin S 1997 J. Chem. Phys. 107 3656
- [16] Komyshev A A 2000 Phys. Rev. E 62 2576
- [17] Allahyarov E and Lowen H 2000 Phys. Rev. E 62 5542
- [18] Komyshev A A 2001 Phys. Rev. Lett. 86 3666
- [19] HillT L 1959 J.Chem. Phys. 30 383
- [20] Fisher M E 1966 J. Chem. Phys. 45 1469