

Cerenkov Emission by Neutral Particles in Gravitoelectromagnetic Fields

X.Q. Li and S.Q. Liu

---

It is shown that under the post-Newtonian approximation the Einstein equations can be reduced to the standard Maxwell-type field equations in a medium; in such a context the Cerenkov emission by a neutral particle gives large energy loss while the particle moves at faster than the phase speed of waves in the medium.

---

KEY WORDS : Gravitoelectromagnetic equations; Emission by moving particles.

It has often been noted that magnetism can be understood as the consequence of electrostatics plus Lorentz invariance. Similarly, Newtonian gravity together with Lorentz invariance in a consistent way must include a gravitomagnetic field. This is the case of gravitoelectromagnetic (GEM) form of the Einstein equations in a medium [1-5]. In such a framework, the refractive index  $n^p$  of a medium is often is more than one [see (27), below]; note that the statement that only accelerated particles radiate is not correct when there are waves with  $n > 1$  [6]. Hence the Cerenkov emission by neutral particles in constant rectilinear motion is unavoidable.

It is well known that in the slow-motion limit of general relativity, accurate to post-Newtonian (PN) order, where the field equations can be reduced as [7]:

$$r^2 = 4G; \quad (1)$$

$$r^2 A = 16G v=c; \quad (2)$$

---

Department of Physics, Nanchang University, Nanchang, P.R.China; to whom correspondence should be addressed: sqliu@ncu.edu.cn

Moreover the harmonic gauge condition and the Lorentz-type force on a unit mass reduced to

$$\nabla_r A + \frac{4}{c} \frac{\partial \phi}{\partial t} = 0; \quad (3)$$

$$\frac{dv}{dt} = \nabla_r \left( \frac{1}{c} \frac{\partial A}{\partial t} + \frac{v}{c} \nabla_r A \right); \quad (4)$$

where  $\phi$  is the Newtonian gravitational potential,  $A = c = g_{i0}$  is the mixed metric and  $G$  is the Newton's constant.

One defines the GEM fields via

$$E_g = \nabla_r \left( \frac{1}{c} \frac{\partial A}{\partial t} \right); \quad \nabla_r A = B_g \quad (5)$$

in direct analogy with electromagnetism; it follows from these definitions the Maxwell-type field equations in the medium [8, 9]

$$\nabla_r B_g = \frac{16}{c} j_r + \frac{4}{c} \frac{\partial E_g}{\partial t}; \quad \nabla_r E_g = 4 G \rho; \quad (6)$$

$$\nabla_r E_g = \frac{1}{c} \frac{\partial B_g}{\partial t}; \quad \nabla_r B_g = 0; \quad (7)$$

where  $\rho = G^{-1}$  is "matter density" and  $j = G^{-1} v$  is "matter current". Then Eq.(4) becomes

$$\frac{dv}{dt} = E_g + \frac{v}{c} B_g; \quad (8)$$

And all terms of  $O(v^4=c^4)$  are neglected in the above analysis. Now one decompose the total matter density and current in equation (6):  $\rho = \tilde{\rho} + \rho_0; j = \tilde{j} + j_0$ , in which the  $\rho_0$  and  $j_0$  are the parts of a external field source in the medium, corresponding to a local additional mass disturbance, for example, by nonlinear interactions between the fields and the medium, or some test particles. In this case it is convenient to define the vector quantity  $D_g(t; r)$  through (in what follows one neglects the marks ' in  $\tilde{\rho}$  and  $\tilde{j}$  for simplicity)

$$D_g = E_g + 4 \int_{-z^t}^z dt^0 j(t^0; r); \quad (9)$$

by use of (9) and the mass continuity equation,

$$\frac{\partial}{\partial t} + r \quad v = 0 \quad ( = g;g_0) \quad ;$$

(6) and (7) are deduced to

$$r \quad B_g = \frac{16}{c} j_0 + \frac{4}{c} \frac{\partial D_g}{\partial t}; \quad r \quad D_g = 4 \quad g_0 \quad ; \quad (10)$$

$$r \quad E_g = \frac{1}{c} \frac{\partial B_g}{\partial t}; \quad r \quad B_g = 0 \quad ; \quad (11)$$

A dimensional analysis of equations (1), (5), (9) and (11) suggests that it is insightful to use the cgs units, since these field variables have the following dimensions:

$$[D_g] \quad [E_g] \quad [B_g] \quad [r] \quad [v] = s;$$

thus, one has

$$\frac{h}{E_g} = \frac{i}{G} \quad \frac{h}{B_g} = \frac{i}{G} \quad \frac{h}{E_g} = \frac{i}{G} \quad \frac{h}{v} = \frac{i}{v^2} \quad ;$$

and by substituting

$$E = \frac{E_g}{4} \frac{P}{G}; \quad B = \frac{B_g}{4} \frac{P}{G}; \quad D = \frac{D_g}{4} \frac{P}{G}; \quad (12)$$

$$j = \frac{j_0}{G} = \frac{P}{G} v; \quad b = \frac{P}{G} = \frac{P}{G} \quad ; \quad (13)$$

the GEM field equations can be expressed in the standard form

$$r \quad B = \frac{4}{c} j + \frac{1}{c} \frac{\partial D}{\partial t}; \quad r \quad E = \frac{1}{c} \frac{\partial B}{\partial t} \quad (14)$$

$$r \quad D = 0; \quad r \quad B = 0; \quad (15)$$

then Eqs.(9) and (8) become

$$D = E + \int_1^{z^t} dt^0 j(t^0; r); \quad (16)$$

$$\frac{dv}{dt} = \frac{p}{4G} E + \frac{v}{c} B \quad (17)$$

To consider the responses of the medium on the GEM fields, one must introduce material relation, which describes the GEM properties of the medium. In view of (16), the states of the medium not only depend on a given time-space point  $(t; r)$  but also depend on previous times and at any point of the medium. Hence, by general reasoning in physics (independent of a specific model for the medium) one can state that this is a non-local linear relation in the limit of linear response; whose Fourier representation is

$$j_i(t; k) = \epsilon_{ij}(t; k) E_j(t; k) \quad (18)$$

In special, for common continuous medium, where the "spatial dispersion" (dependency on  $k$ ) is not important, the relation (18) is deduced to  $(\epsilon_{ij}(t))$

$$j(t) = \epsilon(t) E(t);$$

ie. an "Ohm's gravitational law" [10], where  $\epsilon$  is "gravitational conductivity" [11] (for details see Refs [10,12]). Then by use of Eqs.(16) and (18), one has

$$D_i(t; k) = \epsilon_{ij}(t; k) E_j(t; k); \quad (19)$$

here  $\epsilon_{ij}$ , called the "dielectric tensor", is determined as

$$\epsilon_{ij}(t; k) = \epsilon_{ij} + \frac{i}{\omega} \chi_{ij}(t; k) \quad (\omega \neq 0) \quad (20)$$

In fact, if we consider the kinetic description for a self-gravitating system, on basis of Vlasor equation of the gravity [13,14], in which the force is Lorentz-type one [Eq.(17)], one can expand the distribution function  $f$  in powers of the field  $E$ ; and by Fourier transformation

$$A(r; t) = \int A(t; k) e^{i(t-k \cdot r)} d^3k; \quad A(t; k) = \int A(r; t) e^{i(t-k \cdot r)} \frac{d^3r}{(2\pi)^3}; \quad (21)$$

it yields the material relation (18) for the first order current  $j / \int^R v f_1(r; v; t) dp$ , in which  $\epsilon_{ij}(\omega; k)$  is relative to the particle distribution in the medium.

By applying the Fourier expansion, the field equation (14) becomes

$$\begin{aligned} i(k \cdot B)_k &= \frac{i\omega}{c} \epsilon_{ij} E_j + \frac{4}{c} j_{bi} \\ k \cdot E &= \frac{i\omega}{c} B; \end{aligned}$$

by elimination of B,

$$\epsilon_{ij}(\omega; k) E_j(\omega; k) = \frac{4}{i\omega} j_{bi}(\omega; k); \quad (22)$$

where

$$\epsilon_{ij}(\omega; k) = \frac{k^2 c^2}{\omega^2} \left( \frac{k_i k_j}{k^2} \epsilon_{ij} \right) + \epsilon_{ij}(\omega; k); \quad (23)$$

Multiplying Eq.(22) by any modular vector  $e_{k,ji}$ , with  $e_{k,ji} e_{k,ji} = 1$ ; one obtains

$$k^2 \frac{\omega^2}{c^2} \epsilon_k^T E_k = \frac{4}{c^2} [e_k \cdot j(\omega; k)]; \quad (24)$$

in which

$$\epsilon_k^T \epsilon_k = \epsilon_{ij}(\omega; k) e_{k,ji} e_{k,ij} + \frac{c^2}{\omega^2} (k \cdot k) \epsilon(k \cdot k); \quad (25)$$

and  $\epsilon_k$  is the "dielectric susceptibility". By neglecting the terms in right hand (the nonlinear terms), one gets from Eq.(24) the homogeneous wave equation; and from here one finds the dispersion equation

$$\epsilon_k \frac{k^2 c^2}{\omega^2} = 0; \quad (26)$$

Let  $\omega = \omega(k)$  be its a solution; physically, the progressive wave in the opposite direction is also possible, thus Eq.(26) has another solution with negative frequency  $\omega = -\omega(k)$ .

Consider the lowest order equation of motion for a test particle

$$\frac{dv}{dt} = \frac{p}{4G} E_g;$$

for plane waves  $e^{i\omega t + ik \cdot r}$ , one has:  $i\omega v_{k,i} = \frac{p}{4G} E_{k,i}$ , thus

$$j_{k,i} = \frac{p}{G} v_{k,i} = \frac{p}{G} \frac{p}{4G} E_{k,i} = E_{k,i};$$

ie.

$$j_{k,i} = i\omega E_{k,j}; \quad i\omega = \frac{G}{i} i\omega;$$

For transverse mode ( $k \cdot e = 0$ ), on the other hand, one has from Eq.(20) and Eq.(25)

$$\omega_k^2 = 1 + \frac{4G}{i^2} = 1 + \frac{4G}{i^2}; \quad (27)$$

with the "gravitational conductivity",

$$= i\omega(i\omega; k) e_{k,i}^t e_{k,j}^t;$$

Hence from Eq.(26) we obtain the dispersion relation

$$\omega^2 = k^2 c^2 - \omega_0^2; \quad \omega_0^2 = 4G; \quad (28)$$

$$\omega = kc - (\omega_0^2 - \omega^2)^{1/2};$$

To calculate this emission radiated by the moving external currents, we form the scalar product the first equation of Eq.(14) with  $E$ , of the second one with  $B$  and take the difference; then integrating over a volume and using the Gauss theorem, as a result we obtain

$$\frac{1}{4} \int_V \left[ E \cdot \frac{\partial D}{\partial t} + B \cdot \frac{\partial B}{\partial t} \right] dr + \frac{c}{4} \int_S (E \cdot B) dS = \int_V j \cdot E dr;$$

taking into account that the  $E$  and  $B$  vanish at infinity, thus

$$\frac{\partial W}{\partial t} = \frac{1}{4} [E \frac{\partial D}{\partial t} + B \frac{\partial H}{\partial t}] = \mathbf{j} \cdot \mathbf{E} : \quad (29)$$

i.e., the work done on the gravitoelectric fields per unit time, by the external currents, makes the GEM field energy density ( $[\mathbf{j}_0 \cdot \mathbf{E}] = [\mathbf{j}_g \cdot \mathbf{E}_g = G] = [\dot{\mathbf{v}}] = s$ ) get increments. Hence the GEM wave energy radiated is

$$U = \int_{-Z}^Z dt \int_{-Z}^Z dr E(r;t) \cdot j(r;t) : \quad (30)$$

Using the Fourier transforms of  $E$  and  $j$  and taking account of the condition of real field, we obtain the expression

$$U = 2 \int_{-Z}^Z dt \int_{-Z}^Z dk \operatorname{Re} [j(\mathbf{l};k) \cdot \mathbf{E}(\mathbf{l};k)] : \quad (31)$$

In view of Eq.(24),

$$U = 2 \int_{-Z}^Z dt \int_{-Z}^Z dk \frac{i c^2 \mathbf{j}(\mathbf{k}) \cdot \mathbf{j}(\mathbf{l};k)}{k^2 - \frac{1}{c^2} \omega_k^2} ;$$

by use of the Landau rule and the Plancherel formula [15],

$$\operatorname{Im} \frac{1}{k^2 - \frac{1}{c^2} \omega_k^2} = \operatorname{Im} \frac{1}{\omega_k^2 - k^2 - i0} = \pi \delta(\omega_k^2 - k^2) = \pi k^2 \delta(\omega_k^2 - k^2)$$

one has

$$\begin{aligned} U &= 2 \int_{-Z}^Z dt \int_{-Z}^Z dk \frac{1}{\omega_k^2 - k^2 - i0} \mathbf{j}(\mathbf{l};k) \cdot \mathbf{j}(\mathbf{l};k) = 2 \int_{-Z}^Z dt \int_{-Z}^Z dk \frac{\mathbf{j}(\mathbf{l};k) \cdot \mathbf{j}(\mathbf{l};k)}{\omega_k^2 - k^2 - i0} ; \\ &= 2 \int_{-Z}^Z dt \int_{-Z}^Z dk \frac{\mathbf{j}(\mathbf{l};k) \cdot \mathbf{j}(\mathbf{l};k)}{\omega_k^2 - k^2 - i0} ; \end{aligned} \quad (32)$$

where

$$\omega_k^2 = \omega_{l,k}^2 = \omega_{ij}(\mathbf{l};k) e_{k,i} e_{k,j} ; \quad (33)$$

the second term of Eq.(25) has no contribution to the integration in Eq.(32); and in the last step we consider the solutions  $\omega = \omega(k)$  of the dispersion equation,  $k^2 - \frac{\omega^2}{c^2} n^2 = 0$ ; have the same contribution to the integration.

On the one hand, one may consider any instantaneous velocity as uniform motion at short time intervals, and on the other hand the space should be approximately flat for the slow-motion and weak gravity, hence it is reasonable to study the emission radiated by a neutron particle with mass  $m$  in constant rectilinear motion passing through the medium. Then the current density is

$$j(r;t) = q_0 v(t) \delta(r - r(t)); \quad (34)$$

where  $q_0 = \frac{P}{Gm}$ ,  $r(t)$  is the position vector of the motion particle. Considering that  $r = r_0 + vt$ ;  $v = \text{const}$ ; the Fourier transform of Eq.(34) is

$$j(\omega;k) = \frac{q_0}{(2\pi)^3} v \exp(ik \cdot r_0 - i(\omega - kv)): \quad (35)$$

Because of the delta-function in Eq.(35), for the emission by a uniformly moving neutron particle has to be satisfied the Cerenkov condition:  $\omega = kv$ , or  $v > \omega/k = c/n(\omega)$ , which essentially corresponds to the laws of energy and momentum conservation in the radiation processes. In view of Eq.(27), then  $n(\omega) > 1$ ; therefore, the Cerenkov emission by the neutral particles in constant rectilinear motion is unavoidable.

Substituting the equation Eq.(35) into Eq.(32) the foregoing results yield readily

$$U = \int \frac{dk}{(2\pi)^3} \frac{2}{4\pi(2\pi)^3} \frac{j^t(k) j^t(\omega;k)}{\frac{1}{n^2(\omega)} \left[ \frac{d}{d\omega} n^2(\omega) \right]_{\omega=kv}} : \quad (36)$$

Using the following identity of delta-function

$$[\delta(\omega - kv)]^2 = \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \delta(\omega - kv)$$

and considering the definition of radiated power,  $I = \lim_{t \rightarrow \infty} \frac{U}{t}$  we obtain

$$I^t = \int \frac{2!^t(k) q_0^2 j^t(k) j^t(\omega;k)}{\frac{d}{d\omega} [n^2(\omega)]_{\omega=kv}} \left( \int k \frac{dk}{2} \right) : \quad (37)$$

For transverse waves the polarization vector  $e^t(k)$  in Eq.(37) has two independent components  $(e^{(1)}; e^{(2)})$ , then the total power radiated is obtained by adding the two parts, i.e., proportional to  $e^{(1)}(k) \frac{2}{v} + e^{(2)}(k) \frac{2}{v}$ ; and because  $e^{(1)}; e^{(2)}$  and  $\hat{k}$  ( $k=k$ ) constitute orthogonal frame in the complex space,  $e_i^{(1)} e_j^{(1)} + e_i^{(2)} e_j^{(2)} = \delta_{ij}$   $k_i k_j = k^2$ , one has

$$e^{(1)}(k) \frac{2}{v} + e^{(2)}(k) \frac{2}{v} = v^2 \hat{k} \cdot \hat{v} = v^2 \sin^2 \theta;$$

hence

$$I^t = \frac{G m^2 v^2}{c^2} \frac{d^4 \dots}{v > \frac{c}{n(\omega)}} \frac{c^2}{v^2 n^2(\omega)} \quad (38)$$

It is well known that the gravitational system can emit energy in quadrupole radiation; and the power radiated is proportional to the order of  $c^{-5}$ . But one sees from Eq.(38) that the Cerenkov emission occurs in the order of  $c^{-2}$ . Therefore, in general, the Cerenkov emission is more than the quadrupole radiation. For example, consider a binary system, in which the accreting compact object is a larger black hole with mass  $M = 10^8 M_\odot$  and companion star with mass  $m = M$  moves in nearly circular orbit around the black hole at the radius  $a_0 = 300 r_g$  ( $r_g = 2GM/c^2$ , the Schwarzschild radius). In such a case, from the Kepler law,  $(2\pi p)^2 = G(M+m)a_0^3$ , then one gets  $p = 1.56 \cdot 10^4$  (h); at the same time, we may take  $v = (2GM/a_0)^{1/2} = 1.6 \cdot 10^8$  (cm/s). Furthermore, one can estimate the medium density at  $r = a_0$ ,  $\rho = 2.9 \cdot 10^8 (a_0/3r_g)^{3-2} = 2.9 \cdot 10^5$  (g/cm<sup>3</sup>) [16]. As a result, due to the quadrupole radiation, the energy loss for the system is [17]

$$\frac{dE}{dt} = 3.0 \cdot 10^{33} \left(\frac{m}{M}\right)^2 \left(\frac{M}{M}\right)^{4-3} \left(\frac{p}{1 \text{ hour}}\right)^{10-3} = 1.4 \cdot 10^{30} \text{ (erg/s)};$$

here  $\rho = M m / (M + m)$ . And one gets from Eq.(38) the energy loss in Cerenkov radiation

$$I^0 = 3 \cdot 10^{33} \frac{m}{M} \frac{v}{10^9 \text{ cm/s}} \frac{1}{10^7} = 4.8 \cdot 10^{30} \text{ (erg/s)};$$

we have taken  $\omega = 10^7$  (s<sup>-1</sup>) at the above. Then  $n^2 = \frac{4G}{12} = 2.5 \cdot 10^8$ , thus the Cerenkov condition is satisfied:  $v > c/n(\omega)$ .

## ACKNOWLEDGMENT

This work was partly supported by the National Natural Science Foundation of China and the Natural Science Foundation of Jiangxi Province

## REFERENCES

- [1] Shapiro S.(1996). *Phys.Rev.Lett.* 77 , 4487.
- [2] Bonilla M .A .G .,and Senovilla J M .M .(1997) *Phys.Rev.Lett.* 78,783.
- [3] Senovilla J M .M .(2000) *Mod.Phys.Lett. A* 151,59.
- [4] Ruggiero M .L ., and Tartaglia A .(2002). *Nuovo C im ento B* 117,743.
- [5] Iorio L ., and Lucchesi D .M .(2003). *C lass. Q uantum G rav*20, 2477.
- [6] Melrose D .B . (1986) *Instabilities in space and laboratory plasmas*,  
Cambridge University Press,C am brige.
- [7] Weinberg S. (1972). *G ravitation and C osm ology*, John W iley & Sons, Inc,New York.
- [8] Braginsky V .B ., Caves C .M ., and Thome K ip S. (2003). *Phys.Rev. D* 15, 2045.
- [9] Braginsky V .B ., Pohlarev A .G ., and Thome K ip S.(1984). *Phys.Rev.Lett.*53, 863.
- [10] Ciobotariu C .(1991). *Phys.Lett.A* 158, 27.
- [11] Agop M ., Buzea C .G h., and Nica P.(2000). *Physica C* 339, 120.
- [12] Oprea I., and Agop M .(1998). *Studia G eoph.Et G eod.* 42, 431.
- [13] Li X .Q .(1990). *A stron A strophys*227, 317.
- [14] Li X .Q ., and Ma Y .H .(1993). *A stron A strophys*270, 534.
- [15] Lifshitz E .M ., and P itaevskii L .P.(1981). *Physical K inetics*, Pergam on, O xford.
- [16] Takahara F .(1979). *P rog.Theor.Phys.* 62, 629.
- [17] Peters P .C . and M athews J.(1963). *Phys.Rev.*131, 435.