## The Real Message in the Sky

Douglas Scott\* and J. P. Zibin<sup>†</sup> Department of Physics & Astronomy University of British Columbia, Vancouver, BC, V6T 1Z1 Canada

A recent paper by Hsu & Zee (physics/0510102) suggests that if a Creator wanted to leave a message for us, and she wanted it to be decipherable to all sentient beings, then she would place it on the most cosmic of all billboards, the Cosmic Microwave Background (CMB) sky. Here we point out that the spherical harmonic coefficients of the observed CMB anisotropies (or their squared amplitudes at each multipole) depend on the location of the observer, in both space and time. The amount of observer-independent information available in the CMB is a small fraction of the total that any observer can measure. Hence a lengthy message on the CMB sky is fundamentally no less observer-specific than a communication hidden in this morning's tea-leaves. Nevertheless, the CMB sky does encode a wealth of information about the structure of the cosmos and possibly about the nature of physics at the highest energy levels. The Universe has left us a message all on its own.

PACS numbers: 01.70.+w,95.10.-a,98.80.Bp,84.40.Ua,06.20.Jr

In a recent posting to the physics arXiv [1], Hsu and Zee (HZ) discuss the possibility that if a superior Being created a universe, and wished to inform the inhabitants of that universe of the existence of the creator, then an obvious place to put such a message would be in the anisotropies of the Cosmic Microwave Background (CMB, [2]). This would have the apparent benefit that the encoded information would be readable by all advanced civilisations in that universe. In this short Note we point out that the amount of information that the CMB can encode about fundamental physics is far less than HZ's estimate, and that almost all of the information in the CMB is strongly observer-dependent. While we agree that precision measurements of the CMB anisotropies are a worthwhile endeavour, we stress that there are quite Creator-independent reasons for this.

HZ propose that by tweaking the inflaton potential, a superior Being could encode a surprisingly large message of 100 kbits or so in the CMB  $C_\ell$  spectrum of anisotropies (roughly the size of the raw text file for this Note!). While it is certainly true that the particular realisation of the CMB sky that we observe contains a great deal of information, almost all of it describes the details of the fluctuations within the intersection of our past light cone with the last scattering surface, as we explain below.

The amplitude of primordial fluctuations generated by inflation is determined by the value of the inflaton potential and its slope while the modes are leaving the Hubble radius. However, the connection between the  $C_{\ell}$  spectrum and the inflaton potential is not direct for at least two important reasons. Firstly, the usual identification between the inflaton potential and the fluctuation spectrum P(k) can only be made when the slow roll approximation is valid. One cannot encode an arbitrary pattern

in the primordial spectrum with a corresponding tweak to the potential.

Another important reason the inflaton potential does not mapped directly to the  $C_{\ell}$  spectrum is that the primordial fluctuation spectrum in k-space undergoes a complicated convolution to form the CMB spectrum in  $\ell$ -space. The net result is a smoothed version of the k-space spectrum, which itself does not directly reflect the inflaton potential. Therefore HZ's suggestion, based on their estimate of  $C_{\ell}/\Delta C_{\ell} \simeq \sqrt{\ell}$  independent possible values for each  $\ell$  (where  $\Delta C_{\ell}$  is the inevitable cosmic variance), greatly overestimates the amount of information that can be encoded. No amount of tinkering with the potential can imprint an arbitrary pattern in the  $C_{\ell}$ s – the inherent smoothing results in strong correlations between neighbouring  $\ell$ s.

Furthermore, HZ assume that information can be encoded in the CMB spectrum up to  $\ell_{\rm max} \sim 10^4$ . In practice, the primary CMB power is greatly suppressed and smoothed above  $\ell \simeq 2000$  due to the finite thickness of the last scattering surface. Moreover, due to foregrounds, in practice one can only observe a fraction of the CMB sky, so  $\Delta C_{\ell}$  is somewhat higher. In addition to these problems, we already have measurements approaching cosmic-variance-limited accuracy on the first few hundred  $C_{\ell}$ s [3], and the measured spectrum is consistent with standard  $\Lambda$ CDM models [4]. No message exists in this range of  $\ell$  above the cosmic-variance noise.

It is straightforward to crudely model the effect of the smoothing of the  $C_{\ell}$  spectrum to obtain an improved estimate on the total information that it can encode. Suppose that the smoothing has a characteristic scale of  $\Delta \ell$ . In this case we could distinguish very roughly N distinct spectra above the cosmic variance, with

$$\ln N \sim \ln \prod_{i=1}^{\ell_{\text{max}}/\Delta \ell} \sqrt{i\Delta \ell} \simeq \frac{\ell_{\text{max}}}{2\Delta \ell} \ln(\ell_{\text{max}}/e). \tag{1}$$

Here the product is restricted to every  $\Delta \ell$ th value of  $\ell$ ,

<sup>\*</sup>Electronic address: dscott@phas.ubc.ca †Electronic address: zibin@phas.ubc.ca

and Stirling's approximation was used. Using the values  $\ell_{\rm max} = 1000$  and  $\Delta \ell = 10$  [5] then gives our *conservative* estimate of  $\ln N \sim 300$  bits.

HZ also propose that a superior Being may be advanced enough to directly set the spatial phases of the primordial modes, thereby imprinting a spatial pattern directly on the  $a_{\ell m}$  [6] coefficients of the CMB sky. But this contradicts the authors' intent of finding an observer-independent medium for a message. This is because the CMB sky is not a static billboard in space, but a consequence of the particular set of density variations on the last scattering surface, varying (slowly) in time, and different for each observer.

The specific values of the  $C_{\ell}$ s in the observed CMB angular power spectrum, let us call them  $a_{\ell}^2$ , arise from a projection of the 3-dimensional pattern of density perturbations on the 'last scattering surface' (a spherically shell, with finite thickness, surrounding every observer). In practice there are several contributions to the anisotropies, but for simplicity let us consider the large angle Sachs-Wolfe effect [7] for a flat cosmology:

$$a_\ell^2 \propto \int (dk/k^2) \delta_k^2 j_\ell^2 (kR_{\rm LS}),$$
 (2)

where  $\delta_k^2$  is the primordial power spectrum of density perturbations and  $R_{\rm LS}$  is the distance to last-scattering. The expectation for the  $a_\ell^2$  ( $C_\ell \equiv \langle a_\ell^2 \rangle$ ) is a smooth function of  $\ell$ , which can be calculated precisely for a given set of initial conditions (i.e.  $\langle \delta_k^2 \rangle$ ). However, the actual values of  $a_\ell^2$  depend on the specific  $\delta_k$ s which are sampled on each observer's last scattering surface. This means that different observers (separated by cosmological distances) will observe different values of the  $a_\ell^2$ s. This 'cosmic variance' does *not* limit the precision with which a set of  $a_\ell^2$ s can be determined, it just affects the ability to constrain the underlying expectation values, i.e. the  $C_\ell$ s.

Although one could imagine setting up a set of correlations in the sky [8] among  $a_{\ell m}$ s, it is clearly impossible to arrange for the same set of correlations to be projected on the last-scattering surface for *all* observers, while maintaining anything like statistical isotropy [9].

It's even worse than that though, since the  $a_{\ell m}$ s also change with the time of observation, through a combination of the 3-dimensional density perturbations evolving and the region of space being sampled by the last scattering surface moving. This means that any such communication presented on the CMB sky would be a message placed there for us, at the present epoch.

Hence a 'message in the sky' is no better than a signal hidden in the human genome, or indeed a billboard picture of a plate of spaghetti that a particular set of drivers pass on a specific set of days.

But are there other observer-independent ways of encoding a message? We have been unable to come up with any ideas which do not suffer from some of the same shortcomings as the CMB. We suggest that it may be impossible to think of any signal which a supreme Being could imprint on the universe they have created without it being targetted at particular civilizations or individuals. Given that faith is an important part of religion, then whether this would be a proof for or against the existence of an Intelligent Designer would appear to be a matter of taste. Indeed, we would imagine that any Creator would operate with greater subtlety than running advertisements on billboards.

HZ's suggestion that a supreme Being might encode the Grand Unified gauge group in the CMB sky is an interesting one, because in fact there is real hope that the CMB sky will provide information about Grand Unification, through observables which depend on the inflaton potential. The slope of the initial power spectrum, and particularly the amplitude of the gravity wave contribution (seen through the polarization 'B-modes') hold great promise for probing physics at energy scales  $> 10^{14}$  GeV. So there is indeed a 'message in the sky' – it is the fact that the CMB anisotropies allow us to determine very precisely the large-scale structure of the observable Universe [10], and to probe physics at the highest energies.

## Acknowledgments

We would like to thank the Universe for giving us the CMB [11].

<sup>[1]</sup> S. Hsu, A. Zee, "Message in the Sky", physics/0510102.

<sup>[2]</sup> For general reviews of the physics of CMB anisotropies, see M. White, D. Scott, J. Silk, Ann. Rev. Astron. Astrophys., 32, 319 (1994), W. Hu, S. Dodelson, Ann. Rev. Astron. Astrophys., 40, 171 (2002).

<sup>[3]</sup> D. Scott, G.F. Smoot, in 'The Review of Particle Physics', S. Eidelman et al., Phys. Lett. B592, 1 (2004); in press (2006).

<sup>[4]</sup> Apart from a few glitches, probably due to systematic effects, and in any case carrying very little information.

<sup>[5]</sup> Fast numerical codes for calculating the CMB anisotropy power spectrum use Δℓ as high as 50, e.g. cmbfast: U. Seljak, M. Zaldarriaga, Astrophys. J., 469, 437 (1996).

<sup>[6]</sup> The usual definition is  $T(\theta,\phi) \equiv \sum a_{\ell m} Y_{\ell m}(\theta,\phi)$ . The

 $a_{\ell m}$ s are assumed to be Gaussian random numbers with mean zero and variance  $C_{\ell}$ .

<sup>[7]</sup> R.K. Sachs, A.M. Wolfe, Astrophys. J., 147, 73 (1967).

<sup>[8]</sup> For example encoding the first digits of 3.14159264..., thereby giving ' $\Pi$  in the sky'.

<sup>[9]</sup> And this also ignores the fact that observers living within galaxies (like us within the Milky Way) will have a significant fraction of the sky blocked by bright foregrounds.

<sup>[10]</sup> For a review of the Standard Model of Cosmology, see D. Scott, astro-ph/0510731.

<sup>[11]</sup> This suggests to us the Cosmologist's Anthropic Principle, which states: 'By existing and doing cosmology for a living, we have selected from the multiverse those universes which allow us to find out about them.'