

Properties of quasi-one-dimensional molecules with Feshbach resonance interaction

V. A. Yurovsky

School of Chemistry, Tel Aviv University, 69978 Tel Aviv, Israel

(dated: March 31, 2022)

Bound states and collisions of atoms with two-channel two-body interactions in harmonic waveguides are analyzed. The closed-channel contributions to two-atom bound states become dominant in the case of a weak resonance. At low energies and values of the non-resonant scattering length the problem can be approximated by a one-dimensional resonant model. Three-body problem becomes nonintegrable and the properties of triatomic molecules become different from those predicted by the integrable Lieb-Liniger-McGwire model.

PACS numbers: 03.65.Ge, 03.65.Nk, 03.75.Be, 32.80.Pj

Introduction

Quasi-one-dimensional (1D) molecules, the existence of which was predicted in Ref. [1], have been recently observed in an experiment [2] with two-dimensional optical lattices. Except of lattices (see also Refs. [3, 4, 5]), tight cylindrical confinement, or atomic waveguides, have been realized in elongated atomic traps (see Refs. [6, 7, 8, 9] and references therein), and atomic integrated optics devices (see Refs. [10, 11] and references therein). Unlike weakly bound molecules in free, three-dimensional (3D), space (see Refs. [12, 13] and references therein), existing at positive elastic scattering length only, the quasi-1D ones survive at negative scattering length.

The interaction of two atoms in an atomic waveguide can be considered as a 1D zero-range interaction (see Ref. [14]) whenever the collision or binding energies remain small compared to the transverse waveguide frequency ω_\perp . Due to coupling to excited transverse states, the interaction strength demonstrates a resonant dependence on the ratio of the elastic scattering length a_{3D} to the transverse oscillator length

$$a_\perp = \frac{r_\perp}{m \omega_\perp}; \quad (1)$$

where m is the atomic mass. This confinement-induced resonance can be interpreted as a Feshbach resonance (see Refs. [1, 15]), where the excited transverse states play the role of a closed channel. The interaction strength demonstrates also a dependence on the collision energy, which is a common property of a resonant scattering. Such an energy dependence appears in the scattering amplitudes in Ref. [16], as well as in the bound state energy in Ref. [1]. However, the use of Feshbach resonance for tuning the elastic scattering length, as in Ref. [2], leads to additional energy dependence (see Ref. [18]), which can be important for a weak resonance. A similar effect has been considered in Refs. [19, 20, 21, 22] for a problem of two atoms under 3D harmonic confinement.

A Feshbach resonance appears when the collision energy of an atomic pair in the open channel lies in a vicinity of a bound (molecular) state in a closed channel (see

Ref. [17]). As a result, the quasi-1D molecules are superpositions of the closed channel state and the ground and excited waveguide modes of the open channel. The closed-channel contribution becomes substantial for weak resonances.

The theory of two-body two-channel problem under tight cylindrical harmonic confinement is summarized in Sec. I. This problem can be approximated by a 1D two-channel model. An improved relation between the 1D and 3D scattering parameters, presented in Sec. II below, increases the range of applicability of the 1D model compared to the relations in Ref. [18]. The composition of quasi-1D bound states is analyzed in Sec. III. Three-body 1D molecules are considered in Sec. IV in a way which is similar to the analysis of scattering in Ref. [23]. A system of units with $\hbar = 1$ is used below.

I. FESHBACH RESONANCE IN HARMONIC WAVEGUIDES

The properties of two-atom systems can be described by close-coupled equations for the wavefunction of the open channel $\psi_a(r)$ and the amplitude for the system to be in the closed channel ψ_m , of the form (see Ref. [18]),

$$\begin{aligned} E \psi_a(r) &= \frac{1}{m} r^2 + V_a(r) + V_{\text{conf}}(r) \psi_a(r) \\ &\quad + V_{\text{am}}(r) \psi_m \\ E \psi_m &= D_{3D} \psi_m + V_{\text{am}} \psi_a(0); \end{aligned} \quad (2)$$

Here E and r are, respectively, the energy and coordinate vector of the relative motion and all the energies are measured from the open channel threshold. For a harmonic waveguide the confinement potential has the form

$$V_{\text{conf}} = \frac{m}{4} \omega_\perp^2 z^2; \quad (3)$$

where ρ and z are the cylindrical components of the vector r . The strength of the open channel potential V_a , the coupling strength V_{am} , and the bound state energy in the closed channel D_{3D} can be expressed as (see Ref. [18])

$$V_a = \frac{4}{m} a_{3D}^{-1} \left(1 - \frac{2}{a_{3D} \rho_c} \right)^{-1}$$

$$\mathcal{V}_{\text{am}} = \frac{4}{m} a_{3D} \left(1 - \frac{2}{a_{3D} p_c} \right)^2 \quad (4)$$

$$D_{3D} = B - B_0 + \frac{1}{2} a_{3D} p_c$$

in terms of the phenomenological resonance strength, the difference between the magnetic momenta of an atomic pair in the open and closed channels, the detuning of the external magnetic field B from its resonant value B_0 , and the momentum cutoff p_c . The final results reached below are derived in the limit $p_c \rightarrow \infty$. The δ -function potentials are applicable to two indistinguishable bosons, as well as to bosons or fermions with different spins.

Elimination of m from Eqs. (2) leads to a single equation for $a(r)$. It can be expanded in terms of the transverse Hamiltonian eigenfunctions χ_{0i} with the zeroth angular momentum projection on the waveguide axis z as

$$a(r) = \sum_{n=0}^{\infty} \left(\frac{r}{a_?} \right)^{2n} \sum_{q=0}^{\infty} d_{nq} \chi_{nq}(r) \quad (5)$$

The coefficients d_{nq} satisfy the set of coupled equations (see Ref. [18])

$$\frac{p_n^2}{m} d_{nq} = \frac{1}{2^{2n} a_?^2} V_e(E) \sum_{n^0=0}^{\infty} d_{n^0q} \quad (6)$$

where

$$V_e(E) = V_a + \frac{\mathcal{V}_{\text{am}}}{E - D_{3D}} \quad (7)$$

is a non-normalized energy-dependent interaction strength and

$$p_n = \frac{p}{m} \sqrt{E - (2n+1)\epsilon_?} \quad (8)$$

is the relative axial momentum for the channel corresponding to the transverse excited state χ_{nq} with the excitation energy $(2n+1)\epsilon_?$.

The transition matrix for a two-atom collision in an atomic waveguide has been derived in Ref. [18] as

$$T_{\text{conf}}(p_0) = \frac{4}{m a_?} \frac{a_?}{a_e(E)} + \frac{1}{2} \frac{a_? p_0}{2} \quad (9)$$

Although the collision momentum p_n depends on the channel, the transition matrix is independent of the initial and final transverse states n, n^0 for all open channels ($n, n^0 < (E - \epsilon_?)^{1/2} = 2$) and is expressed by Eq. (9) in terms of p_0 . It is a consequence of the use of zero-range potentials in Eqs. (2). The energy-dependent length

$$a_e(E) = a_{3D} \left(1 + \frac{B - B_0}{E} \right) \quad (10)$$

replaces the elastic scattering length in Bethe-Peierls boundary condition. The Hurwitz zeta function is defined as (see Refs. [16, 24]),

$$\zeta(s, \nu) = \sum_{n=0}^{\infty} \frac{1}{(n + \nu)^s} \quad (11)$$

with $2 < \arg(\nu) < \infty$.

II. RELATION TO THE ONE-DIMENSIONAL PROBLEM

The confined two-body problem can be interpreted as a 1D one described by the Schrodinger equation

$$E_c' a(z) = \frac{1}{m} \frac{d^2 a}{dz^2} + U_e(E_c) a(z) \quad (12)$$

with a zero-range interaction, where the interaction strength U_e depends on the collision energy $E_c = p_0^2/m$. The 1D transition matrix corresponding to Eq. (12),

$$T_{1D}(p_0) = U_e^{-1}(E_c) + \frac{im}{2p_0} \quad (13)$$

coincides with Eq. (9) for

$$U_e(E_c) = \frac{1}{2! a_e(E_c + \epsilon_?)^2} + \frac{m a_?}{4} \frac{1}{2!} \frac{E_c}{2!} \frac{im}{2p_0} \quad (14)$$

The case of low collision energies $E_c \rightarrow 0$ can be analyzed using the expansion (see Refs. [14, 16])

$$\frac{1}{2!} \frac{1}{a_e(E_c + \epsilon_?)^2} = \frac{C}{C^0} \frac{1}{2!} \frac{1}{a_e(E_c + \epsilon_?)^2} \quad (15)$$

where $C = \frac{1}{2} \Gamma(1/2) \Gamma(3/2) \approx 1.4603$, $C^0 = \frac{1}{2} \Gamma(3/2) \approx 1.3062$, and $\Gamma(x)$ is the Gamma function (see Ref. [24]).

Following expressions attain a simpler form written in terms of dimensionless parameters (the scattering momentum k , the elastic scattering length a , the detuning b^0 , and the resonance strength d), defined as

$$k = \frac{p_0 a_?}{2}; \quad a = \frac{a_{3D}}{a_?}; \quad b^0 = \frac{B - B_0}{2! \epsilon_?}; \quad d = \frac{a_{3D}}{2a_? \epsilon_?} \quad (16)$$

The parameters k , a , and d have been used previously in Ref. [18]. The detuning b^0 , measured from the crossing point of the closed-channel bound state and the continuum threshold ($D_{3D} = \epsilon_?$), is related to the detuning b of Ref. [18], measured from the scattering length resonance, as $b^0 = b - d = a$.

Substitution of the expansion (15) into Eq. (14) leads to the expression of the interaction strength in the form

$$U_e^{-1}(E_c) = \frac{4}{m a_?} \frac{a}{C^0 a k^4 + k^2 (1 - C a) b - d a} \quad (17)$$

where

$$= 1 - C a - C^0 a b^0: \quad (18)$$

In a wide range of the parameters the interaction strength can be approximately expressed as

$$U_e(E_c) = U_a + \frac{2\hbar^2 g^2}{E_c D_{1D}}: \quad (19)$$

This form corresponds to a two-channel 1D problem (see Ref. [18]), described by the coupled equations

$$E'_0(z) = \frac{1}{m} \frac{d^2 \psi_0}{dz^2} + U_a \psi_0(0) + \frac{\hbar}{2g} \psi_1^{am}(z) \quad (20)$$

$$E'_1^{am} = D_{1D} \psi_1^{am} + \frac{\hbar}{2g} \psi_0(0)$$

for the open- and closed-channel coefficients $\psi_0(z)$ and ψ_1^{am} , respectively. The non-resonant interaction strength U_a , the channel coupling g , and the detuning D_{1D} will be further related to the 3D scattering parameters. Equation (12) with U_e given by Eq. (19) can be obtained by the elimination of the closed channel from Eqs. (20).

The ratio of the first term in the denominator of Eq. (17) to the other terms does not exceed the order of magnitude of ak^2 . It can be neglected whenever $a < 1$ and $k \ll 1$, leading to an expression of the form of Eq. (19) with

$$U_a = \frac{4a}{m a_2}; \quad \hbar^2 g^2 = 4\hbar^2 \frac{d + C^0 (ab^0)^2}{m a_2^2} \\ D_{1D} = 2\hbar^2 \frac{(1 - C a) ab^0 + d}{a}: \quad (21)$$

For the case of a relatively small detuning, $C^0 \hbar^2 ab^0 \ll \hbar^2 C a$, or

$$\hbar^2 (B - B_0 + \hbar^2) \ll \hbar^2 \frac{a_2 \hbar^2}{a_{3D}} - 1 - C \frac{a_{3D}}{a_2} \hbar^2 \quad (22)$$

the terms proportional to b^0 in the parameter can be neglected. The parameters U_a and D_{1D} are expressed then by Eqs. (43) and (45) in Ref. [18], while Eq. (44) therein will be valid whenever $d \ll a^2 b^0$, or $2(B - B_0 + \hbar^2) \ll a_2 \hbar^2 \ll a_{3D} \hbar^2$.

Unlike Eqs. (43)–(45) in Ref. [18], the relations (21) demonstrate a non-linear dependence of all three parameters U_a , g , and D_{1D} on the elastic scattering length, detuning, and resonance strength. These relations substantially increase the applicability range of the 1D approximation, as is demonstrated by Fig. 1 using the example of binding energy $E_b = \hbar^2 E = 2\hbar^2 x^2$. The parameter x here is the ratio of a_2 to the bound state axial size. For the confined system it is determined as a solution of the transcendent equation (see Eq. (55) in Ref. [18])

$$\frac{ax^2 + ab^0 + d}{x^2 + b^0} = \hbar^2 \frac{1}{2} x^2; \quad x > 0; \quad (23)$$




FIG. 1: The binding energy E_b calculated as a function of the dimensionless resonance strength d [see Eq. (16)] at the scattering length resonance $B = B_0 + \hbar^2$, or $b^0 = d = a$, with the dimensionless non-resonant interaction strength $a = 0.1$. The solid, dashed, and dot-dashed lines present, respectively, solutions of the exact equation (23), the 1D approximation (24), and the open-channel model (25).

while for the related 1D system it is evaluated as a solution of the cubic equation

$$x^3 + ax^2 + [(1 - C a)b^0 + d = a]x + ab^0 = 0: \quad (24)$$

This equation determines poles of the 1D T-matrix (13) with the interaction strength (19) and the resonance parameters given by Eq. (21). A similar equation has been considered in Ref. [25].

Substitution of the resonant scattering length a_{3D} [$\hbar^2 = (B - B_0 + \hbar^2)$], which takes into account the resonance shift by \hbar^2 due to confinement, into the equations of Ref. [1], leads to the following equation for the parameter x

$$\frac{ab^0 + d}{b^0} = \hbar^2 \frac{1}{2} x^2; \quad x > 0; \quad (25)$$

This approximation, corresponding to a single-channel confined problem with an energy-independent interaction, is called here the "open-channel model". Figure 1 demonstrates that this model is applicable to strong resonances only.

Various approximations for the binding energy are compared in Fig. 2 for a strong resonance. The results demonstrate good agreement between solutions of the exact equation (23), the 1D approximation (24), and the open-channel model (25) for $B < B_0 + \hbar^2$, when a weak bound state exists. However, the open-channel model predicts a non-physical singularity at $B = B_0 + \hbar^2$. Higher above the resonance the result of the open-channel model tends to the energy of the deep quasi-3D bound state. The latter state is not described




FIG. 2: The binding energy E_b calculated as a function of the external magnetic field for the 202 G resonance in K with $\Delta = 69.2$ KHz. The solid line, pluses, and crosses present, respectively, solutions of the exact equation (23), the 1D approximation (24), and the open-channel model (25).




FIG. 3: The binding energy E_b calculated as a function of the external magnetic field for the 543 G resonance in ^6Li with $\Delta = 200.2$ KHz. The solid line, pluses, and crosses present, respectively, solutions of the exact equation (23), the 1D approximation (24), and the open-channel model (25).

by the 1D approximation. The exact equation (23) gives correctly both the deep and weak bound states.

The 1D approximation agrees to the exact equations also in the case of a weak resonance (see Fig. 3). However the open-channel model, neglecting energy dependence of the interaction strength, is applicable in this case only within a small interval of B close to $B_0 + \Delta =$.

The applicability range of the 1D approximation is improved because the approximation (15) has a root at $\alpha = 0.294$, close to exact value of 0.303 (cf. the value of 0.468 provided by two-term approximation used in Ref.

[18]). The accuracy of the binding energy demonstrates the applicability of the 1D approximation to the σ -shell T-matrix, which is important for applications to many-body problems (see also discussion in Ref. [26]). The only criteria of applicability are $\alpha < 1$ and $E_b \ll \Delta$ (or $E_c \ll \Delta$ for collisions).

Equation (17) can demonstrate an energy-dependence of the form of Eq. (19) for a weak resonance, whenever $\Delta \gg \alpha \Delta$, or

$$\mathcal{B} \approx B_0 + \Delta = j \quad (26)$$

and the term $\Delta \alpha$ in the denominator of Eq. (17) can be neglected. In this case the 1D parameters can be estimated as

$$U_a \approx 0; \quad \mathcal{D}_{1D} \approx \frac{4\Delta}{C_0 m a^2}; \quad \mathcal{D}_{1D} \approx 2\Delta \frac{1}{C_0 a}. \quad (27)$$

They are independent of the Feshbach resonance parameters. Therefore, in agreement with Ref. [1], a confinement-induced resonance can be interpreted as a two-state Feshbach resonance even for non-resonant 3D scattering. It is a consequence of the approximation (15). It should be noted that the closed channel involves in this case a superposition of all transverse excitations. However, the detuning \mathcal{D}_{1D} then substantially exceeds the transverse frequency Δ and the energy-dependence is very weak in the quasi-1D regime, whenever the energy is less than Δ .

III. BOUND STATE COMPOSITION

The bound states of two atoms in atomic waveguides are superpositions of the closed and open channels of the Feshbach resonance. The open-channel component is a superposition of all transverse modes. The size of the closed-channel component is negligibly small compared to a , this component is not affected by the waveguide, and does not need an expansion in terms of the transverse modes.

For a bound state with energy $E < \Delta$ all momenta p_n , defined by Eq. (8), are imaginary and the solution of Eq. (6) has the form

$$\tilde{\psi}_n(q) = \frac{C_a}{p_n^2 + q^2}. \quad (28)$$

The probability to find the molecule in the n -th transverse mode of the open channel can then be expressed as

$$W_n = \frac{C_a^2}{2 p_n^2}. \quad (29)$$

Equations (2) and (5) allow to relate the closed-channel amplitude ψ_m to the open-channel wavefunction as

$$\psi_m = \frac{V_{am}}{E - D_{3D}} \frac{C_a}{2 a^2} \sum_{n=0}^{\infty} \tilde{\psi}_n(q) dq. \quad (30)$$

Although the sum here diverges, the renormalization procedure of Ref. [18] leads to a finite probability to find the molecule in the closed channel

$$W_c = \sum_m \sum_j = \frac{\sum_a \sum_j a_j^2}{2m a_{3D} [E - (B - B_0)]^2}: \quad (31)$$

The final results can be expressed in terms of the dimensionless parameters (16)

$$\frac{W_c}{W_0} = \frac{2dx^3}{a^2(x^2 + b^2)^2}; \quad \frac{W_e}{W_0} = x^3 - \frac{3}{2}x^2 - 1; \quad (32)$$

where x is the solution of Eq. (23), $W_e = \sum_{n=1}^p W_n$ is the total contribution of all excited transverse states, and the contributions are normalized as $W_0 + W_e + W_c = 1$. An approximate expression, used in following calculations,

$$1 - \frac{\frac{3}{2}x^2 - \frac{1+2x^2}{x^3}}{0.5 + 0.54884x + 2.5636x^2 + 0.12172x^3 + 4x^4}$$

gives a relative error of less than $3 \cdot 10^{-4}$.

Bound states of the related 1D system are superpositions of open and closed channels, where the closed channel effectively incorporates contributions of the excited transverse states and the closed channel of the confined system. The two contributions can be respectively expressed as

$$W_c^{1D} = \sum_1^{\infty} \int_0^{\infty} |f_1^{\text{am}}|^2; \quad W_o^{1D} = \sum_1^{\infty} \int_0^{\infty} |f_0(z)|^2 dz; \quad (33)$$

where the bound-state solution of Eqs. (20) has the form

$$f_1^{\text{am}} = \frac{p}{E - D_{1D}} \frac{2g'_0(0)}{D_{1D}}; \quad f_0(z) = f_0(0) \exp \left(-\frac{p}{m E_b} \sum_j j \right); \quad (34)$$

The ratio of the contributions can be expressed as

$$\frac{W_c^{1D}}{W_o^{1D}} = 2x^3 \frac{d + C^0 a^2 b^2}{a^2 (x^2 + b^2)^2}; \quad (35)$$

A direct evaluation demonstrates that

$$\frac{W_c + W_e}{W_0} \xrightarrow{x \rightarrow 0} \frac{W_c^{1D}}{W_o^{1D}}, \text{ where } \frac{3}{2}x^2 \xrightarrow{x \rightarrow 0} x^3 + \frac{3}{2}; \quad (36)$$

The contributions to the bound states are presented in Fig. 4 for two resonances: the strong one in K with $ad = 1.6$ and the weak one in ^6Li with $ad = 4.1 \cdot 10^{-4}$. The results demonstrate that $W_c^{1D} = W_e + W_c$ and $W_o^{1D} = W_0$ with a good accuracy over a wide range of parameter values. The closed channel and excited states of the open channel yield the dominant contribution far below the




FIG. 4: (color online) The contributions of the ground (dot-dashed line) and excited (dashed line) states of the open channel and of the closed channel (solid line) to the weak bound state calculated as a function of the external magnetic field. The dot-dot-dot-dashed and dotted lines present, respectively, contributions of the open and closed channels in the 1D model. The parts (a) and (b) are related, respectively, to the 202 G resonance in K with $ad = 69.2$ KHz and to the 543 G resonance in ^6Li with $ad = 200.2$ KHz. The dot-dashed and dot-dot-dot-dashed lines almost coincide in the part (b).

bound state threshold at $B = B_0 + \frac{1}{2} \mu_B$, while the ground state of the open channel becomes dominant near the threshold. For the strong resonance the contribution of the closed channel in the confined system is always small, and the bound state consists mostly of the ground and excited states of the open channel. However, for the weak resonance the major contribution is yielded by the closed channel and the ground state of the excited channel. The closed channel and excited states of the open channel yield similar contributions far below the bound state threshold for $ad = m a_{3D}^2 = 4 \cdot 0.1$. The resonances with higher or lower values of the product ad can be called, respectively, as open-channel or closed-

Figure 5 presents the scaled binding energy $\epsilon_3 = \epsilon_m^{-1/3} |\epsilon_j|^{4/3} E$ calculated by a numerical solution of Eq. (41). A related problem with non-resonant interactions only, the Lieb-Liniger-McGuire model [28, 29], has an exact solution. The binding energies for two- and three-body bound states in that model are expressed as $\epsilon_2 = \epsilon^2/2$ and $\epsilon_3 = 2\epsilon^2$, respectively, and therefore $\epsilon_3 = 4\epsilon_2$. In the present resonant case $\epsilon_3 < 4\epsilon_2$ only at large positive detunings. For large negative detunings the two-body bound state contains mostly the 1D closed-channel contribution, and the three-body bound state has a form of the two-body state with a third atom weakly bounded to it.

Conclusions

Two atoms with a Feshbach resonant interaction in an atomic waveguide can be approximated by a 1D reso-

nant model at low energies and values of the non-resonant scattering length. In the case of a strong resonance two-atom bound states contain mostly the contributions of the open channel. The closed channel contribution becomes dominant in weak resonances, such that the 543 G resonance in ^6Li . In the case of triatomic molecules a resonant interaction leads to properties different from those predicted by the Lieb-Liniger-McGuire model.

Acknowledgments

The author is very grateful to Maxim Olshanii and Abraham Ben-Reuven for helpful discussions and to Andrea Simonini for providing parameters [30] of the Feshbach resonance in ^6Li .

-
- [1] T. Bergeman, M. G. Moore, and M. Olshanii, Phys. Rev. Lett. 91, 163201 (2003).
 - [2] H. Moritz, T. Stoferle, K. Guenter, M. Kohl, and T. Esslinger, Phys. Rev. Lett. 94, 210401 (2005).
 - [3] M. Greiner, I. Bloch, O. Mandel, T. W. Hansch, and T. Esslinger, Phys. Rev. Lett. 87, 160405 (2001).
 - [4] T. Kinoshita, T. Wenger, and D. Weiss, Science 305, 1125 (2004).
 - [5] C. D. Fertig, K. M. O'Hara, J. H. Huckans, S. L. Rolston, W. D. Phillips, and J. V. Porto, Phys. Rev. Lett. 94, 120403 (2005).
 - [6] A. Gorlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, Phys. Rev. Lett. 87, 130402 (2001); A. E. Leanhardt, A. P. Chikkatur, D. K. Jin, Y. Shin, T. L. Gustavson, W. Ketterle, and D. E. Pritchard, Phys. Rev. Lett. 89, 040401 (2002).
 - [7] K. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, Nature 417, 150 (2002).
 - [8] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon, Science 296, 1290 (2002).
 - [9] S. Richard, F. Gerbier, J. H. Thywissen, M. Hugbart, P. Bouyer, and A. Aspect, Phys. Rev. Lett. 91, 010405 (2003); M. Hugbart, J. A. Retter, F. Gerbier, A. Varon, S. Richard, J. H. Thywissen, D. Clement, P. Bouyer, and A. Aspect, physics/0501456 (2005).
 - [10] R. Föllman, P. Krueger, J. Schmiedmayer, J. Denschlag, and C. Henkel, Adv. At. Mol. Opt. Phys. 48, 263 (2002).
 - [11] Y. Shin, C. Sanner, G.-B. Jo, T. A. Pasquini, M. Saba, W. Ketterle, D. E. Pritchard, M. Vengalattore, and M. Prentiss, cond-mat/0506464 (2005).
 - [12] K. Goral, T. Kohler, S. A. Gardiner, E. Tiesinga, and P. S. Julienne, J. Phys. B 37, 3457 (2004).
 - [13] C. Chin, cond-mat/0506313.
 - [14] M. Olshanii, Phys. Rev. Lett. 81, 938 (1998).
 - [15] M. Girardeau, Hieu Nguyen, and M. Olshanii, Optics Communications 243, 3 (2004).
 - [16] M. Moore, T. Bergeman, and M. Olshanii, J. Phys. (Paris) IV 116, 69 (2004).
 - [17] E. Timmermans, P. Tommasini, M. Hussein, and A. Kerman, Phys. Rep. 315, 199 (1999).
 - [18] V. A. Yurovsky, Phys. Rev. A 71, 012709 (2005).
 - [19] E. Tiesinga, C. J. Williams, F. H. Mies, and P. S. Julienne, Phys. Rev. A 61, 063416 (2000).
 - [20] D. Blume and C. H. Greene, Phys. Rev. A 65, 043613 (2002).
 - [21] E. L. Borda, E. Tiesinga, and P. S. Julienne, Phys. Rev. A 66, 013403 (2002); E. L. Borda, E. Tiesinga, and P. S. Julienne, Phys. Rev. A 68, 032702 (2003).
 - [22] M. G. Moore, cond-mat/0506383 (2005).
 - [23] V. A. Yurovsky, A. Ben-Reuven, and M. Olshanii, physics/0512033.
 - [24] H. Bateman and A. Erdelyi, Higher Transcendental Functions, Vol. 2 (McGraw-Hill, New York, 1953).
 - [25] K. V. Khurutsyan and P. D. Drummond, Phys. Rev. A 58, 2488 (1998).
 - [26] A. Abdurakhimov and A. L. Zubarev, Z. Phys. A 322, 523 (1985).
 - [27] W. G. Lockett, The Quantum Mechanical Few-Body Problem (Springer, Berlin, 1983).
 - [28] E. H. Lieb and W. Liniger, Phys. Rev. 130, 1605 (1963).
 - [29] J. B. McGuire, J. Math. Phys. 5, 622 (1964).
 - [30] A. Simonini (unpublished).