

Laying the ghost of the relativistic temperature transformation

Peter T. Landsberg* [†] and George E.A. Matsas[‡]

Instituto de Física Teórica, Universidade Estadual Paulista

Rua Pamplona 145, 01405-900–São Paulo, SP

Brazil

Abstract

Using the Unruh-DeWitt detector, it is shown that a universal and continuous Lorentz transformation of temperature cannot exist for black-body radiation. Since any valid Lorentz transformation of temperature must be able to deal with black-body radiation, it is concluded that a universal and continuous temperature transformation does not exist.

*Permanent address: Faculty of Mathematical Studies, University of Southampton, Southampton SO17 1BJ, England.

[†]e-mail: ptl@maths.soton.ac.uk

[‡]e-mail: matsas@axp.ift.unesp.br

This note represents yet another attempt to lay the ghost of the relativistic temperature transformation which has motivated a number of papers by the founders of relativity and quantum mechanics [1]– [5], more than 60 contributions during 1963-1968 [6] (see e.g. [7]– [15]), and a continuing trickle ever since (see e.g. [16]– [28] and references therein). The new result which makes the present considerations worthwhile is the explicit formula for the excitation rate of an Unruh–DeWitt detector, *i.e.* a two-level monopole [29], which has a single proper energy gap $\hbar\omega'$, and detects massless scalar particles or, equivalently, “spinless photons”. Suppose that black-body radiation with proper temperature T is at rest in some inertial frame S . The excitation rate of an Unruh-DeWitt detector moving with constant velocity v through it is found, with the aid of quantum field theory, to be proportional to the particle number density [26]

$$n'(\omega', T, v)d\omega' = \frac{\omega'kT\sqrt{1-v^2/c^2}}{4\pi^2c^2v\hbar} \ln \left[\frac{1 - e^{-(\hbar\omega'\sqrt{1+v/c})/(kT\sqrt{1-v/c})}}{1 - e^{-(\hbar\omega'\sqrt{1-v/c})/(kT\sqrt{1+v/c})}} \right] d\omega'. \quad (1)$$

Here we use “'” to refer to quantities as measured in S' , which is the inertial frame in which the detector is assumed to be at rest. This result, while reducing to the Planckian form

$$n(\omega, T)d\omega = \frac{\omega^2/c^3}{2\pi^2(e^{\hbar\omega/kT} - 1)} d\omega \quad (2)$$

in the limit $v \rightarrow 0$, *has not itself this form.*

By a black body we mean a system which has a Planckian spectrum, and by black-body temperature the parameter which characterizes this spectrum; so if one looks for the temperature of a black-body as defined in the moving frame S' , one needs to express (1) in the form

$$n'_{bb}(\omega', T')d\omega' = \frac{\omega'^2/c^3}{2\pi^2(e^{\hbar\omega'/kT'} - 1)} d\omega', \quad (3)$$

for some continuous function $T' = T'(T, v)$. If one succeeds, then one would be able to say that temperature transforms under boosts according to the law $T' = T'(T, v)$. *However, such a transformation is easily shown mathematically not to exist.* In particular, the popular transformations

$$T'(T, v) = \gamma^a T \quad (4)$$

where $\gamma \equiv (1 - v^2/c^2)^{-1/2}$, and $a = -1$ (see e.g. [1]- [5], [11], [28]), $a = +1$ (see e.g [7], [9]) or $a = 0$ (see e.g. [12], [15], [21]) are useless in this context because, again, they do not reduce (1) to (3). Since any universal continuous relativistic temperature transformation has to be able to deal at least with the black-body case, we conclude that *such a transformation does not exist*.

What *can* be said is that an observer at rest in S' , and looking at the radiation in a frequency interval $d\omega'$ coming from the solid angle interval $d\Omega'$, finds indeed a variant of (3):

$$n'_{\theta'}(\omega', T'_{\theta'}) d\omega' d\Omega' = \frac{\omega'^2/c^3}{2\pi^2(e^{\hbar\omega'/kT'_{\theta'}} - 1)} d\omega' d\Omega'. \quad (5)$$

Here $T'_{\theta'}$ is a “*directional*” temperature defined by

$$T'_{\theta'}(T, v, \theta') = \frac{T\sqrt{1 - v^2/c^2}}{1 - \cos\theta' v/c}, \quad (6)$$

and θ' is the angle between the axis of motion and the direction of observation [3]. (This result was rediscovered later in connection with studies of the 2.7 K background radiation, e.g. [30], and so has become well-known.) Thus, it should be clear that, regrettably, one cannot discover the desired transformation $T'(T, v)$ used in Eq. (3) by averaging the elegant result (5) over solid angles. One finds instead

$$\frac{1}{4\pi} \int n'_{\theta'}(\omega', T'_{\theta'}) d\omega' d\Omega' = n'(\omega', T, v) d\omega'. \quad (7)$$

This states what is at first sight a surprising result: Eq. (1) is found exactly; this time, therefore, without any appeal to quantum field theory.

Any definition based on some operational procedure, like: “*Temperature is what some pre-chosen device in S' measures*” will be arbitrary because different “thermometers”, and measuring procedures will lead to different functional dependencies. For example, if we construct a thermometer which measures the average of (6), $T'_{\text{aver1}}(T, v) \equiv \langle T'_{\theta'} \rangle$, which is perfectly possible, its reading would satisfy

$$T'_{\text{aver1}}(T, v) = \frac{1}{4\pi} \int T'_{\theta'} d\Omega' = \frac{cT}{2v\gamma} \ln \frac{1+v/c}{1-v/c}. \quad (8)$$

Another thermometer constructed to measure $T'_{\text{aver2}}(T, v) \equiv \langle T'^4_{\theta'} \rangle^{1/4}$, leads, of course, to a completely different result (p. 31 of [31])

$$T'_{\text{aver2}}(T, v) = \left[\frac{1}{4\pi} \int T'^4_{\theta'} d\Omega' \right]^{1/4} = T \left[\frac{1+v^2/3c^2}{1-v^2/c^2} \right]^{1/4}. \quad (9)$$

Clearly $T'_{\text{aver1}}(T, v=0) = T'_{\text{aver2}}(T, v=0) = T$. In fact, a good thermometer is a device which measures a temperature T [see Eq. (2)], when it is at rest with the thermal bath. Other operational procedures to measure temperature based on Unruh-DeWitt detectors are equally possible, and also give different results [26].

Previous writers were often concerned with the manipulation of Lorentz transformations of thermodynamical variables such as energy, volume, entropy, etc. This approach leads to doubtful results [see the discussion of Eq. (4)] unless the theory is made intrinsically covariant (see e.g. [15], [23] for former, and [24] for more recent formulations). In this case four-tensors S^μ , Q^μ , etc, replace some of the usual thermodynamical variables like entropy, heat, etc, and their significance then goes beyond the normal thermodynamics. (For example, by writing the entropy four-flux as $S^\mu = Su^\mu$, where u^μ is the four-velocity associated with equilibrium states, the usual entropy is given by projecting S^μ on u^μ .) It will be noted that our approach has by-passed this problem since we are discussing this issue from a microscopic rather than a macroscopic point of view, and no explicitly thermodynamical arguments are needed.

It might be thought that a rough way of seeing that no black-body equivalent temperature exists for a radiation enclosure moving relative to an observer may be by noting simply that, because of the angle dependence in (5), this equation cannot be associated with a legitimate thermal bath (which is necessarily isotropic). While this view is correct, we have in this paper gone beyond this observation by obtaining the angle-averaged Eq. (5) in the form of Eq. (1). This has the merit of linking the above simple observations with earlier works and making the case against the temperature transformation much more explicit.

In summary, our main conclusion is that because the temperature concept of a black body is unavoidably associated with the Planckian thermal spectrum, and because a bath which is thermal in an inertial frame S is non-thermal in an inertial frame S' , which moves with some velocity $v \neq 0$ with respect to S , a universal relativistic temperature transformation $T' = T'(T, v)$ cannot exist. Thus the proper temperature T alone is left as the only temperature of universal significance. This seems to complete a story started 90 years ago [1] of how the *usual* temperature transforms, and to conclude a controversy [7] of 33 years' standing.

Acknowledgements: PL would like to acknowledge Fundação de Amparo à Pesquisa do Estado de São Paulo, and the European Union through contracts ERB CH RXCT 92 0007 and CIPA-CT-92-4026, and GM would like to acknowledge Conselho Nacional de Desenvolvimento Científico e Tecnológico for partial financial support.

REFERENCES

- [1] A. Einstein, *Jahrb. der Radioaktivität und Elektronik*, **4** (1907) 411.
- [2] M. Planck, *Ann. der Physik*, **26** (1908) 1.
- [3] W. Pauli, *Theory of Relativity*, (Pergamon Press, London, 1958); *Die Relativitätstheorie*, *Encycl. d. math. Wiss.* **V(2)** (Teubner, Leipzig, 1921).
- [4] R.C. Tolman, *Relativity, Thermodynamics and Cosmology*, (Clarendon Press, Oxford, 1934).
- [5] M. Von Laue, *Die Relativitätstheorie*, 1 Band, 7th Ed. (Vieweg, Braunschweig, 1961).
See also earlier editions, e.g. 5th Ed. (1951).
- [6] H. Arzeliès, *Thermodynamique relativiste et quantique*, (Gauthier–Villars, Paris, 1968).
- [7] H. Ott, *Zeits. Phys.*, **175** (1963) 70.
- [8] H. Arzeliès, *Nuovo Cimento*, **35B** (1965) 792.
- [9] T.W.B. Kibble, *Nuovo Cimento*, **41B** (1966) 72.
- [10] F. Rohrlich, *Nuovo Cimento*, **45B** (1966) 76.
- [11] R.K. Pathria, *Proc. Phys. Soc.*, **88** (1966) 791.
- [12] P.T. Landsberg, *Nature*, **212** (1966) 571; *Nature*, **214** (1967) 903; P.T. Landsberg, and
K.A. Johns, *Proc. Roy. Soc. A*, **306** (1968) 477.
- [13] L.A. Schmid, *Nuovo Cimento*, **47B** (1967) 1.
- [14] Editorials, *Nature*, **214** (1967) 1069; *Nature*, **221** (1969) 1100; *Nature*, **224** (1969) 219.
- [15] N.G. van Kampen, *Phys. Rev.*, **173** (1968) 295.
- [16] I. Brevik, *Nuovo Cimento*, **63B** (1969) 250.
- [17] S. Nakajima, *Prog. Theor. Phys.*, **41** (1969) 1450.

- [18] N.L. Balasz; R. Balescu; P.T. Landsberg; L.A. Schmid, in: *A Critical Review of Thermodynamics*, eds. E.B. Stuart, B. Gal-or, and A.J. Brainard, (Monobook Corporation, USA, 1970).
- [19] J. Bicák, *Lett. Nuovo Cimento*, **1** (1969) 302.
- [20] R. Hakim, and A. Mangeney, *Lett. Nuovo Cimento*, **1** (1969) 429.
- [21] P.T. Landsberg, *Phys. Rev. Lett.*, **45** (1980) 149.
- [22] O. Grøn, *Nuovo Cimento*, **17B** (1973) 141.
- [23] P.T. Landsberg, *Essays in Phys.*, **2** (1970) 93;
- [24] W. Israel, *J. Non-Equilib. Thermodyn.*, **11** (1986) 295.
- [25] R. Aldrovandi, and J. Gariel, *Phys. Lett.*, **A170** (1992) 5.
- [26] S.S. Costa, and G.E.A. Matsas, *Phys. Lett.*, **A209** (1995) 155.
- [27] A. Komar, *Gen. Rel. Grav.*, **27** (1995) 1985.
- [28] Ch. Fenech, and J.P. Vigièr, *Phys. Lett.*, **A215** (1996) 246.
- [29] W.G. Unruh, *Phys. Rev. D*, **14** (1976) 870; B.S. DeWitt, in: *General Relativity*, eds. S.W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1979).
- [30] R.N. Bracewell, and E.K. Conklin, *Nature*, **219** (1968) 1343; P.J.E. Peebles, and D.T. Wilkinson, *Phys. Rev.*, **174** (1968) 2168; G.R. Henry, R.B. Feduniak, J.E. Silver, and M.A. Peterson, *Phys. Rev.*, **176** (1968) 1451.
- [31] A.P. Lightman, H.W. Press, R.H. Price, and S.A. Teukolsky, *Problem Book in Relativity and Gravitation* (Princeton University Press, Princeton, 1975)