Structural and E lectronic P roperties of Sm all N eutral (M gO)_n C lusters

- E.de la Puente¹, A. Aguado¹, A. Ayuela² and J.M. Lopez¹
 - 1. Departamento de F sica Teorica, Facultad de Ciencias, Universidad de Valladolid. 47011 Valladolid, Spain.
 - 2. Institut für Theoretische Physik, Technische Universität Dresden, 01062 Dresden, Germany.

January 29, 2022

A bstract

Ab initio Perturbed Ion (PI) calculations are reported for neutral stoichiom etric (M gO)_n (n 13) clusters. An extensive number of isom er structures was identi ed and studied. For the isom ers of (M gO)_n (n 7) clusters, a full geom etrical relaxation was considered. Correlation corrections were included for all cluster sizes using the C oulom b-H artree-Fock (C H F) m odel proposed by C lem enti. The results obtained compare favourably to the experimental data and other previous theoretical studies. Inclusion of correlation is crucial in order to achieve a good description of these systems. We not an important number of new isom ers which allows us to interpret the experimental magic numbers without the assumption of structures based on (M gO)₃ subunits. Finally, as an electronic property, the variations in the cluster ionization potential with the cluster size were studied and related to the structural isom or properties.

PACS: 36.40.+d; 61.50 Lt; 61.60.+m; 79.60 Eq Keywords: Clusters. Magnesium oxide. (MgO)_n clusters.

1 Introduction

C lusters provide a new state of aggregation from which the condensed matter properties eventually emerge, being thus of form idable interest in order to investigate the transition from the molecular world to the solid state physics. Small clusters often present substantial deviations in their physical and them ical properties when compared both to the molecule and the bulk phase. Rationalizing the evolution of those properties with the cluster size is a major challenge to the now adays science. As an example, for an intermediate size cluster, the number of dierent isomers which may coexist in a small energy range may be quite large. Though it is a very dicult task, a full characterization of those isomer congurations for each cluster size would be highly desirable for a good understanding of cluster growing processes.

In this paper we focus our interests in small stoichiom etric (M gO)_n clusters. These have received special attention in the last few years both from experimentalists and theoreticians. Saunders[1, 2] published mass-spectra and collision-induced-fragmentation (CIF) data for sputtered M gO cluster ions, and found enhanced stabilities for (M gO)_n⁺ clusters with n = 6, 9, 12 and 15. The results were interpreted there in terms of (M gO)₃ subunits from which the clusters built up. Ziem ann and Castlem an [3, 4, 5] performed experimental measurements by using laser-ionization time-of-ight mass spectrometry [6, 7]. They found \magic clusters" at n = 2, 4, 6, 9, 12 and 15. In order to explain the features observed in their mass spectra, they performed theoretical calculations by using the rigid ion and the polarizable ion shell models [8, 9, 10]. Their main conclusion was that the clusters form compact cubic structures similar to pieces of the M gO crystal lattice. Self consistent calculations on these clusters were performed by M oukouri and

Noguera [11, 12] with the use of a sem iem pirical tight binding approach. The necessity of going beyond a fully ionic model like that of pair potential interactions was there remarked. Recio et al[13, 14] reported ab initio molecular orbital calculations on neutral clusters of M gO containing up to 26 atoms. These calculations included correlation e ects in the clusters with n=1-6 molecules. Their results were in agreement with Saunders' hypothesis, in the sense that structures based on the (M gO)₃ subunit were preferred over the cubic like ones. Very recently, Malliavin et al[15] have performed calculations on (M gO)_n (n=6) clusters using the DMOL method. The geometries obtained in that work are in total agreement with those presented by Recio et al.

Each particular work from the above mentioned ones represented, in principle, an improvement over the preceding investigations. Pair potential models provided the st theoretical attempt to identify and classify di erent isom ers and calculate the energy di erences between them . Nevertheless, som e selected set of empirical parameters for M q^{q+} and O q = 1,2 ions has to be used as input, and that set is the sam e for ions which are in nonequivalent cluster sites. The studies by M oukouri and N oguera added self consistency to the calculations, but they were not free of empirical parameters. Finally, the rst ab initio study on these systems was carried out by Recio et al. Several approximations had to be done in their calculations, however, in order to maintain the computational time at reasonable values: (a) Large geom etrical distortions in their considered isom ers were not allowed, as the optim ization process was carried out just by varying the nearest neighbour distance R (for ringlike structures, the stacking distance between rings was also independently optim ized) and, for some structures, even this value was kept xed; (b) Several other isomers, which may be important in a more

com plete characterization of the di erent isom er structures adopted by sm all $(M \ gO)_n$ clusters, were not considered; (c) Correlation corrections could be included (at the $M \ P \ 2$ level) only in clusters containing up to 12 atom s.

The present work undertakes a new extensive and system atic ab initio study of (M gO)_n clusters with n up to 13. We have used the ab initio Perturbed Ion (aiPI) model[16], which is based on the Theory of Electronic Separability (TES)[17, 18] and the ab initio Model Potential (aiMP) approach of Huzinaga et al[19], within the Restricted Hartree-Fock (RHF) approximation. The model has been used by our group in several studies on (N aC l)_n [20] and (N aI)_n [21] clusters, and also in a global study of alkalichalide clusters[22]. Although we do not expect to obtain accurate results for the only partially ionic M gO molecule, our main concern is by far not the molecule, but intermediate size clusters where a full range of different isomer structures may be studied. The solid M gO has been excellently described by the aiPI model[23], and precise results are achieved for (M gO)_n clusters from n = 3 on.

Our calculations represent a major advance with respect to pair potential or sem iem pirical methods. The ion-cluster consistency achieved in the calculations results in a dierent ab initio description of each nonequivalent ion in the cluster, without requiring the use of any empirical parameter [20, 21, 22]. On the other hand, they represent an alternative description to the molecular orbital models, and complete the results obtained by Recio et al in the following aspects: (a) For (MgO)_n (n 7) clusters, a full geometrical relaxation of the dierent isomers has been considered; (b) We have studied a more complete set of isomers; (c) Correlation corrections have been included for all cluster sizes (up to 26 atoms). Finally, some electronic properties as ionization potentials have been also considered.

Results presented here are also aim ed to assist in the interpretation of possible future experimental investigations on these clusters. Renewed interest in isom er geometries has recently emerged because of the drift tube experimental studies, which, by measuring the mobility of cluster ions through an inert bu er gas under the in uence of a weak electric eld, provide valuable information about the cluster geometries [24, 25, 26]. However, we have no know ledge on experimental studies of this kind on (MgO)_n clusters.

The rest of the paper is structured in several sections. In the next one, we give a brief resum e of the aiP Im odel for the study of clusters (the interested reader can obtain more details in our previous works[20, 21, 22]), and some computational details. Section III deals with the principal structural and electronic results of the present study. Conclusions are given in the last section.

2 ComputationalMethod

The ab initio perturbed ion model[16] was originally designed for the description of ionic solids[27], and subsequently adapted to the study of clusters in our group [20, 21, 22]. Its theoretical foundation lies in the theory of electronic separability [28] for weakly overlapping groups[29], and its practical implementation in the Hartree-Fock (HF) version of the TES[17, 18]. The HF equations of the cluster are solved in localized Fock spaces, by breaking the cluster wave function into local nearly orthogonal group functions (ionic in nature in our case). When the self-consistent process (see details below) nishes, the outputs are the total cluster energy $E_{\rm clus}$ and a set of localized wave functions for each geometrically nonequivalent ion in the cluster. The

cluster energy can be written as a sum of ionic additive energies [20, 21, 22]:

$$E_{\text{chis}} = \sum_{R=1}^{X^{N}} E_{\text{add}}^{R};$$
 (1)

where the sum runs over all ions in the cluster, and the contribution of each particular ion to the total cluster energy $(E_{\rm add}^{\rm R})$ can be expressed in turn as a sum of intraionic (net) and interionic contributions:

$$E_{\text{add}}^{R} = E_{\text{net}}^{R} + \frac{1}{2} \sum_{S \in R}^{X} E_{\text{int}}^{RS} = E_{\text{net}}^{R} + \frac{1}{2} E_{\text{int}}^{R}$$
 (2)

Once E $_{\rm clus}$ has been obtained, the binding energy perm olecule of the (M gO) $_n$ cluster with respect to the dissociation process (M gO) $_n$! nM g^2+ + nO + ne + nE $_{\rm bind}$, is given by:

$$E_{bind} = \frac{1}{n} E_{clus} \quad nE_0 (M g^{2+}) \quad nE_0 (O);$$
 (3)

where E_0 (M g^{2+}) and E_0 (O) are the energies of the M g^{2+} and O free ions, respectively.

The localized nature of the aiP I procedure has some advantages over the usual molecular orbital models. As in weakly overlapping systems the correlation energy correction is almost intraionic in nature (being therefore a sum of contributions from each ion), the localized cluster-consistent ionic wave functions may be used to attain good estimations of this correction. In this paper, the correlation energy correction is obtained through C lementi's C oulom b-H artree-Fock method [30, 31]. Besides, it also allows the development of computationally e cient codes [32] which make use of the large multi-zeta basis sets of C lementi and Roetti [33] for the description of the ions. At this respect, our optimizations have been performed using basis sets (5s4p) for M g^{2+} and (6s4p) for O g^{2+} respectively. Inclusion of di use basis

functions has been checked and shown unnecessary. A general discussion on the election of the most appropriate basis set within the aiP I model has been given elsewhere [22].

A few comm ents regarding the weak-overlap assumption and the description of the 0² anions are worth of mention. 0² is an unstable anion in free space, so a basic problem in an ionic description of M gO clusters is the correct description of the 02 wave function. Luana et al have shown how the 0² anion is stabilized by the action of the crystal environm ent in solid M gO [23]. Speci cally, the 2p oxygen orbital experiences a large contraction in the lattice, which makes it stable as an embedded anion. The resulting 0^2 wave function leads to a diam agnetic susceptibility and an electron kinetic energy increase in quantitative agreem ent with the experim ent. Thus, the quality of that wave function is asserted. An analysis of the PI results reveals that the orbital contraction is mainly due to the action of the projection operators. The same conclusion has been achieved for the M gO clusters studied here. The projection operator supplies two major e ects[29]: On one hand, it tries to maintain the strong-orthogonality hypothesis, being therefore responsible for the orbital contraction. On the other hand, it has a well-de ned physical meaning, namely, its expectation value coincides exactly with the overlap energy if we assume that this overlap is \weak". By \weak" it is meant that the products of two overlap integrals $(S_{AB})_{ij}(S_{CD})_{km}$ with A \in B and C \in D are negligible $((S_{AB})_{ij} = \langle A_j j_j \rangle, where A_i is$ the ith orbital of the ion A). This corresponds to the following truncation of the Low din expansion $\beta 4$ for S^{-1} :

$$S^{-1}$$
 I T $(T = S I)$: (4)

W ithin this context, the hypothesis of weakly overlapping ions does not im -

ply that the overlap contribution to the cluster energy is negligible. What is neglected is those contributions to the overlap energy coming from second and higher orders in the Low din expansion. The overlap energies of the 0^2 anions in the (MgO)₃ clusters are just 0.5eV larger than the corresponding value in the solid (12.84 eV for 0^2 : MgO), and they are decreasing functions of the cluster size. The assumption of weakly overlapping ions is good in the solid (the results are exceptional), and we consider that such an small increase in the overlap energies does not invalid that assumption for cluster studies. Our basis set for oxygen anion is exible enough because by adding di use basis functions the cluster energy does not lower anymore, nor the projection operator is more elective in projecting the frozen orbitals out of the active space (we would like to remark that an extensive set of dilerent basis functions was considered in test calculations on (MgO)₂ and (MgO)₆ isomers. The ones selected gave the lowest cluster energies and achieved saturation in the value of the overlap energies).

We have used the following self-consistent method: for a given distribution of the ions forming the cluster, we consider one of them as the active ion R (for instance, a particular oxygen anion), and solve the Self-Consistent-Field equations for anion R in the eld of the remaining ions, which are considered frozen at this stage. The solution obtained is transferred to all the anions equivalent to anion R, that is, to the anions which have equivalent positions in the cluster. Next, we take a non equivalent oxygen anion (anion S) as the active ion and repeat the same process. Evidently, since anions S are not equivalent to anions R, the energy eigenvalues and wave functions of electrons in anions S are di erent from those of anions R. We continue this process in the same way until all the inequivalent anions have been exhausted. The same procedure is then followed for the magnesium

cations. The process just described is a PI cycle. We iterate the PI cycles until convergence in the total energy of the cluster is achieved.

As input geometries we considered cubic structures resembling pieces of bulk M gO crystal, ring structures (mainly hexagonal, but also some octagonal and decagonal), m ixed structures m ade up usually of a regular isom er with a molecule attached to it, and some more open structures like the truncated octahedron for n = 12 or the wurtzite piece for n = 7. These trial geom etries have been either taken from the geom etries considered in pair potential calculations on M gO and alkali halide clusters or intuitively guessed with the experience gathered in our previous works on alkali halide clusters. The input geom etries have been fully relaxed (that is, the total cluster energy has been minimized with respect to variations in all the (3N independent variables, where N is the number of ions) for (M gO) $_{\rm n}$ (n 7) clusters, at the HF level. A simplex downhill algorithm has been used in these calculations [35, 36]. For clusters with n = 8-13 this procedure became com putationally quite expensive and we turned to geom etrical optim izations (also at the HF level) with respect to a limited number of relevant parameters. Cuboid structures have been relaxed with respect to a single param eter, the rst-neighbour cation-anion distance, whereas in the ring structures the cation and anion distances to the center of the ring, and the distance between rings have been independently varied. For the mixed structures, the part of the regular structure which is nearest (rst and second neighbours) to the attached molecule is relaxed together with the molecule, and the rest of the isom er is kept xed at the geom etrical con guration with the minimum energy found for that isomer without the molecule attached. The CHF correlation correction is then introduced for all the studied isomers at the equilibrium geometries found at the HF level. These geometries are further scaled (without shape modi cations) until the minimum CHF energy is achieved. We do not expect, from the results obtained by Recio et al[13], that inclusion of correlation substantially modi es the isomers shape. We have carried out these scaling calculations mainly because they can a ect dierently to ringlike and cubiclike structures (and therefore to the energetic isomer ordering) and also in order to assess its in wence on the averaged cluster bond length, and its evolution towards the bulk value.

3 Results.

3.1 Lowest energy structures and isom ers.

Results concerning the structural properties of small neutral (M gO)_n clusters, obtained by following the method explained in the previous section, are shown in gure 1. Small and large spheres are used to represent M g and O ions, respectively. The most stable CHF structure (rst isomer) is shown on the left side for each cluster size. The rest of the structures are the low-lying isomers obtained in the calculation. For clusters with 3, 4 and 5 m olecules we show two dierent views of some isomers. The numbers given below each isomer are the total cluster energy dierences with respect to the ground state (denoted GS in the gure), at the CHF (rst row) and HF (second row) levels.

For small clusters sizes (n = 2-6) only two isomers have been considered. The ground state geometries in the HF calculations are mainly rings: hexagonal for n = 3, decagonal for n = 5 and a stacking of two hexagonal rings for n = 6. However for n = 4 the GS is a cube. The inclusion of correlation does not change the GS structures except for n = 5, where a distorted cube with a M gO molecule attached to it becomes energetically more stable. These re-

sults are in total agreement with the results of Recio et al[14] and Malliavin et al[15]. However, the isomers obtained in our calculations have large geometrical distortions compared to those obtained by Recio et al, mainly those which have non-ringlike congurations: the rst excited isomer for n=3 was obtained after a full relaxation of a planar piece of the MgO bulk lattice; this planar piece, with no distortions, is also the rst excited isomer in Recio et el results, but we can see in g. 1 that the nalgeometry is greatly distorted with respect to the perfect bulk structure. This kind of deformations has been also obtained in lithium halide [22] and sodium iodide [23] clusters. The above results show that the PI calculations agree with those of other more \popular" ab-initio calculations: GAUSSIAN [13, 14] and DMOL [15]. That means that the PI method is a good candidate for obtaining the ground state and other geometrical and electronic properties of clusters with ionic bonding like the MgO)_n clusters considered in this paper.

For the clusters with n=7-13 we have considered a larger number of initial geometries in our calculations. O verall, the ground state geometries obtained in the HF calculations are the same that those obtained by Recio et al. We appreciate dierences only for those cluster sizes in which the ground state structures that we obtain were not considered by Recio et al in their calculations (n=11,12,13).

The inclusion of correlation in the calculations changes the GS geom etries for n=7 and 8. These results can not be compared with the calculations performed by Recio et albecause they did not include the correlation contribution in this range of sizes; neither with the calculations of Malliavin et albecause they did not perform any calculations in this size range. For n=7 the HF ground state isomer is the wurtzite piece whereas the CHF ground state is the bulk-like $(M \ gO)_6$ piece with a M gO molecule attached

to it. For n = 8 the HFGS is a stacking of two octagonal rings whereas the CHFGS is a mixed structure formed by \pining" the most stable (M gO)6 and $(M \text{ gO})_4$ isom ers. The last structure is also the second isom er in the HF results. For n = 9 the ground state isom er is a stacking of three hexagonal rings, the same as in the HF calculations; the other isomers considered are a piece of M gO bulk and the (M gO), HFGS with a molecule attached to it. Forn = 10 the GS geometry is a piece of the M gO lattice. The other isomers considered in our calculations are, in order of increasing energy: a stacking of two decagonal rings, a piece formed by \jpining" two stackings of two hexagonal rings by a face, and a piece of bulk M gO lattice with a molecule attached to it (attaching a molecule to the (M gO)₉ GS leads to an isomer which lies higher in energy). For n = 11 the GS isomer is a mixed structure com posed by a stacking of two hexagonal rings plus a bulk-like part, and the other isomers considered are either from mixed type or bulk-like. A truncated octahedron is obtained as the GS for n = 12. This isomer has been also found to be specially stable in $(N aI)_{12}$ clusters [21]. It was not considered in ref.[14]. Finally, a mixed structure composed by a piece of the M gO lattice and an octagon is found as the $(M gO)_{13}$ ground state. Our second isomer is the defect-cuboid structure, which was the single isomer tried in ref.[14]. A point which is worth of mention from these two nalcluster sizes is that there is the possibility to obtain dierent isom ers with the same geometrical shape but with cations and anions interchanged. The electrostatic interaction tends to stabilize energetically anions [20, 21], so those isomers with a central oxygen are more stable.

We compare now the preceding results to the results obtained with the PI model for alkali-halide (AX)_n clusters[20, 21, 22]. In our previous works we found evidence of a clear-cut separation between the structural properties of

di erent alkali-halide clusters. (K X) $_{\rm n}$ and (R bX) $_{\rm n}$ clusters showed almost from the very beginning a preference to adopt minimum energy structures which are fragments of the bulk lattice. However, (LiX)n clusters adopted preferentially more open structures. (NaX)_n showed an intermediate behaviour. Our structural results for (M gO), clusters show, in outline, large sim ilarities with those of (LiX)_n or (N aI)_n clusters. These are the least ionic compounds in the alkali-halide series. Rounded geometries had been proposed by Twu et al[7] in those cases showing a decreased ionic bonding character. Then, our minimum energy isomers are consistent with a reduced ionic character in (M gO), clusters. This non-fully ionic character was also pointed out in the analysis of Recio et al[14]. The discussion there was based on a comparison with the classical results of Ziem ann et al[3], who performed pair potential calculations by using two dierent (but both fully ionic) models, with dierent nominal charges assigned to the ions. The calculations for $(M g^{+1} O^{-1})_n$ clusters revealed mainly cubic structures, whereas in $(M g^{+2}O^{-2})_n$ calculations a set of much more open structures was obtained. As the $(M g^{+1} O^{-1})_n$ calculations explained better the experimentalmass spectra, Ziemann et alconcluded that pieces of M gO bulk lattice are obtained for small $(M \neq 0)_n$ clusters. Both our calculations and those of refs.[14, 15] obtain results which are in between these two lim it cases, so nom inal charges between jljand jljshould be ascribed to M g=0 ions in classical pair potential simulations. However, our calculations show that a description in terms of weakly overlapping M g²⁺ and O² groups is completely appropriate provided that all the relevant quantum -m echanical interactions (overlap included) are accounted for properly and a full ion-cluster consistency is achieved. It is worth of stressing that such a description does not enter in con ict with a possible assignment of fractional charges to each ion.

A population analysis is best carried out directly in term softhe total density, following the Bader's scheme for deriving atom ic properties from the topology of the charge density [37]. That has been recently done in PI calculations on solids [38], and fractional ionic charges have been derived for a variety of ionic crystals.

From the study of small (M gO)_n (n = 2-13) clusters we conclude that, in the earliest stages of cluster growing, the bulk lattice fragments are not the GS isomers except for some particular values of n. For the rest of cluster sizes other (m ore open) structures are preferred. Thus, the convergence to the bulk structure is not reached yet in this size range (the solid M gO crystallizes in the f.c. structure [39]).

We can appreciate in g.1 for n=6, 9 and 12 (and also for other cluster sizes), that the inclusion of correlation enhances the stability of the hexagonal isom ers over the cubic ones. Though in the size range covered open structures are preferred over the cubic ones, eventually there will be a cluster size where the cubic structures become energetically more stable, and this special value of n will depend on the inclusion or not of correlation in the calculations. The above observation suggests that this \critical n will be larger upon inclusion of correlation, that is, pieces of the MgO bulk lattice will appear as GS isomers for values of n which are larger in CHF calculations than in HF calculations.

The magic numbers for small (M gO) $_n^+$ clusters emerging from mass spectra experiments have been explained by Saunders[1] and by Recio et al[14] as resulting from specially enhanced structures based on the stacking of (M gO) $_3$ subunits. Our results show that, at least for neutral (M gO) $_n$ clusters (and also for ionized (M gO) $_n^+$ clusters if the vertical approximation holds[14]), this conclusion is not universal. The truncated octahedron obtained as the ground

state isom er for n = 12 in our calculations indicates that this interpretation should be taken with some care.

3.2 Inter-ionic distances and binding energies.

In qure 2 we present the average interionic CHF distance d between M q and 0 nearest neighbours for the ground state geom etry as a function of the cluster size n. d presents an irregular behaviour as a function of n, due to the fact that the ground state geom etries for the di erent values of n do not correspond to the sam e type of structural fam ily. The points that correspond to the same structural family have been pined by the two dashed lines, and it can be appreciated that the variation of d is smoother for each structural fam ily. The value of dincreases with the number of ions in the cluster. The cubic clusters have an interatom ic distance larger than the hexagonal ones. The isomers which resemble pieces of the bulk M gO lattice are the important set in order to compare with the bulk value. In the M gO crystald = 2.106 A [39], and we obtain a value of d = 1.999 A for the GS isom erwith n = 10. The saturation value is not achieved yet, but the trend is the correct one. The contraction of the distances induced by the inclusion of correlation is m ore important in cubiclike pieces than in ringlike ones: the average e ect in cubic structures is a contraction of approximately 4%, while in hexagonal structures the contraction is only of som e 2 %. The inclusion of correlation is necessary in order to obtain quantitative agreem ent with the experim ental bulk value in the lim it of n large.

In Figure 3, we represent E_{bind} as a function of n, both for the HF and CHF calculations. The general trend is an increase of the binding energy with n in both cases. Superposed to that general trend, especially stable clusters are predicted for certain values of n (\magnitude agic numbers"); these are identified

in the gures as maxima or pronounced changes in the slope. We observe that both type of calculations do not predict the same set of magic numbers. The HF curve is quite smooth for small values of n, and maxima are observed only at n=9, 12 (pronounced slope changes are not observed). The inclusion of correlation elects (CHF curve) is crucial to achieve agreement with the experimental results [1, 3], which predict magic clusters at n=2, 4, 6, 9, 12, 15... The maxima at n=6, 9, 12 are clear, and a pronounced slope change is appreciable at n=4. The abundance maximum obtained experimentally at n=2 should be probably related to the singly-ionized stoichiometric clusters and not to the neutrals [14].

3.3 Cluster ionization potentials.

In this subsection we analyse the variation of the cluster ionization potential (\mathbb{P}) with the cluster size. The output of the PI calculation contains a fully cluster-consistent wave function for each inequivalent ion in the cluster. In our model, the electron must be extracted from a special localized anionic orbital when the cluster is ionized. The vertical \mathbb{P} of the cluster is calculated as the smallest binding energy of a 2p anionic electron (K copmans' approximation). The results are given in gure 4, where we have plotted the eigenvalues of the 2p orbitals of 0^2 anions as a function of p for the most stable structure of each cluster size in the HF (p.4a) and in the CHF (p.4b) calculations. We appreciate a band of eigenvalues for each cluster size because the anions can occupy inequivalent positions in the cluster. These band gaps are indicative of the different reactivity of each nonequivalent cluster site, they can determ ine the preferred adsorption sites, and could be helpful in photoionization spectroscopy studies.

The dashed lines, which join the lowest 2p energy eigenvalues for each

cluster size, indicate the variation of the HF (4a) and CHF (4b) ionization potentials with the cluster size. The variation of the vertical cluster ionization potential with n is related to the relative stabilities of neutral and singlyionized stoichiom etric clusters [14]: When there is a maximum in the IP for a cluster size n, the ionized structure is less stable than the neutral one against loss of a M gO m onom er; the opposite occurs at the m inim a in the IP curve. For the HF results, a set of minima at n = 4, 7, and 11 and maxima at n = 3, 5, 8, and 10 is observed, in agreement with the results of Recio et al. The di erence is that they obtain a maximum at n = 12; but the results are not directly comparable for these cluster sizes (n = 12, 13), because the isom ers that we obtain as the GS were not considered in their calculations. This pattern changes upon inclusion of correlation e ects (see CHF curve): m in im a are found at sizes n = 5, 7, and 11, and maxim a at n = 3, 6, and 10. These changes are directly related to the change in the GS isom er for n =5. We can give a clear geom etrical interpretation to the minima in the light of the localized picture provided by the PIm odel. At the minim a (n = 5, 1)7, and 11), m ixed structures were found as the most stable isomers (q.1). The least bound anionic p-electron in those clusters corresponds always to that oxygen with the smallest number of neighbouring ions. That specic oxygen is less stabilized than the other oxygens in the cluster because of the reduced electrostatic interaction for those cluster sites. An electron will be rem oved from it more easily than for the rest of cluster sizes (for which the GS isomer is more \compact"). Such a clear geometrical interpretation of the maxima in the IP's is not apparent for the small size range covered in this work: the opposite situation (maxima in the IP for those ground states with the highest sym m etries) is not always observed. As the main contribution to the anionic binding energy is the M adelung potential, and this is the largest

for the bulklike structures, we expect maxima to appear precisely in compact cubic structures for larger cluster sizes.

In simple metallic clusters, the variations in the cluster ionization potential are correlated with the magic numbers, indicating that the magic numbers are electronic in nature [40]. There is not such a denite connection in ionic clusters, and the variations observed in the cluster ionization potential are related mainly to structural features. From our discussion, the existence of such a connection between electronic (IP's) and geometrical properties in (MgO)_n clusters becomes clear. The question of the interconnection between electronic, energetic, and geometrical cluster properties has been more thoroughly addressed in ionic alkali-halide materials [22].

4 Conclusions

Ab initio Perturbed Ion (PI) calculations have been carried out in order to investigate the structural, energetic and some electronic properties of small neutral stoichiom etric (M gO)_n clusters. The set of studied isomers has been more complete than in previous calculations, and some of the not previously considered isomers have resulted to be either the ground state structure or a low-lying isomer for several cluster sizes. Specifically, we have shown that the interpretation of the magic numbers of small (M gO)_n clusters in terms of stackings of (M gO)₃ subunits is not universal, because a truncated octahedron is more stable for n = 12 than the corresponding hexagonal isomer. General structural distortions have been considered for (M gO)_n (n 7) clusters, and these deformations have been shown to be really in portant in those isomers with the lowest symmetries. Correlation corrections have been included for all cluster sizes with the C lementi's C oulomb-Hartree-Fock (C HF)

m ethod. Their inclusion has been very important in the determination of some ground state isomers and of the energy dierences between isomers for each cluster size. Specifically, at n=5, 7, 8, and 10 dierent ground state structures are obtained after accounting for correlation elects. Comparison with previous ab initio calculations has been made and the agreement has been good whenever the comparisons have been possible.

The evolution of the average inter-ionic distance with the cluster size has been studied. Clusters from the same structural family show a smooth variation of their inter-ionic distances with n. The correlation-induced distance contractions are the largest for those isomers which are fragments of the M gO bulk lattice. Energetically, however, the correlation correction stabilizes m ore the hexagonal isom ers, so the transition towards bulk structures will take place at larger cluster sizes than HF calculations would predict. This result contrasts with the situation encountered in alkali halide clusters, where the inclusion of correlation induces a larger energetical stabilization of the cuboid isom ers [22, 41]. The (M gO), m agic num bers have been identi ed as maxima or pronounced slope changes in the binding energy versus cluster size curve. They are n = 4, 6, 9, and 12, in agreem ent with the experim ental results n = 2, 4, 6, 9, 12, 15, ... (n = 2 has been considered to be due to a larger enhancement of the singly-ionized (M gO) $_2^+$ species). Inclusion of correlation has also shown to be completely necessary in order to reproduce the correct (experim ental) magic numbers.

The interconnection between electronic and structural cluster properties has been studied by considering the variations in the cluster ionization potential (\mathbb{P}) with the cluster size n. For the minima in the \mathbb{P} , which occur for the lowest symmetry ground state isomers at n=5, 7, and 11, a clear geometrical interpretation can be given: in those isomers, there is always

an oxygen anion with a number of neighbouring ions notably reduced, from which is relatively easy (compared to the rest of cluster sizes) to remove an electron and create a singly-ionized isomer. Such a denite connection has not been found for the maxima in the size range considered in this work. The corresponding opposite situation of maxima in the IP emerging for all those ground state isomers with the highest symmetries is not observed. The increased classical electrostatic (Madelung) energy of isomers resembling pieces of the MgO crystal explains the maximum observed at n=10. Then, maxima for larger cluster sizes should be expected to appear whenever the ground state becomes a compact fragment of the MgO bulk lattice.

ACKNOW LEDGEMENTS. This work has been supported by DGI-CYT (Grant PB95-D720-C02-01). One of us (A. Aguado) acknowledges a predoctoral Grant from Junta de Castilla y Leon.

R eferences

- [1] W .A. Saunders, Phys. Rev. B 37, 6583 (1988).
- [2] W .A. Saunders, Z. Phys. D 12, 601 (1989).
- [3] P.J. Ziem ann and A.W. Castlem an, Jr., Z. Phys. D 20, 97 (1991).
- [4] P.J.Ziem ann and A.W. Castlem an, Jr., J.Chem. Phys. 94, 718 (1991).

- [5] P.J. Ziem ann and A.W. Castlem an, Jr., Phys. Rev. B 44, 6488 (1991).
- [6] C.W.S.Conover, Y.A.Yang and L.A.Bloom eld, Phys. Rev. B 38, 3517 (1988).
- [7] Y.T.Twu, C.W.S.Conover, Y.A.Yang and L.Bloom eld, Phys. Rev. B 42,5306 (1990).
- [8] T.P.Martin, Phys. Rep. 95, 168 (1983).
- [9] J.D iefenbach and T.P.M artin, Surf. Sci. 156, 234 (1984).
- [10] N.G.Phillips, C.W.S.Conover and L.A.Bloom eld, J.Chem.Phys. 94,7498 (1991).
- [11] S.M oukouri and C.Noquera, Z.Phys.D 24, 71 (1992).
- [12] S.M oukouri and C.Noquera, Z.Phys.D 27, 79 (1993).
- [13] J.M. Recio and R. Pandey, Phys. Rev. A 47, 2075 (1993).
- [14] J.M. Recio, R. Pandey, A. Ayuela and A. B. Kunz, J. Chem. Phys. 98, 4783 (1993).
- [15] M. J. Malliavin and C. Coudray, J. Chem. Phys. 106, 2323 (1997).
- [16] V. Luana and L. Pueyo, Phys. Rev. B 41, 3800 (1990).
- [17] S. Huzinaga and A. P. Cantu, J. Chem. Phys. 55, 5543 (1971).
- [18] V. Luana. Thesis Dissertation, Universidad de Oviedo (1987).
- [19] S. Huzinaga, L. Seijo, Z. Barandiaran and M. Klobukowski, J. Chem. Phys. 86, 2132 (1987).

- [20] A. Ayuela, JM. Lopez, JA. Alonso and V. Luana, [a] Z. Phys. D 26, S213 (1993); [b] An. Fis. (Spain), 90, 190 (1994); [c] Physica B 212, 329 (1996).
- [21] A. Aguado, A. Ayuela, JM. Lopez and JA. Alonso, J. Phys. Chem., accepted for publication.
- [22] A. Aguado, A. Ayuela, JM. Lopez and JA. Alonso, Preprint
- [23] V. Luana, J. M. Recio and L. Pueyo, Phys. Rev. B 42, 1791 (1990).
- [24] G. von Helden, M. T. Hsu, P. R. Kemper and M. T. Bowers, J. Chem. Phys. 95, 3835 (1991).
- [25] M.F. Jarrold, J. Phys. Chem. 99, 11 (1995).
- [26] M. Maier-Borst, P. Loer, J. Petry and D. Kreisle, Z. Phys. D (in press).
- [27] V. Luana, M. Florez, E. Francisco, A. Mart n Pendas, J. M. Recio, M. Bermejo and L. Pueyo, in Cluster Models for Surface and Bulk Phenomena, edited by G. Pacchioni, P. S. Bagus and F. Parmigiani (Plenum, New York, 1992), p. 605
- [28] R. McW eeny, Methods of molecular quantum mechanics, Academic Press. London (1994).
- [29] E. Francisco, A. Mart n Pendas and W. H. Adams, J. Chem. Phys. 97, 6504 (1992).
- [30] E.C. lem enti, IBM J. Res. Develop., 9, 2 (1965).
- [31] S.J.C. hakravorty and E.C. lem enti, Phys. Rev. A, 39, 2290 (1989).

- [32] V. Luana, A. Mart n Pendas, J. M. Recio and E. Francisco, Comput. Phys. Commun. 77, 107 (1993)
- [33] E.C. Lem enti and C. Roetti, At. Data Tables, 14, 177 (1974).
- [34] P.O.Lowdin, Adv. Phys., 5, 1 (1956).
- [35] JA. Nelder and R. Mead, Comput. J. 7, 308 (1965).
- [36] William H.Press and Saul A. Teukolsky, Computers in Physics, 5, 426, (1991).
- [37] R.F.W. Bader, Atoms in Molecules (Oxford University Press, Oxford, 1990).
- [38] A.Mart n Pendas, A.Costales and V.Luana, Phys. Rev. B, 55, 4275 (1997).
- [39] N. M. Ashcroft and N. D. M. erm in, Solid State Physics, Holt, Rinehart and W. inston, New York (1976).
- [40] W.A.de Heer, W.D.Knight, M.Y.Chou and M.L.Cohen, Solid State Physics 40, 93 (1987).
- [41] C.Ochsenfeld and R.Ahlrichs, Ber. Bunsenges Phys. Chem. 98, 34 (1994).

Captions of gures

Figure 1. Isom er geom etries for (M gO)_n clusters. The most stable CHF structure (rst isom er) is shown on the left side. Total energy di erences (in eV) with respect to the most stable structure are given for each isom er (rst row: CHF; second row: HF). M g^{2+} , small spheres; O², large spheres. For n = 3 5, two di erent views of som e isom ers are provided.

Figure 2. A veraged nearest neighbour M g-O distance for the CHF ground state structures of Figure 1. The two lines join cubic and hexagonal clusters.

Figure 3. Binding energy permolecule as a function of the cluster size.

Figure 4.0 rbital energies (with opposite sign) of the 2p levels of 0 2 anions as a function of the cluster size for the ground state structures of (M gO) $_{\rm n}$ clusters. The dashed line joins the vertical cluster ionization potentials. (a) HF results; (b) CHF results.













