

An Alternative Basis for the Wigner-Racah Algebra of the Group $SU(2)$

M. KIBLER¹ and M. DAOUD²

¹Institut de Physique Nucléaire de Lyon
IN2P3-CNRS et Université Claude Bernard
43 bd du 11 novembre 1918, 69622 Villeurbanne Cedex, France

²Laboratoire de Physique Théorique
Université Mohammed V
Avenue Ibn Batouta, B.P. 1014, Rabat, Morocco

Abstract

The Lie algebra of the classical group $SU(2)$ is constructed from two quon algebras for which the deformation parameter is a common root of unity. This construction leads to (i) a (not very well-known) polar decomposition of the generators J_- and J_+ of the $SU(2)$ Lie algebra and to (ii) an alternative to the $\{J^2, J_3\}$ quantization scheme, viz., the $\{J^2, U_r\}$ quantization scheme. The key ideas for developing the Wigner-Racah algebra of the group $SU(2)$ in the $\{J^2, U_r\}$ scheme are given. In particular, some properties of the coupling and recoupling coefficients as well as the Wigner-Eckart theorem in the $\{J^2, U_r\}$ scheme are briefly discussed.

Paper to appear in the Proceedings of the International Conference **Quantum Groups, Deformations and Contractions** (Istanbul, Turkey, 17 - 24 September 1997). The Proceedings of the Conference will be published in the Turkish Journal of Physics.

1 Motivations and Introduction

In recent years, intermediate statistics and deformed statistics were the object of considerable interest [1-19]. The use of deformed oscillator algebras proved to be useful in parastatistics, anyonic statistics and deformed statistics. In particular, one- and two-parameter deformations of the Bose-Einstein statistics (more precisely, deformations of the relevant second quantization formalism) were studied by several authors [6-19]. A common characteristic of most of these studies is that it is possible to obtain a Bose-Einstein condensation of a free gas of bosons in $D = 2$ and 3 dimensions. However, in $D = 3$ dimensions, the q -deformed Bose-Einstein (B-E) temperature is generally greater than the classical (corresponding to $q = 1$) B-E temperature. In the specific case of ^4He super-fluid in phase II, the usual q -deformations, i.e., the *à la* Biedenharn [20] and *à la* Macfarlane [21] q -deformations, yield the following inequality :

$$(T_{\text{B-E}})_{q \neq 1} > (T_{\text{B-E}})_{q=1} > (T_{\text{B-E}})_{\text{exp}}$$

so that we do not gain anything when passing from $q = 1$ to $q \neq 1$. On the other hand, by using a *à la* Rideau [22,23] deformation, it is feasible to lower the critical temperature $(T_{\text{B-E}})_{q \neq 1}$ due to the occurrence of a second parameter ν'_0 in addition to the deformation parameter q . This result corresponds to the model M_1 introduced in ref.[19]. For this model, we can obtain couples (ν'_0, q) for which $(T_{\text{B-E}})_{q \neq 1}$ is in agreement with the experimental value $(T_{\text{B-E}})_{\text{exp}} \sim 2.17$ K. However, as a drawback, the model M_1 depends on two parameters. Although it is possible to find a physical interpretation (in terms of the chemical potential) of the deformation parameter q , there is up to now no satisfying interpretation of the phenomenological parameter ν'_0 .

The just mentioned difficulty to interpret the parameter ν'_0 was the starting point of an investigation of alternative deformations of the second quantization formalism. More specifically, we investigated the *à la* Arik and Coon [24] deformation but in the case where q is a root of unity. (In the original work by Arik and Coon, the deformation parameter q is a real number : The reality of q ensures that the creation and annihilation operators are connected via Hermitean conjugation.) Thus, we arrived at the conclusion that it is necessary to simultaneously consider two quon algebras A_q and $A_{\bar{q}}$ in order to obtain a convenient framework for obtaining B-E condensation of quons.

As a first by-product, we were naturally left to the definition and study of operators, referred to as k -fermion operators, that interpolate between boson and fermion operators. These new operators arise through the consideration of two non-commuting quon algebras A_q and $A_{\bar{q}}$ for which $q = \exp(2\pi i/k)$ with $k \in \mathbf{N} \setminus \{0, 1\}$. The case $k = 2$ corresponds to fermions and the limiting case $k \rightarrow \infty$ to bosons. Generalized coherent states (connected to k -fermionic states) and super-coherent states (involving a k -fermionic sector and a purely bosonic sector) were examined. In addition, the operators in the k -fermionic algebra were used to find realizations

of the Dirac quantum phase operator and of the W_∞ Fairlie-Fletcher-Zachos algebra [25]. All these matters were discussed in Bregenz (at the Symposium *Symmetries in Science X*), Dubna (at the VIII International Conference on *Symmetry Methods in Physics*) and Istanbul (at the International Workshop *Quantum Groups, Deformations and Contractions*) and shall be reported elsewhere [26,27].

In the present paper, we would like to deal with a second by-product of our quon approach. Here, instead of considering two non-commuting quon algebras A_q and $A_{\bar{q}}$, we shall consider two realizations of two commuting quon algebras corresponding to the same root of unity $q = \exp(2\pi i/k)$ with $k \in \mathbf{N} \setminus \{0, 1\}$. We shall see how to construct (in Section 2) the Lie algebra of $SU(2)$ from these two quon algebras ; how to obtain (in Section 3) an alternative to the $\{J^2, J_z\}$ scheme of $SU(2)$; and how to develop (in Section 4) the Wigner-Racah algebra of $SU(2)$ in this new scheme. In a last section (Section 5), we shall indicate some perspectives and briefly discuss some open problems.

2 A Quon Approach to $SU(2)$

We start with two commuting quon algebras $A_i = \{a_{i-}, a_{i+}, N_i\}$, with $i = 1$ and 2 , for which the generators satisfy

$$a_{i-}a_{i+} - qa_{i+}a_{i-} = 1, \quad [N_i, a_{i\pm}] = \pm a_{i\pm} \quad (1)$$

where the deformation parameter

$$q = \exp\left(\frac{2\pi i}{k}\right) \quad \text{with} \quad k \in \mathbf{N} \setminus \{0, 1\} \quad (2)$$

(the same for A_1 and A_2) is a root of unity. As constraint relations, compatible with (1) and (2), we take the nilpotency conditions

$$(a_{i+})^k = (a_{i-})^k = 0 \quad \text{with} \quad k \in \mathbf{N} \setminus \{0, 1\} \quad (3)$$

Grassmannian realizations of eqs.(1) and (3) are obtainable from ref.[26]. In this work, we take the representations of A_1 and A_2 defined by

$$a_{1+}|n_1\rangle = |n_1 + 1\rangle, \quad a_{1+}|k-1\rangle = 0$$

$$a_{1-}|n_1\rangle = [n_1]_q |n_1 - 1\rangle, \quad a_{1-}|0\rangle = 0$$

$$a_{2+}|n_2\rangle = [n_2 + 1]_q |n_2 + 1\rangle, \quad a_{2+}|k-1\rangle = 0$$

$$a_{2-}|n_2\rangle = |n_2 - 1\rangle, \quad a_{2-}|0\rangle = 0$$

$$N_1|n_1\rangle = n_1|n_1\rangle, \quad N_2|n_2\rangle = n_2|n_2\rangle$$

on a Fock space $\mathcal{F} = \{|n_1 n_2\rangle = |n_1\rangle \otimes |n_2\rangle : n_1, n_2 = 0, 1, \dots, k-1\}$ of finite dimension ($\dim \mathcal{F} = k^2$). We use here the notation

$$[x]_q = \frac{1 - q^x}{1 - q} \quad \text{for } x \in \mathbf{R}$$

so that $[n]_q = 1 + q + \dots + q^{n-1}$ for $n \in \mathbf{N}^*$.

We now define the two following linear operators

$$H = \sqrt{N_1(N_2 + 1)}$$

and

$$U_r = \left[a_{1+} + \exp\left(i\frac{\phi_r}{2}\right) \frac{(a_{1-})^{k-1}}{[k-1]_q!} \right] \left[a_{2-} + \exp\left(i\frac{\phi_r}{2}\right) \frac{(a_{2+})^{k-1}}{[k-1]_q!} \right]$$

where the real parameter ϕ_r is taken in the form

$$\phi_r = \pi(k-1)r \quad \text{with } r \in \mathbf{R}$$

and the q -deformed factorial is defined by

$$[n]_q! = [1]_q [2]_q \cdots [n]_q \quad \text{for } n \in \mathbf{N}^* \quad \text{and} \quad [0]_q! = 1$$

The action of U_r on \mathcal{F} is easily found to satisfy

$$U_r|n_1 n_2\rangle = |n_1 + 1, n_2 - 1\rangle \quad \text{for } n_1 \neq k-1 \quad \text{and} \quad n_2 \neq 0 \quad (4)$$

and

$$U_r|k-1, 0\rangle = \exp(i\phi_r)|0, k-1\rangle \quad (5)$$

while for H we have

$$H|n_1 n_2\rangle = \sqrt{n_1(n_2 + 1)}|n_1 n_2\rangle \quad (6)$$

By using the Schwinger trick

$$j = \frac{1}{2}(n_1 + n_2), \quad m = \frac{1}{2}(n_1 - n_2) \quad \Rightarrow \quad |n_1 n_2\rangle = |j + m, j - m\rangle \equiv |jm\rangle$$

we can rewrite eqs.(4) and (5) as

$$U_r|jm\rangle = [1 - \delta(m, j)]|j, m+1\rangle + \delta(m, j)\exp(i\phi_r)|j, -j\rangle$$

Similarly, eq.(6) can be rewritten

$$H|jm\rangle = \sqrt{(j+m)(j-m+1)}|jm\rangle$$

Furthermore, we have

$$U_r^\dagger |jm\rangle = [1 - \delta(m, -j)] |j, m - 1\rangle + \delta(m, -j) \exp(-i\phi_r) |jj\rangle$$

where U_r^\dagger stands for the adjoint of U_r . For a fixed value of k , we take

$$2j = k - 1 \quad \text{with} \quad k \in \mathbf{N} \setminus \{0, 1\}$$

We can thus have $j = \frac{1}{2}, 1, \frac{3}{2}, \dots$. The case $j = 0$ corresponds to the limiting situation where $k \rightarrow \infty$.

It is obvious that the operator H is Hermitean and the operator U_r is unitary. The action of U_r on \mathcal{F} is cyclic. As a further property of U_r , we have

$$(U_r)^{2j+1} = \exp(i\phi_r)$$

that reflects the cyclical character of U_r .

Let us introduce the three operators

$$J_+ = HU_r, \quad J_- = U_r^\dagger H \quad (7)$$

and

$$J_3 = \frac{1}{2} (N_1 - N_2) \quad (8)$$

It is immediate to check that the action on the state $|jm\rangle$ of the operators defined by eqs.(7) and (8) is given by

$$J_\pm |jm\rangle = \sqrt{(j \mp m)(j \pm m + 1)} |j, m \pm 1\rangle$$

and

$$J_3 |jm\rangle = m |jm\rangle$$

Consequently, we have the commutation relations

$$[J_3, J_\pm] = \pm J_\pm, \quad [J_+, J_-] = 2J_3$$

which correspond to the Lie algebra of the group $SU(2)$. As a result, the non-deformed Lie algebra $\mathfrak{su}(2)$ is obtained from two q -deformed oscillator algebras.

To close this section, it is interesting to note that we can generate the infinite dimensional Lie algebra W_∞ from the generators of A_1 and A_2 . Indeed, by putting

$$U = U_r, \quad V = q^{N_1 - N_2}$$

and

$$T_{(m_1, m_2)} = q^{m_1 m_2} U^{m_1} V^{m_2}$$

we can prove that

$$[T_m, T_n] = -2i \sin\left(\frac{2\pi}{k} m \times n\right) T_{m+n} \quad (9)$$

where we use the abbreviations

$$m = (m_1, m_2), \quad n = (n_1, n_2)$$

and

$$m + n = (m_1 + n_1, m_2 + n_2), \quad m \times n = m_1 n_2 - m_2 n_1$$

Equation (9) shows that the operators T_ℓ span the algebra W_∞ introduced by Fairlie, Fletcher and Zachos [25]. This result parallels a similar result obtained in ref.[26] in the study of k -fermions and of the Dirac quantum phase operator.

3 A New Basis for SU(2)

At this stage, it is important to establish a link with the work by Lévy-Leblond [28]. The decomposition (7), in terms of H and U_r , coincides with the polar decomposition, described in ref.[28], of the shift operators J_+ and J_- of the Lie algebra $\mathfrak{su}(2)$. This is easily seen by taking the matrix elements of U_r and H and by comparing these elements to the ones of the operators Υ and J_T in [28]. This yields $H \equiv J_T$; furthermore, by identifying the arbitrary phase φ of [28] to $\phi_r = 2\pi jr = \pi(k-1)r$, we obtain that U_r turns out to be identical to the operator Υ of [28]. Equation (7) constitutes an important original result of ref.[28].

It is easy to prove that the Casimir operator

$$J^2 = \frac{1}{2}(J_+ J_- + J_- J_+) + J_3^2 = H^2 + J_3^2 - J_3$$

commutes with U_r for any value of r . (Note that the commutator $[U_r, U_s]$ is different from zero for $r \neq s$.) Therefore, for fixed r , the commuting set $\{J^2, U_r\}$ provides us with an alternative to the familiar commuting set $\{J^2, J_3\}$ of angular momentum theory. The (complete) set of commuting operators $\{J^2, U_r\}$ can be easily diagonalized. This leads to the following result.

Result : The spectra of the operators U_r and J^2 are given by

$$U_r |j\alpha; r\rangle = q^{-\alpha} |j\alpha; r\rangle, \quad J^2 |j\alpha; r\rangle = j(j+1) |j\alpha; r\rangle \quad (10)$$

where

$$|j\alpha; r\rangle = \frac{1}{\sqrt{2j+1}} \sum_{m=-j}^j q^{\alpha m} |jm\rangle \quad (11)$$

with the range of values

$$\alpha = -jr, -jr+1, \dots, -jr+2j, \quad 2j \in \mathbf{N}$$

The parameter q in eqs.(10) and (11) is

$$q = \exp\left(\frac{2\pi i}{2j+1}\right) \quad (12)$$

(cf. eq.(2) with $k = 2j + 1$ for $k \in \mathbf{N} \setminus \{0, 1\}$ and $k \rightarrow \infty$ for $j = 0$).

It is important to note that in eqs.(10) and (11) the label α goes, by step of 1, from $-jr$ to $-jr + 2j$. (It is only for $r = 1$ that α goes, by step of 1, from $-j$ to j .) The inter-basis expansion coefficients

$$\langle jm|j\alpha; r\rangle = \frac{1}{\sqrt{2j+1}} q^{\alpha m}$$

(with $m = -j, -j+1, \dots, j$ and $\alpha = -jr, -jr+1, \dots, -jr+2j$) in eq.(11) define a unitary transformation that allows to pass from the well-known orthonormal standard basis $\{|jm\rangle : 2j \in \mathbf{N}, m = -j, -j+1, \dots, j\}$ of the space \mathcal{F} to the orthonormal non-standard basis $B_r = \{|j\alpha; r\rangle : 2j \in \mathbf{N}, \alpha = -jr, -jr+1, \dots, -jr+2j\}$. Then, the expansion

$$|jm\rangle = \frac{1}{\sqrt{2j+1}} \sum_{\alpha=-jr}^{-jr+2j} q^{-\alpha m} |j\alpha; r\rangle$$

with

$$m = -j, -j+1, \dots, j, \quad 2j \in \mathbf{N}$$

is the inverse of eq.(11).

We thus foresee that it is possible to develop the Wigner-Racah algebra (WRa) of the group $SU(2)$ in the $\{J^2, U_r\}$ scheme. This furnishes an alternative to the WRa of $SU(2)$ in the $SU(2) \supset U(1)$ basis corresponding to the $\{J^2, J_3\}$ scheme.

4 A New Approach to the Wigner-Racah Algebra of $SU(2)$

In this section, we give the basic ingredients for the WRa of $SU(2)$ in the $\{J^2, U_r\}$ scheme. The Clebsch-Gordan coefficients (CGc's) adapted to the $\{J^2, U_r\}$ scheme are defined from the $SU(2) \supset U(1)$ CGc's adapted to the $\{J^2, J_3\}$ scheme. The adaptation to the $\{J^2, U_r\}$ scheme afforded by eq.(11) is transferred to $SU(2)$ irreducible tensor operators. This yields the Wigner-Eckart theorem in the $\{J^2, U_r\}$ scheme.

4.1 Coupling and Recoupling Coefficients in the $\{J^2, U_r\}$ Scheme

The CGC's or coupling coefficients $(j_1 j_2 \alpha_1 \alpha_2 | j \alpha; r)$ in the $\{J^2, U_r\}$ scheme are simple linear combinations of the $SU(2) \supset U(1)$ CGC's. In fact, we have

$$(j_1 j_2 \alpha_1 \alpha_2 | j \alpha; r) = \frac{1}{\sqrt{(2j_1 + 1)(2j_2 + 1)(2j + 1)}} \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} \sum_{m=-j}^j \\ \times q^{\alpha m} q_1^{-\alpha_1 m_1} q_2^{-\alpha_2 m_2} (j_1 j_2 m_1 m_2 | j m)$$

where q , q_1 and q_2 are given by eq.(12) in terms of j , j_1 and j_2 , respectively. The symmetry properties of the coupling coefficients $(j_1 j_2 \alpha_1 \alpha_2 | j \alpha; r)$ cannot be expressed in a simple way (except the symmetry under the interchange $j_1 \alpha_1 \leftrightarrow j_2 \alpha_2$). Let us introduce the f_r symbol via

$$f_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix} = (-1)^{2j_3} \frac{1}{\sqrt{2j_1 + 1}} (j_2 j_3 \alpha_2 \alpha_3 | j_1 \alpha_1; r)^* \quad (13)$$

where the star indicates the complex conjugation. Its value is multiplied by the factor $(-1)^{j_1 + j_2 + j_3}$ when its two last columns are interchanged. However, the interchange of two other columns cannot be described by a simple symmetry property. Nevertheless, the f_r symbol is of central importance for the Wigner-Eckart theorem in the $\{J^2, U_r\}$ scheme (see eq.(17) below).

Following ref.[29], we define a more symmetrical symbol, namely the \bar{f}_r symbol, through

$$\bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix} = \frac{1}{\sqrt{(2j_1 + 1)(2j_2 + 1)(2j_3 + 1)}} \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} \sum_{m_3=-j_3}^{j_3} \\ \times q_1^{-\alpha_1 m_1} q_2^{-\alpha_2 m_2} q_3^{-\alpha_3 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \quad (14)$$

where the parameters q_i are given by eq.(12) with $q \equiv q_i$ and $j \equiv j_i$ for $i = 1, 2, 3$. The $3 - jm$ symbol on the right-hand side of eq.(14) is an ordinary Wigner symbol for the group $SU(2)$ in the $SU(2) \supset U(1)$ basis. As a matter of fact, it is possible to pass from the f_r symbol to the \bar{f}_r symbol and vice versa by means of a metric tensor. The \bar{f}_r symbol is more symmetrical than the f_r symbol. The \bar{f}_r symbol exhibits the same symmetry properties under permutations of its columns as the $3 - jm$ Wigner symbol : Its value is multiplied by $(-1)^{j_1 + j_2 + j_3}$ under an odd permutation and does not change under an even permutation. In addition, the orthogonality properties of the highly symmetrical \bar{f}_r symbol easily follow from the corresponding properties of the $3 - jm$ Wigner symbol. Thus, we have

$$\sum_{j_3 \alpha_3} (2j_3 + 1) \bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix}^* \bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha'_1 & \alpha'_2 & \alpha_3 \end{pmatrix} = \delta(\alpha'_1, \alpha_1) \delta(\alpha'_2, \alpha_2) \quad (15)$$

and

$$\sum_{\alpha_1 \alpha_2} \bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix} \bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3' \\ \alpha_1 & \alpha_2 & \alpha_3' \end{pmatrix}^* = \frac{1}{2j_3 + 1} \Delta(0|j_1 \otimes j_2 \otimes j_3) \delta(j_3', j_3) \delta(\alpha_3', \alpha_3) \quad (16)$$

where $\Delta(0|j_1 \otimes j_2 \otimes j_3) = 1$ or 0 according to as the Kronecker product $(j_1) \otimes (j_2) \otimes (j_3)$ contains or does not contain the identity irreducible representation (0) of SU(2). Observe that the real number r is the same for all the \bar{f}_r symbols occurring in eqs.(15) and (16).

The values of the SU(2) CGC's in the $\{J^2, U_r\}$ scheme as well as of the f_r and \bar{f}_r coefficients are not necessarily real numbers. For instance, we have the following property under complex conjugation

$$\bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix}^* = (-1)^{j_1 + j_2 + j_3} \bar{f}_r \begin{pmatrix} j_1 & j_2 & j_3 \\ \alpha_1 & \alpha_2 & \alpha_3 \end{pmatrix}$$

Hence, the value of the \bar{f}_r coefficient is real if $j_1 + j_2 + j_3$ is even and pure imaginary if $j_1 + j_2 + j_3$ is odd. Then, the behavior of the \bar{f}_r symbol under complex conjugation is completely different as the one of the ordinary 3 - jm Wigner symbol.

Finally, it is worth to mention that the recoupling coefficients of the group SU(2) can be expressed in terms of coupling coefficients of SU(2) in the $\{J^2, U_r\}$ scheme. For example, the 9 - j symbol can be expressed in terms \bar{f}_r symbols by replacing, in its decomposition in terms of 3 - jm symbols, the 3 - jm symbols by \bar{f}_r symbols. On the other hand, the decomposition of the 6 - j symbol in terms of \bar{f}_r symbols requires the introduction of six metric tensors corresponding to the six arguments of the 6 - j symbol. These matters may be developed by following the approach initiated in refs.[29-32].

4.2 Wigner-Eckart Theorem in the $\{J^2, U_r\}$ Scheme

From the spherical components $T_q^{(k)}$ (with $q = -k, -k + 1, \dots, k$) of an SU(2) irreducible tensor operator $\mathbf{T}^{(k)}$, we define the $2k + 1$ components

$$T_\alpha^{(k)}(r) = \frac{1}{\sqrt{2k + 1}} \sum_{m=-k}^k q^{\alpha m} T_m^{(k)}$$

with

$$\alpha = -kr, -kr + 1, \dots, -kr + 2k, \quad 2k \in \mathbf{N}$$

In the $\{J^2, U_r\}$ scheme, the Wigner-Eckart theorem reads

$$\langle \tau_1 j_1 \alpha_1; r | T_\alpha^{(k)}(r) | \tau_2 j_2 \alpha_2; r \rangle = (\tau_1 j_1 || T^{(k)} || \tau_2 j_2) f_r \begin{pmatrix} j_1 & j_2 & k \\ \alpha_1 & \alpha_2 & \alpha \end{pmatrix} \quad (17)$$

where $(\tau_1 j_1 || T^{(k)} || \tau_2 j_2)$ denotes an ordinary reduced matrix element. Such an element is basis-independent. Therefore, it does not depend on the labels α_1, α_2 and α . On the contrary, the f_r coefficient in eq.(17), defined by eq.(13), depends on the labels α_1, α_2 and α .

5 Concluding Remarks

In this paper, we have developed a quon approach to the Lie algebra of the classical (not quantum!) group $SU(2)$. Such an approach leads to the polar decomposition of the generators J_+ and J_- of $SU(2)$, a decomposition originally introduced by Lévy-Leblond [28].

The familiar $\{J^2, J_3\}$ quantization scheme with the (usual) standard spherical basis $\{|jm\rangle : 2j \in \mathbf{N}, m = -j, -j + 1, \dots, j\}$, corresponding to the canonical chain of groups $SU(2) \supset U(1)$, is thus replaced by the $\{J^2, U_r\}$ quantization scheme with a (new) basis, namely, the non-standard basis $B_r = \{|j\alpha; r\rangle : 2j \in \mathbf{N}, \alpha = -jr, -jr + 1, \dots, -jr + 2j\}$. We have given the premises of the construction of the Wigner-Racah algebra of the group $SU(2)$ in the B_r basis. Of course, there exists an infinity of B_r bases due to the fact that $r \in \mathbf{R}$. The case $r = 1$ probably deserves a special attention. We shall give elsewhere a complete development of the Wigner-Racah algebra of $SU(2)$ in the B_1 basis. In particular, the calculation and the properties, including Regge symmetry properties, of the coupling coefficients (\bar{f}_1 and f_1 symbols and CGc's in the $\{J^2, U_1\}$ scheme) shall be the object of a forthcoming paper.

As a further interesting step, it would be interesting to find realizations of the B_r basis (i) on the sphere S^2 for j integer and (ii) on the Fock-Bargmann spaces (of entire analytical functions) in 1 and 2 dimensions for j integer or half of an odd integer. In this respect, the problem of finding a differential realization of the operator U_r on S^2 and of expressing its eigenfunctions

$$[y_r]_{\ell\alpha}(\theta, \varphi) = \frac{1}{\sqrt{2\ell+1}} \sum_{m=-\ell}^{\ell} q^{\alpha m} Y_{\ell m}(\theta, \varphi) \quad (18)$$

with

$$\alpha = -\ell r, -\ell r + 1, \dots, -\ell r + 2\ell, \quad \ell \in \mathbf{N}$$

as special functions is very appealing. (In eq.(18), $Y_{\ell m}$ denotes a spherical harmonic.)

6 Acknowledgments

One of the authors (M.K.) wishes to thank Jean-Marc Lévy-Leblond (from the Université Sophia Antipolis, Nice, France) for attracting his attention on ref.[28] sixteen years ago. He also wishes to thank Jacob Katriel (from the Israel Institute of Technology, Haifa, Israel) for discussions which contributed to clarify the question of the spectrum of the operator $U_{r=0}$ eight years ago. Finally, he would like to thank the organizing committee of the International Workshop *Quantum Groups, Deformations and Contractions* for inviting him to give a talk from which the present paper is a by-product.

References

- [1] J.M. Leinaas and J. Myrheim, *Nuovo Cimento* **B37** (1977) 1.
- [2] G.A. Goldin, R. Menikoff and D.H. Sharp, *J. Math. Phys.* **21** (1980) 650 ; **22** (1981) 1664.
- [3] J. Beckers and N. Debergh, *Nucl. Phys.* **B340** (1990) 767.
- [4] D. Bonatsos, P. Kolokotronis and C. Daskaloyannis, *Mod. Phys. Lett.* **A10** (1995) 2197.
- [5] A. Mostafazadeh, *Int. J. Mod. Phys.* **A11** (1996) 2957.
- [6] M.-l. Ge and G. Su, *J. Phys.* **A24** (1991) L721.
- [7] C.R. Lee and J.-P. Yu, *Phys. Lett.* **A164** (1992) 164.
- [8] G. Su and M.-l. Ge, *Phys. Lett.* **A173** (1993) 17.
- [9] J.A. Tuszyński, J.L. Rubin, J. Meyer and M. Kibler, *Phys. Lett.* **A175** (1993) 173.
- [10] V.I. Man'ko, G. Marmo, S. Solimeno and F. Zaccaria, *Phys. Lett.* **A176** (1993) 173.
- [11] R.-R. Hsu and C.-R. Lee, *Phys. Lett.* **A180** (1993) 314.
- [12] Ya.I. Granovskii and A.S. Zhedanov, *Mod. Phys. Lett.* **A8** (1993) 1029.
- [13] M. Chaichian, R.G. Felipe and C. Montonen, *J. Phys.* **A26** (1993) 4017.
- [14] S. Vokos and C. Zachos, ANL-HEP-CP-93-39.
- [15] R.K. Gupta, C.T. Bach and H. Rosu, *J. Phys.* **A27** (1994) 1427.
- [16] M.A. R.-Monteiro, I. Roditi and L.M.C.S. Rodrigues, *Phys. Lett.* **A188** (1994) 11.
- [17] R.-S. Gong, *Phys. Lett.* **A199** (1995) 81.
- [18] M. Daoud and M. Kibler, *Phys. Lett.* **A206** (1995) 13.
- [19] M.R. Kibler, J. Meyer and M. Daoud, *On qp -Deformations in Statistical Mechanics of Bosons in D Dimensions*, in *Symmetry and Structural Properties of Condensed Matter*, eds. T. Lulek, W. Florek and B. Lulek (World Scientific, Singapore, 1977), page 460.
- [20] L.C. Biedenharn, *J. Phys.* **A22** (1989) L873.
- [21] A.J. Macfarlane, *J. Phys.* **A22** (1989) 4581.
- [22] G. Rideau, *Lett. Math. Phys.* **24** (1992) 147.
- [23] M.R. Kibler, *Introduction to Quantum Algebras*, in *Symmetry and Structural Properties of Condensed Matter*, eds. W. Florek, D. Lipiński and T. Lulek (World Scientific, Singapore, 1993), page 445.
- [24] M. Arik and D.D. Coon, *J. Math. Phys.* **17** (1976) 524.
- [25] D.B. Fairlie, P. Fletcher and C.K. Zachos, *J. Math. Phys.* **31** (1990) 1088.
- [26] M. Daoud, Y. Hassouni and M. Kibler, *The k -Fermions as Objects Interpolating between Fermions and Bosons*, in *Symmetries in Science X*, eds. B. Gruber and M. Ramek (Plenum Press, New York, 1998).

- [27] M. Daoud, Y. Hassouni and M. Kibler, *Generalized Super-Coherent States*, *Yad. Fiz.* (submitted for publication).
- [28] J.-M. Lévy-Leblond, *Rev. Mex. Física* **22** (1973) 15.
- [29] M. Kibler, *J. Molec. Spectrosc.* **26** (1968) 111 ; *Int. J. Quantum Chem.* **3** (1969) 795.
- [30] M. Kibler, *C. R. Acad. Sci. (Paris)* **B268** (1969) 1221.
- [31] M.R. Kibler, *J. Math. Phys.* **17** (1976) 855 ; *J. Molec. Spectrosc.* **62** (1976) 247 ; *J. Phys.* **A10** (1977) 2041.
- [32] M.R. Kibler and P.A.M. Guichon, *Int. J. Quantum Chem.* **10** (1976) 87 ; M.R. Kibler and G. Grenet, *Int. J. Quantum Chem.* **11** (1977) 359 ; M.R. Kibler, *Int. J. Quantum Chem.* **23** (1983) 115.