Towards an understanding of lineage speci cation in hem atopoietic stem cells: A mathematical model for the interaction of transcription factors GATA -1 and PU .1

Ingo Roeder Ingm ar Glauche

Institute for M edical Inform atics, Statistics and Epidem iology, University of Leipzig, H aertelstr. 16/18, D-04107 Leipzig, G erm any

Abstract

In addition to their self-renewal capabilities, hem atopoietic stem cells guarantee the continuous supply of fully di erentiated, functional cells of various types in the peripheral blood. The process which controls di erentiation into the di erent lineages of the hem atopoietic system (erythroid, m yeloid, lym phoid) is referred to as lineage speci cation. It requires a potentially multi-step decision sequence which determ ines the fate of the cells and their successors. It is generally accepted that lineage speci cation is regulated by a complex system of interacting transcription factors. How ever, the underlying principles controlling this regulation are currently unknown.

Here, we propose a simple quantitative model describing the interaction of two transcription factors. This model is motivated by experimental observations on the transcription factors GATA -1 and PU 1, both known to act as key regulators and potential antagonists in the erythroid vs. myeloid di erentiation processes of hem atopoietic progenitor cells. We demonstrate the ability of the model to account for the observed switching behavior of a transition from a state of low expression of both factors (undi erentiated state) to the dominance of one factor (di erentiated state). Depending on the parameter choice, the model predicts two di erent possibilities to explain the experimentally suggested, stem cell characterizing priming state of low level co-expression. Whereas increasing transcription rates are su - cient to induce di erentiation in one scenario, an additional system perturbation (by stochastic uctuations or directed impulses) of transcription factor levels is required in the other case.

Keywords: lineage speci cation, hem atopoietic stem cell, transcription factor network, PU 1, GATA-1

1 Introduction

The hem atopoietic system consists of a variety of functionally di erent cell types, including mature cells such as erythrocytes, granulocytes, platelets, or lym phocytes, as well as several di erent precursor cells (i.e., prem ature cell stages) and hem atopoietic stem cells (HSC) (Lord, 1997; Orkin, 2000). Most mature cell types have limited life spans ranging from a few hours to several months, which implies the existence of a source capable of replenishing these di erentiated cells throughout the life span of an individual. This supply is realized by the population of HSC, which is maintained and even regenerated after in jury or depletion throughout the whole life of the organism . This selfrenewal property is a major characteristic de ning HSC (Loe er and Roeder, 2002; Lord, 1997; Potten and Loe er, 1990). A second major characteristic of HSC is their ability to contribute to the production of cells of all hem atopoietic lineages, thus ensuring the supply of functionally di erentiated cells meeting the needs of the organism. The process controlling the developm ent of undifferentiated stem or progenitor cells into one speci c functional direction (i.e., one speci c hem atopoietic lineage) is called lineage speci cation. It is generally accepted that the process of lineage speci cation is governed by the interplay of m any di erent transcription factors (A kashi, 2005; C antor and O rkin, 2002; Cross et al., 1994; Orkin, 1995, 2000; Tenen, 2003). Experim ental results suggest that a number of relevant transcription factors are expressed simultaneously in HSC, although at a low level (A kashiet al., 2003; Hu et al., 1997). Som e authors refer to this state of a low level co-expression as prim ing behavior (A kashi, 2005; C ross and E nver, 1997; E nver and G reaves, 1998). During di erentiation the balanced co-expression of these potentially antaqonistic transcription factors is assumed to be broken at some point (or even multiple points). Thereafter, the system is supposed to be characterized by an up-regulated level of some transcription factors, specic for a particular lineage, while other transcription factors are down-regulated. These observations suggest a transcription factor network, capable of switch-like behavior by changing from unspeci c co-expression to di erent states of speci c expression. How ever, the general underlying principles of the regulatory mechanism s are currently unknown. Particularly, it is unclear whether the assumption of a dynam ically balanced low level co-expression state is justilled or whether priming should rather be interpreted as the result of an inactive transcription factor network overlaid by stochastic uctuations of transcription factor expression.

In this paper we propose a simple mathematical model describing di erent

Corresponding author.Tel.: + 49 (0)341 97 16 111; fax: + 49 (0)341 97 16 109 Em ailaddresses: ingo.roeder@imise.uni-leipzig.de (Ingo Roeder),

ingmar.glauche@imise.uni-leipzig.de (Ingm arG lauche).

interaction scenarios of two transcription factors. Biologically, this simple two component network model is motivated by experimental observations on the transcription factors GATA-1 and PU 1, known to be involved in the process of lineage speci cation of HSC (Duetal, 2002; O ikawa et al., 1999; Rekhtm an et al., 1999; Rosm arin et al., 2005; Tenen, 2003; Voso et al., 1994). The zinc nger factor GATA -1 is reported to be required for the differentiation and m aturation of erythroid/m egakaryocytic cells, while the Etsfam ily transcription factor PU 1 supports the development of myeloid and lym phoid cells (reviewed by C antor and O rkin, 2002; Tenen, 2003). For both, GATA-1 and PU 1, it has been demonstrated that they are able to stimulate their own transcription through an auto-catalytic process (Chen et al., 1995; N ishim ura et al., 2000; O kuno et al., 2005; T saiet al., 1991). A dditionally, there are physical interactions between GATA -1 and PU .1 which induce a mutual inhibition and, therefore, favor one lineage choice at the expense of the other (erythroid/m egakaryocyte vs. m yeloid) (Du et al, 2002; N erbv et al, 2000; Rekhtm an et al., 1999, 2003; Voso et al., 1994; Yam ada et al., 1998; Zhang et al., 1999, 2000). In particular, two di erent mechanisms for the mutual inhibition of these two transcription factors have been suggested by experim ental observations: O n one hand, GATA -1 binds to the 3=4 region of PU.1 (complex 1) and displaces the PU.1 co-activator c-Jun from its binding site, thereby, inhibiting the transcription initiation of PU 1 (Zhang et al., 1999). On the other hand, the inhibition of GATA-1 transcription is mediated by the binding of the N-term inal region of PU 1 to the C-nger region of GATA-1 (complex 2), thus blocking the binding of GATA-1 to its prom oter (Zhang et al., 2000). That means, although both inhibition mechanisms are interfered through the form ation of PU 1/GATA-1 heterodim ers, the two com plexes are structurally di erent. W hereas com plex 1 (inhibition of PU.1 transcription by GATA-1) is known to bind to DNA, thus occupying a PU 1 prom oter site, DNA-binding of com plex 2 (inhibition of GATA-1 transcription by PU 1) has not been reported so far.

The mechanisms of antagonistic interdependence together with positive autocatalytic regulation provide a fram ework for the theoretical investigation of di erent scenarios of transcription factor interaction and their implications for the explanation of lineage speci cation control. Applying a mathematical model, which formalizes the described interactions, it is now possible to analyze di erent combinations of transcription factor activation and inhibition on a qualitative and quantitative level. The proposed model relies on principles suggested for the description of general genetic switches (e.g. Becskei et al., 2001; C inquin and D emongeot, 2002, 2005; G ardner et al., 2000).

In this paper it is our objective to exam ine the following questions within the fram ework of this model structure:

A re the experim entally described interactions of the two transcription fac-

tors su cient to generate a switching behavior between a stable co-expression of two factors and the dom inance of one of these factors?

W hat are the conditions inducing such a qualitative change in the system behavior?

Is there evidence for a functional role of the (experim entally suggested) priming status?

To answer these question the following strategy is applied. Firstly, the model equations are derived on the basis of the described biological mechanisms of transcription factor interaction for GATA -1 and PU 1 (Section 2). Secondly, this model is analyzed with respect to the existence of steady state solutions and their dependence on the model parameters. A coording to our objective, to understand the mechanisms leading to switches between di erent stable system states, we focus our analysis particularly on the determination of bi-furcation conditions, considering di erent scenarios of transcription factor interaction (Section 3). Finally, the obtained results are discussed in relation to the ongoing debate about lineage speci cation control in the hem atopoietic system, speci cally with respect to potential explanations of the experim entally suggested low level co-expression of transcription factors (priming) in undi erentiated progenitors and stem cells (Section 4).

2 M odel description

A lthough our analysis is motivated by experimental observations of specic transcription factor interactions (GATA-1 and PU 1), the model may also be applied in the general context of two interacting transcription factors. In the following, the two transcription factors are denoted by X and Y.

2.1 General assumptions

The general design of the model structure is based on the following assum p-tions which are motivated by the experim ental observations outlined in Section 1:

Both transcription factors, X and Y, are able to act as activator m olecules:

- If bound to their own promoter region, X and Y introduce a positive feedback on their own transcription. This process is referred to as speci c transcription (Fig. 1(a)).
- X and Y are both able to induce an overall transcription which also e ects potentially antagonistic transcription factors. A lthough such an interaction is most likely indirect, for the model we consider a mutual

activation of X and Y by the opposing transcription factor, which we refer to as unspeci c transcription (Fig. 1(b)).

W e assume that transcription initiation is only achieved by the simultaneous binding of two X and Y molecules, respectively (i.e., binding cooperativity c = 2). This assumption is motivated by the result that a binding cooperativity c > 1 is a necessary condition for the existence of system bistability (see e.g. Becskei et al., 2001; C inquin and D emongeot, 2005; G ardner et al., 2000).

There is a mutual inhibition of X and Y.W ithin this context, two possible mechanisms, based on the formation of two structurally dierent complexes of X and Y, are considered:

- Joint binding of X and Y molecules to promoter sites (Fig. 1(c)). Here, the DNA-bound X Y -complex (Z₁) acts as a transcription repressor, which blocks the binding sites. This represents a mode of competition for free binding sites.
- Form ation of another X Y -com plex, called Z₂, which neither binds to X nor Y DNA binding site (Fig. 1 (d)). In contrast to Z₁, this represents a competition for free transcription factor molecules.

Both inhibition mechanisms (including combinations of them) are considered for X as well as for Y .

To facilitate the analysis of the mathematical model we make the following simplications:

Post-transcriptional regulation is neglected, i.e., the transcription of a gene is considered to ultimately result in the production of the corresponding protein (here, a transcription factor).

T in e delays due to transcription and translation processes are neglected. Simultaneous binding of X /Y m onom ers together with a Z_1 -heterodim er, of two Z_1 -heterodim ers, as well as of a X and a Y m onom er at the same prom oter are excluded from the analysis.

Interactions of X, Y as well as the prom oter regions of the coding genes with further transcription factors are neglected.

Throughout the paper the follow ing notations are used:x;y denote the m olecule concentrations of X and Y, respectively. Z_1 denotes the DNA bound XY - com plex and Z_2 the structurally di erent XY - com plex, which is not able to bind to prom oter DNA. $D_{x=y}$ denotes free DNA binding sites within the promoter region of X and Y, respectively. In contrast, binding sites occupied by X or Y molecules or by the XY -com plex Z_1 are denoted as $D_{x=y}^{xx=yy=xy}$.



Figure 1. Principles of transcription initiation and inhibition for X and Y. (a) Speci c transcription, i.e. auto-catalysis by the transcription factor itself; (b) Unspeci c transcription, i.e. transcription initiated by another transcription factor; (c-e) Suggested mechanisms of transcription inhibition for X and Y by formation of X Y -com plexes: (c) A X Y -com plex, called Z_1 , bound to the promoter regions acts as a repressor; (d) The formation of a structurally di erent X Y -com plex (Z_2) com petitively inhibits the DNA binding of X and Y molecules; (e) C om bination of (c) and (d) as suggested for GATA-1 and PU 1 (Zhang et al., 1999, 2000)

2.2 Model equations

W ith these assumptions one can write down a set of them ical reaction equations which underly the system dynamics.

The processes of specic and unspecic transcription activation (see Fig.

1(a), (b)) are described by equations (1)-(4).

$$X + X + D_x^{K_1} D_x^{XX}; D_x^{XX}; D_x^{XX} + X$$
 (1)

$$Y + Y + D_x \stackrel{K_2}{\longrightarrow} D_x^{YY}; \quad D_x^{YY} \stackrel{\mu_x}{:} D_x^{YY} + X$$
(2)

$$Y + Y + D_{y} \stackrel{K_{3}}{\longrightarrow} D_{y} \stackrel{Yy}{\longrightarrow} D_{y} \stackrel{Yy}{\longleftarrow} D_{y} \stackrel{Yy}{\longrightarrow} + Y$$
(3)

$$X + X + D_{y}^{K_{4}} D_{y}^{xx}; D_{y}^{xx} \stackrel{U_{y}}{!} D_{y}^{xx} + Y$$
(4)

Herein we made the simplifying assumption that the DNA binding of X and Y always occurs as the binding of hom odim ers. That means, that the sequential binding of two monom ers, as the second possibility of DNA binding, is not consider. The process of dimerization, as well as the DNA binding and dissociation, are regarded to be in quasi steady state.

Here and throughout the paper, the K_i = $k_i = k_i$ (i = 1; :::;7) denote the equilibrium (dissociation) constants of the reactions, with k_i and k_i representing the forward and backward reaction rate constants, respectively. Finally, it is assumed that both transcription factor monomers, X and Y, decay with rst order kinetics at rates k_0^x and k_0^y , whereas dimension plexes are assumed to be stable.

The di erent mutual transcription inhibition mechanisms are illustrated in Figs.1(c)-(e).First, we consider the form ation the X Y -complex Z_2 (see Fig. 1(d))

$$X + Y \xrightarrow{K_5} Z_2$$
: (5)

Under the quasi steady state assumption Z_2 does not contribute to the mathematical description of the system dynamics.

As shown in Fig. 1(c), there is also the possibility that X and Y form a structurally di erent heterodim er Z_1 , which is able to bind to the promoter regions, acting as a repressor for X and Y transcription, respectively:

$$X + Y + D_{x} \stackrel{K_{6}}{\longrightarrow} D_{x}^{xy};$$
 (6)

$$X + Y + D_{y} \stackrel{K_{\gamma}}{\rightarrow} D_{y} \stackrel{xy}{\rightarrow}$$
(7)

As with the promoter binding of X and Y, we collapse dimerization, which is assumed to be in quasi steady state, and DNA binding into one process, neglecting the sequential binding of monomers.

Under the posted quasi steady state assumptions, equations (1)-(7) lead to

the following set of ordinary di erential equations:

$$\frac{dx}{dt} = k_{0_x}x + \frac{s_xK_1x^2 + u_xK_2y^2}{1 + K_1x^2 + K_2y^2 + K_6xy}$$
(8)

$$\frac{dy}{dt} = k_{0_y}y + \frac{s_yK_3y^2 + u_yK_4x^2}{1 + K_3y^2 + K_4x^2 + K_7xy}$$
(9)

Details of the derivation are given in Appendix A.

3 Results

3.1 Symmetric system

To analytically derive steady state as well as potential bifurcation conditions, we restrict ourself in this section to the special case of a completely symmetric system, i.e.: $k_{0_x} = k_{0_y} = k_0$, $s_x = s_y = s$, $u_x = u_y = w$, $K_{p1} = K_3$, $K_{2} = \frac{K_4}{K_1}$, and $K_6 = K_7$. Using these relations, together with $x = \frac{P}{K_1} \frac{K_1}{K_1} x$, $y = \frac{K_1}{K_1} y$, $k_u = K_2 = K_1$, $k_r = K_6 = K_1$, $s = \frac{P}{K_1} \frac{s}{K_1} s = k_0$, $u = \frac{P}{K_1} \frac{w}{K_1} = k_0$, and $k_0 = k_0 t$, the system (8), (9) can be written in a dimensionless form as

$$\frac{dx}{d} = x + \frac{sx^2 + uk_u y^2}{1 + x^2 + k_u y^2 + k_r x y};$$
(10)

$$\frac{dy}{d} = y + \frac{sy^2 + uk_u x^2}{1 + k_u x^2 + y^2 + k_r x y};$$
(11)

Equations (10) and (11) are a pair of coupled rst order di erential equations. The steady state ($\underline{x} = \underline{y} = 0$) is de ned in plicitly by

$$\mathbf{x} = \frac{\mathbf{s}\mathbf{x}^{2} + \mathbf{u}\mathbf{k}_{u}\mathbf{y}^{2}}{1 + \mathbf{x}^{2} + \mathbf{k}_{u}\mathbf{y}^{2} + \mathbf{k}_{r}\mathbf{x}\mathbf{y}};$$
(12)

$$y = \frac{sy^2 + uk_u x^2}{1 + k_u x^2 + y^2 + k_r x y}$$
 (13)

The domain of these nullclines for x and y is restricted by the choice of parameters as outlined in Appendix B. The intersections of the nullclines correspond to the xed points (x; y) of the dimensial equations (10) and (11). Fixed points on the diagonal (x; x) are traced under the simplifying condition x = y. In this case, equations (12) and (13) can be summarized by

$$\mathbf{x} = \frac{\mathbf{x}^{2} (\mathbf{s} + \mathbf{u} \mathbf{k}_{u})}{1 + \mathbf{x}^{2} (1 + \mathbf{k}_{u} + \mathbf{k}_{r})} ;$$
(14)

The rst (trivial) xed point of equation (14) is $x_1 = 0$. Having eliminated this solution, the remaining quadratic equation yields two further non-trivial xed points at

$$\mathbf{x}_{2=3} = \frac{(s + uk_u)}{2(1 + k_u + k_r)^2} \frac{q}{(s + uk_u)^2} \frac{4(1 + k_u + k_r)}{2(1 + k_u + k_r)} :$$
(15)

 $(x_2;x_2)$ and $(x_3;x_3)$ are real xed points on the diagonal for

s
$$uk_u + 2 \frac{q}{1 + k_u + k_r}$$
: (16)

B ifurcation points can be characterized by nullclines intersecting with equal slopes. The derivatives of equations (10) and (11) are evaluated to determ ine explicit conditions for bifurcation occurrence on the diagonal, considering s as the bifurcation parameter¹. For simplicity the denom inators in equations (12) and (13) are de ned as $P_x = 1 + x^2 + k_u y^2 + k_r xy$ and $P_y = 1 + k_u x^2 + y^2 + k_r xy$. The partial derivative of equation (12) with respect to y leads to

$$x^{0} = \frac{(2sxx^{0} + 2uk_{u}y)P_{x}}{P_{x}^{2}} (sx^{2} + uk_{u}y^{2})P_{x}^{0}}$$
(17)

with $x^0 = 0x = 0y$ and $P_x^0 = 0P_x = 0y = x^0(2x + k_ry) + 2k_uy + k_rx$. Solving for x^0 yields

$$x^{0} = \frac{2uk_{u}yP_{x}}{P_{x}^{2}} \frac{(sx^{2} + uk_{u}y^{2})(2k_{u}y + k_{r}x)}{(sx^{2} + uk_{u}y^{2})(2x + k_{r}y)};$$
(18)

For bifurcation points on the diagonal (x = y), where the denom inators P_x and $P_y \sin p \operatorname{lify}$ to $P = 1 + x^2 (1 + k_u + k_r)$, equation (18) can be rewritten as

$$x^{0}(P^{2} 2sxP + x^{3}(s+uk_{u})(2+k_{r})) = 2uk_{u}xP x^{3}(s+uk_{u})(2k_{u}+k_{r})$$
: (19)

Inserting P in the form $P = x(s + uk_u)$ derived from equation (14) and neglecting the trivial solution the equality now reads

$$x^{0}(uk_{u} + x(2 + k_{r})) = 2uk_{u} + x(2k_{u} + k_{r})$$
: (20)

To nd the bifurcation points on the diagonal one needs to study the two distinct cases for $x^0 = 1$ and $x^0 = -1$ (see Appendix C).

Case I $(x^0 = 1)$:

¹ Param eter s is chosen to account for changes in the transcriptional activity by enhancer actions or modi cations in chromatin structure. Furthermore, s is the critical param eter that gives rise to the di erent distinct domains for the nullclines as outlined in Appendix B.

Equation (20) satis es the condition $x^0 = 1$ at

$$\mathbf{x}_{\mathbf{x}^{0}=1} = \frac{\mathbf{s} + \mathbf{u}\mathbf{k}_{u}}{2(1 + \mathbf{k}_{u} + \mathbf{k}_{r})}$$
(21)

This only coincides with the xed points $x_{2=3}$ derived in equation (15) if the expression under the radical in equation (15) vanishes, i.e., $x_2 = x_3$. This is true for q

$$s_1 = uk_u + 2 \frac{1}{1 + k_u + k_r}$$
; (22)

which corresponds to the condition de ned in (16). This implies that the \birth" of the xed points $(x_2;x_2)$ and $(x_3;x_3)$ coincides with the bifurcation condition $x^0 = 1$. The sequence of nullclines in Fig. 2(a)-(c) illustrates this behavior for an unspeci c transcription rate u = 1. The nullclines in Fig. 2(a) do not intersect for $s < s_1$, i.e., there is no non-trivial xed point along the diagonal. For $s = s_1$ one common xed point at $x_2(s_1) = x_3(s_1) = (s_1 + uk_u)=2(1 + k_u + k_r)$ exists, which marks the bifurcation point depicted in Fig. 2(b). For $s > s_1$ two distinct xed points $(x_2;x_2)$ and $(x_3;x_3)$ exist on the diagonal, shown in Fig. 2(c). Whereas the upper point at $(x_2;x_2)$ is stable, the lower one at $(x_3;x_3)$ is unstable. The nullclines change qualitatively for a further increase in the bifurcation parameter s as shown in the sequence Fig. 2(d), (e).

In the case of a smaller unspecie c transcription rate u = 0.4 the corresponding bifurcation is illustrated in Fig. 3(d). The two xed points $(x_2; x_2)$ and $(x_3; x_3)$ generated at the diagonal are both unstable as depicted in Fig. 3(e), (f). The qualitative dimensions between the scenarios for small and large unspecie c transcription u are more thoroughly investigated in the subsequent paragraphs.

Case 2 (
$$x^0 = 1$$
)

W hen $x^0 = 1$ equation (20) simplies to

:

$$x_{x^{0}=1} = \frac{3uk_{u} \quad s}{2k_{u} \quad 2}$$
(23)

Equating $x_{x^{0}=1} = x_{2=3}$ from equation (15) leads to a dependency on the parameters s, u, k_u and k_r . Two further bifurcation points are obtained at:

$$s_{2} = \frac{uk_{u} (1 + 3k_{r} + 5k_{u}) + 2(k_{u} - 1)^{q} \overline{1 - k_{r} - 3k_{u} + 4k_{u}^{2}u^{2}}}{(1 + k_{r} + 3k_{u})}$$
(24)

$$s_{3} = \frac{uk_{u}(1+3k_{r}+5k_{u}) \quad 2(k_{u} \quad 1)^{4} \overline{1 \quad k_{r} \quad 3k_{u}+4k_{u}^{2}u^{2}}}{(1+k_{r}+3k_{u})}$$
(25)

To guarantee the existence of these bifurcations at the diagonal, $s_{2=3} = s_1$ is required. The case $s_{2=3} < s_1$ indicates that bifurcations occur of the diagonal.











Figure 2. Deform ation of the nullclines for increasing values of the bifurcation parameter s. The occurrence of the st bifurcation at s is depicted in (b), the second bifurcation at $s_1 = s_2$ in (g). Parameters are $k_u = 1$, $k_r = 0$ and u = 1. The bifurcation parameter is set to s = 1:7 (a), $s = s_1 = 1 + 2$ 2 1.83 (b), s = 1:9 (c), s = 1:99 (d), s = 2:01 (e), s = 2:6 (f), $s = s_2 = s_3 = 3u = 3$ (g), s = 3:8 (h). Fixed points are marked as follows: trivial xed point (0;0) - , stable/unstable xed point (x;x_2) - N/4, unstable xed point (x;x_3) - 0, stable/unstable xed

points o the diagonal – /, bifurcation point – .



Figure 3.D eform ation of the nullclines for increasing values of the bifurcation parameters for unspeci c transcription rate u = 0.4. Parameters are $k_i = 1$ and $k_r = 0$. The bifurcation parameter is set to s = 1.9 (a), s = 2.04 (b), s = 2.1 (c), $s = s_1 = 2.43$ (d), s = 2.8 (e), s = 3.2 (f). Fixed points are marked according to the caption in Fig.2.

For the special case $k_u = 1$ the conditions for the occurrence of these bifurcations simplify to $s_2 = s_3 = 3u$ (given $s_2 = s_3 > s_1$ which is true for u > 1 = 2). Since this condition is valid for both xed points, $(x_2;x_2)$ and $(x_3;x_3)$, it indicates that the bifurcations occur at the same bifurcation parameter s = 3u. Fig. 2 (f)-(h) depicts the bifurcations for both xed points in the case u = 1, $k_u = 1$. A fier a deform ation of the nullclines the intersections in Fig. 2 (f) still represent the xed points $(x_2;x_2)$ and $(x_3;x_3)$ for $s < s_2 = s_3$. In Fig. 2 (g) the nullclines for $s = s_2 = s_3$ intersect with the same bocal slope at x_2 as well as at x_3 . This marks the bifurcation point for both xed points, that coincides for $k_u = 1$. Fig. 2 (h) illustrates the new xed points o the diagonal, which are stable bifurcating from $(x_2;x_2)$ and unstable bifurcating from $(x_3;x_3)$. The xed point $(x_2;x_2)$ itself changes the stability and becomes unstable, $(x_3;x_3)$ remains unstable as before.

For small u the condition for the occurrence of further bifurcations $s_{2=3} = s_1$ is violated. Numerical results indicate that two saddle-node bifurcations form



Figure 4.B ifurcation diagram s x vs. s with $k_u = 0.8$, $k_r = 0$, u = 1 (a) and 0 = 0.4 (b). The stability of the steady states is coded as follow s: solid line - stable, dashed line - unstable.

xed points o the diagonal at $s < s_1$ as depicted in the sequence of nullclines in Figs. 3 (b), (c). The saddle-node bifurcation on the diagonal is observed at s_1 . For large s these scenarios show a comparable pattern of two up-regulated steady states with one high and one low expressed component and a further stable xed point at (0;0) (compare Figs. 2 (h) and 3 (f)).

The bifurcation diagram s in Fig. 4 comprise the above ndings for $k_u = 0.8$. The x-coordinate for the xed points is shown depending on the bifurcation parameter s. For u = 1 (Fig. 4(a)) the birth of two xed points through a saddle-node bifurcation can be seen at s_1 , given by equation (22). Condition (24) de nes the occurrence of the pitchfork bifurcation on the upper branch $(x_2;x_2)$ at s_2 , whereas condition (25) is the equivalent for the lower branch $(x_3; x_3)$ at s_3 . Note that the additional condition $s_{2=3}$ s_1 is fulled. The upper branch gives rise to three xed points, one unstable (arising from the existing stable xed point) at the diagonal at x_2 and two new stable xed points branching o this axis. For the lower case all three xed points are unstable for $s > s_3$. The inset in Fig. 4 (a) enlarges this bifurcation occurring at s_3 . Fig. 4 (b) illustrates the equivalent scenario for u = 0.4. The saddle-node bifurcation at s_1 represents the form ation of the unstable xed points $(x_2;x_2)$ and $(x_3; x_3)$. In addition, two further saddle node bifurcations exist that also form stable xed points. Since $s_{2=3} < s_1$ these bifurcations do not occur on the diagonal. All branches in Fig. 4 that do not represent xed points on the diagonal were determ ined num erically.

Fig. 5 provides an overview of regions of multi-stability in the phase space u vs. s.D istinct regions with di erent numbers of stable steady states are identi ed depending on the combination of the dimensionless parameters. Lines of separation are determined by equations (22), (24) and, for the lower branch,



Figure 5.P hase space diagram u vs.s with $k_u = 1$, $k_r = 0$. The lines separating the distinct regions ofm ulti-stability are determined by equations (22), (24) and, for the lower branch, numerical results. In the lower left region only one stable xed point at (0;0) exists. In the region marked with \two stable xed points" one additional up-regulated stable xed point exists besides the one at (0;0). In \three stable xed points" region two additional up-regulated stable xed points exist. The dashed horizontal lines correspond to the sequences of nullclines in Figs. 2 and 3.

num erical results. The sequence of nullclines given in Fig. 2 is illustrated by the dashed line at u = 1, with the dots referring to the sub gures for varying s. The dashed line at u = 0.4 gives a similar representation, with its correspondence in Fig. 3.

Figs. 2 and 3 both indicate that the basin of attraction for the xed point at the origin (0;0) is separated from the basins of attraction of the up-regulated stable states by a set of unstable xed points. The sequences of graphs also illustrate that these unstable xed pointsm ove towards the xed point at (0;0) for increasing s, thus continously reducing the size of its basin of attraction. However, this size characterizes the stability of the xed point at the origin (0;0) in response to external perturbations. Unlike the interm ediate stable steady state, arising from the bifurcation at s₁ depicted in Fig. 4 (a), where a dynam ically increasing s inevitably leads to one of the two up-regulated xed points, the escape from the xed point (0;0) needs to be triggered by a perturbation that exceeds the size of its basin of attraction. G iven the position of the unstable xed point at the diagonal (x₃; x₃) as a function of s in equation (15), an appropriate measure for the size of the basin of attraction is provided.

3.2 A sym m etric system

As indicated by Zhang et al. (1999, 2000) the inhibition of PU 1 by GATA – 1 and the converse are based on di erent mechanisms. The formation of the PU 1-GATA –1 complex, which we refer to as a Z₂-complex, prevents free transcription factors from binding to their speci c DNA binding sites. A com – petitive inhibition in this form a ects both transcription factors, although Zhang et al. (2000) do not explicitly outline the consequences of binding of the PU 1-GATA –1 complex to the PU 1 binding site. On the other hand, GATA – 1 prevents the binding of c-Jun to the DNA bound PU 1 protein and thus disables the transcription initiation of PU 1. This process explicitly targets the PU 1 binding sites and introduces a functional asymmetry of inhibition mechanisms.

The mathematical counterpart of this asymmetry is a specic binding rate $K_6 > 0$ while keeping $K_7 = 0$ (see equations (8), (9)). In terms of the dimensionless formulation in equations (10) and (11) this translates into two dimensional constants $k_{r_x} > 0$ and $k_{r_y} = 0$. The additional binding mode ($k_{r_x} > 0$) can be interpreted as a reduction in the transcriptional activity of the X gene conferring a disadvantage relative to Y.

For any $k_{r_x} > 0$ there is a symmetry breaking which shifts the previously observed bifurcations of the diagonal and destroys the pitchfork bifurcation observed in the symmetric case for large u. The two up-regulated stable fields are not created instantaneously by the transformation of a previous stable state at the diagonal, but the initial stable point remains unchanged while a further (saddle node) bifurcation forms the second up-regulated stable point alongside with one unstable for the point. This scenario is shown in the sequence of nullclines in Fig. 6 (a)-(c). The parameter k_{r_x} regulates the distance between the up-regulated stable points and the extension of their basins of attraction. This is visualized in the bifurcation diagrams in Fig. 6 (d)-(f) for different values of k_{r_x} .

For sm allunspeci c transcription rates u, where in the symmetric case the additional up-regulated stable states are created of the diagonal, no qualitative changes are introduced by the functional asymmetry.

The introduction of asymmetry is not necessarily based on dimension of a erent interaction mechanisms. It is plausible that auto-regulative transcription activation does not require identical transcription rates for the genes of interest. This can be described by relaxing the symmetry assumption of Section 2.1, which leads to gene speci c transcription rates s_x and s_y . This asymmetry in transcriptional activity results in a qualitatively similar symmetry breaking as in the case of the mechanistic asymmetry: the pitchfork bifurcation occurring for large



Figure 6. The deform ation of the nullclines for the case $k_{r_x} = 0.1$ is depicted in gures (a) to (c). Parameters are $k_{r_y} = 0$, u = 0.8, $k_u = 1$, and s = 1.99 (a), s = 3.0 (b), s = 6.8 (c). The trivial xed point at (0;0) is marked by , the stable/unstable xed points o the diagonal by / . The qualitative change in the bifurcation behavior is shown in gures (d) to (f). The bifurcation diagram s are shown for increases in the asymmetry parameter k_{r_x} . Parameters are $k_{r_y} = 0$, u = 1, $k_u = 0.8$ and $k_{r_x} = 0.0$ (d), $k_{r_x} = 0.01$ (e), and $k_{r_x} = 0.1$ (f). Solid lines indicate stable, dashed lines unstable xed points.

u is replaced by a remaining stable state alongside a saddle-node bifurcation form ing the second up-regulated stable state (data not show n). The magnitude of the di erence in the speci c transcriptions rates s_x and s_y regulates the distance between the up-regulated stable states in the phase plane.

In a scenario where asymmetry of interaction mechanisms occurs alongside an asymmetry in the specic transcription rates, the elects on the system behavior combine, either amplifying or compensating each other.

3.3 Over-expression scenarios

Induced over-expression of a certain critical component is a common experimentalmethod to study interaction dynamics between dierent transcription

factors and has also been applied to the GATA-1 / PU 1 system (Nerlov et al., 2000; Rekhtm an et al., 1999; Zhang et al., 2000). These experiments provide insight in the stability of the system, interaction time scales, and the role of co-factors and interaction mechanisms. We have applied an over-expression in pulse of am plitude and duration doe to the model system given in equations (10), (11). Characteristics of the dynam ic response are only valid under the outlined steady state assumptions. A qualitative overview of the simulation results is presented in Fig. 7. Starting from a fully symmetric system as studied in Section 3.1 where, for large s, the system is in one of the two up-regulated states (characterized by one high and one low expressed transcription factor) two modes of over-expression are applied: a short impulse over-expression of the lower expressed component and a long and steady overexpression of the same component. Not surprisingly, the model reacts to the over-expression with two distinct scenarios, depending on the intensity of the in pulse. For a subcritical over-expression the system returns to the previous expression level (indicated in Figs. 7(a) and (d)), whereas for a supercritical situation the form er expression state is reversed (indicated in Figs 7 (b), (c), (e) and (f). Translating this picture into the x vs. y phase plane, the supercritical over-expression corresponds to a change from one basin of attraction to another, induced by a crossing of the separatrix. Most available experim ental techniques to articially induce gene expression lead to a massive over-expression that signi cantly exceeds physiological levels, a scenario still underestim ated by Figs. 7 (c) and (f). A sensitively tuned expression experim ent is more promising to elucidate critical intensities and time scales necessary to induce a perm anent shift in the genetic expression patterns and thus to characterize the stability of the initial states.

4 D iscussion

The presented model of transcription factor interaction is based on principles of coupled feedback regulations, which have previously been proposed for the description of general genetic switches (Becskei et al., 2001; C inquin and D em ongeot, 2005; Francois and H akim, 2004; G ardner et al., 2000; G lass and K au m an, 1973) and the modeling of prokaryotic gene regulation (M cA dam s and A rkin, 1998; Santillan and M ackey, 1998, 2001, 2004). Here, speci c experimental know ledge of activation and inhibition mechanisms of two transcription factors (G ATA -1 and PU .1), which play a key role in the myeloid/erythroid differentiation process of hem atopoietic progenitor cells, is incorporated in this general fram ework.

Our model analysis particularly focuses on the investigation of the steady states of transcription factor expression and there dependence on parameter changes. In this context, we are able to analyze the experimentally suggested



Figure 7. Scenarios for sub- and supercritical over-expression. In the subcritical scenarios (a) and (d) the transcription factor concentrations remain at the same xed point, whereas in the supercritical scenarios (b), (c), (e) and (f) the over-expression leads to a change of the basin of attraction, resulting in an di erent nal value of the transcription factor concentrations. O ver-expression is applied as a short term in uence at time t = 30 ((a), (b) and (c)) and long term in uence starting at time t = 20 ((d), (e) and (f)). Parameters are s = 5, u = 1, $k_u = 1$, and $k_r = 0$. The over-expression is applied with amplitude a_{oe} and duration $d_{oe} = 3$ and $d_{oe} = 1$ (a), $a_{oe} = 4$ and $d_{oe} = 1$ (b), $a_{oe} = 8$ and $d_{oe} = 1$ (c), $a_{oe} = 0.3$ and $d_{oe} = 50$ (d), $a_{oe} = 0.32$ and $d_{oe} = 50$ (e), $a_{oe} = 2.5$ and $d_{oe} = 50$ (f).

feedback structures and their e ects on the system behavior under various conditions.

To facilitate the m athem atical analysis, a num ber of sim pli cations have been m ade. W e interpret the transcription factors described in the m odel (X and Y) as representatives of a m ore com plex factor form ation rather than an explicit m odel of PU 1 and GATA-1 abne. Also, we are aware that m ost of the statem ents resulting from the m odel analysis are only sem i-quantitative in the sense that for all m odel parameters, as there are DNA binding-, decay-, and transcription-rates, no experimentally determ ined estimates are available for the investigated system. In the same line of argum entation, details of the transcription factor m olecules and the delay induced by the processes of transcription and translation, have been excluded from the analysis. A lthough such phenom ena

can in uence the dynam ics of the system (Bundschuh et al., 2003; V ilar et al., 2002), these e ects are speculative since detailed inform ation about relevant rates and time scales are not available. The simplications arising from the quasi steady state assumption outlined in Section 2 for dimerization and DNA binding impose further limitations on our model with respect to the exact description of the system dynamics (c.f. P irone and E lston, 2004). However, these simplications do not e ect the steady state behavior, and, thus, do not alter the results derived in Section 3.

The functional role of the so called priming behavior is a question of particular biological relevance which is addressed by this model. It has been suggested that low level co-expression of multiple transcription factors, speci c for di erent lineages, m ight be a characteristic of (hem atopoietic) tissue stem cells (Akashi, 2005; Cross and Enver, 1997; Orkin, 2000). However, it is currently unclear whether priming corresponds to a stable state of low level co-expression or to a truly zero-expression overlaid by some random expression noise. Furtherm ore, there is a hypothesis that lineage speci cation induction m ight be a two stage process with a primary initialization of transcription factor network interaction (i.e., a transition from no expression to low level co-expression) and a secondary network-induced di erentiation process (Enver and G reaves, 1998). This perspective immediately leads to the questions under which conditions such a two stage process can be established and whether such a sequence of di erent activation states of the transcription factor network requires (multiple) external induction signals or whether it represents a system inherent developm ent.

The suggested model generates two characteristic modes of system stability depending on the magnitude of the speci c transcription rate s: For small s only the trivial xed point (0;0) exists; for large s two additionalup-regulated stable states are observed that are marked by the dom inance of one factor over the other (dom inated co-expression). These modes are maintained independently of a mechanistic or parametric asymmetry. A ssum ing a di erentiation initiation by increasing the transcription rate s (e.g. by changes in chromatin structure (Berger and Felsenfeld, 2001; R osmarin et al., 2005) or by alterations in activation/inhibition com plexes (Hume, 2000)), the transition between the di erent stable states is the centralmechanism characterizing lineage speci cation.

W ithin the proposed biological fram ework the trivial xed point at (0;0), which exists for all values of s, can be idential with the undimentiated state of a cell where neither activation nor decision processes are observed. It should be mentioned that stability of this xed point is specific for the outlined model and has not been observed for the general case of a toggle switch (c.f. G ardner et al., 2000). In logical extension, the two up-regulated stable xed points, observed for large s, would be interpreted as expression states promoting one or the other lineage. These distinct states are characterized by a high auto-regulative expression of one dom inating factor and a reduced expression of the antagonistic factor.

In Sections 3.1 and 3.2 it has been dem onstrated that, for increased unspeci c transcription rates u (but only in this case!), a further stable xed point exists for interm ediate s prior to the form ation of the two distinct states of dom inated co-expression. This particular xed point is characterized by a balanced low level co-expression of the two antagonistic factors where no nal com m it-m ent decision has been m ade. The resulting transition sequence between three distinct regions of multi-stability can be interpreted as a possible explanation for a two stage di erentiation process mentioned above.

The induction of a system change from the stable trivial xed point to the dom inated or, if existent, balanced low level co-expression state, needs to be triggered either by a stochastic background expression or by an active in pulse on the system. The unstable xed point separating the trivial from the up-regulated stable states is an indicator of the size of the basins of attraction. The observation that the unstable xed point approaches the trivial one for increasing s indicates that the magnitude of the perturbation to introduce a transition from the zero-state to the co-expression states decreases in the same fashion: for a su ciently large s even a small perturbation is able to initiate di erentiation.

Concluding from these results, there are two di erent scenarios to explain the experimentally suggested priming behavior within the proposed model fram ework: (1) Prim ing m ight be considered as the existence of perturbations in the expression of transcription factors, in posed on a zero-expression state represented by the trivial xed point at (0;0), either in the form of stochastic background uctuations (functional noise) or by active in pulses. In this seenario, the perturbations are necessary components of the regulatory system to induce a di erentiation process. It points to the potential role of stochastic e ects in the context of decision making in stem cell di erentiation as frequently suggested (see K aem et al. (2005) for a review). (2) In contrast to this scenario, priming can also be explained by the balanced low level coexpression state, which becomes unstable for increasing speci c transcription rates. Due to this parameter dependent loss of stability, this scenario would lead to di erentiation without the need for external perturbations². How ever, the balanced low level co-expression state is only existent if there is a certain degree of unspeci c transcription.

Currently, our results do not allow to decide between the two scenarios. The

 $^{^2}$ To be precise: An in nitesimal perturbation is required to escape from the unstable xed point. Fluctuations of this magnitude are present in any \realworld" system .

introduction of arti cial di erentiation in pulses of di erent intensities on uncom m itted cells m ight be an appropriate way to tackle this question experimentally. W hereas, a low level co-expression priming (like in scenario (2)) would be una ected by these perturbations, the system could be enforced to escape the priming status in scenario (1). Moreover the existence and the stability of the di erent stable system states depend sensitively on the model parameters. Due to the lack of available data on transcription and binding rates, we are currently not able to specify the biological relevant regimes more rigorously. A ny experimental approximation of binding and transcription rates for the involved components supports the identication of the nature of priming.

The over-expression scenarios presented in Section 3.3 fail to explain experimental ndings described by several authors (Nerby et al, 2000; Rekhtman et al, 2003; Zhang et al., 2000). In spite of the induced up-regulation of one transcription factor it was observed that the transcription level of the antagonistic transcription factor remained more or less constant. These observations are in contrast to the model results presented here, in which the induced over-expression of the initially low expressed factor shifts the equilibrium to the opposing co-expression state. Retaining our model assumptions, a potential interpretation can be given as follows: One of the major functions of transcriptional regulators like GATA-1 and PU 1 is the activation of a set of lineage-specic genes which include further transcription and growth factors as well as functional components of the committed lineages (Tenen, 2003). In Sieweke and Graf (1998) and Tsaietal. (1991), the authors point to a continuously modulated set of cooperative lineage-inherent transcription factors changing with the state of di erentiation. Such secondary com plexes of transcription factors could in turn act as activators of the initial transcription factor, substituting for a simple auto-regulation and thus stabilizing the initial up-regulation pattern (Hume, 2000). In such a scenario our model would only account for the initial switching process. The experimentally observed stable transcription level of the antagonistic factor in over-expression experiments could be interpreted as a substitution of the auto-regulation by secondary transcription factor com plexes.

Sum marizing, the presented model is able to provide a quantitative explanation for possible mechanisms underlying lineage speci cation control in eukaryotic systems. It is able to generate parameter dependent changes in the system behavior, with alteration of the number of possible stable steady states. Speci cally, the model explains states of stable co-expression as well as the situation characterized by an over-expression of one factor over the other. The conditions inducing shifts from one to another stable state (e.g. parameter choice, degree of system disturbances), how ever, depend in a sensitive manner on the assumed activation and inhibition mechanisms. U sing the mathematical model, we were able to test several combinations of experimentally described feedback mechanisms with respect to their in uence on the resulting stable states and provide possible explanations for the experimentally suggested differentiation priming of stem cells.

A cknow ledgm ent

The authors thank M ichael M ackey for his encouragement and critical discussions in the process of preparing this manuscript, and M ichael C ross for his explanations of many biological details. Furthermore we acknow ledge the H um an Frontier Science P rogram which supported initiation of this work by short-term fellow ship to IR. at the C entre for N onlinear D ynam ics, M cG ill U niversity, M ontreal.

A Derivation of transcription factor dynam ics

It is assumed that the transcription of transcription factors X and Y requires the existence of activator complexes, i.e., the binding of X or Y dimers to the promoter regions of X (D_x) and Y (D_y) , respectively. As described in Section 2.2, we distinguish between a speci c (see equations (1),(3)) and an unspeci c (equations (2),(4)) transcription activation. Furtherm ore, there is the possibility that X and Y can act jointly as a repressor dimer Z_1 , inhibiting the DNA binding of the X and Y activator dimers (see equations (6),(7)).

The totalam ount of prom oter sites for X and Y can be specified as the sum of unbound (free) and occupied (by repressor or activator molecules) promoter regions, i.e.,

$$D_{x=y}^{tot} = D_{x=y} + D_{x=y}^{xy} + D_{x=y}^{xx} + D_{x=y}^{yy}$$
 : (A.1)

U sing the equilibrium (dissociation) constants

$$K_{1} = \frac{D_{x}^{xx}}{D_{x}x^{2}}; \quad K_{2} = \frac{D_{x}^{yy}}{D_{x}y^{2}}; \quad K_{3} = \frac{D_{y}^{yy}}{D_{y}y^{2}}; \quad K_{4} = \frac{D_{y}^{xx}}{D_{y}x^{2}}; \quad K_{6} = \frac{D_{x}^{xy}}{D_{x}xy}; \text{ and } K_{7} = \frac{D_{y}^{xy}}{D_{y}xy}; \quad (A.2)$$

obtained from assuming equations (1)-(4), (6), (7) to be in a quasi-steady state, the fraction of promoter sites contributing to active X and Y transcription is

given by

$$\frac{D_{x}^{xx} + D_{x}^{yy}}{D_{x}^{tot}} = \frac{K_{1}x^{2}D_{x} + K_{2}y^{2}D_{x}}{D_{x} + K_{1}x^{2}D_{x} + K_{2}y^{2}D_{x} + K_{6}xyD_{x}}$$

$$= \frac{K_{1}x^{2} + K_{2}y^{2}}{1 + K_{1}x^{2} + K_{2}y^{2} + K_{6}xy}$$
(A.3)

and

$$\frac{D_{y}^{yy} + D_{y}^{xx}}{D_{y}^{tot}} = \frac{K_{3}y^{2}D_{y} + K_{4}x^{2}D_{y}}{D_{y} + K_{3}y^{2}D_{y} + K_{4}x^{2}D_{y} + K_{7}xyD_{y}}$$

$$= \frac{K_{3}y^{2} + K_{4}x^{2}}{1 + K_{3}y^{2} + K_{4}x^{2} + K_{7}xy}$$
(A.4)

respectively.

Taking the (rst order) decay rates of X and Y into account, one immediately obtains equations (8), (9) by writing down the balance equations for X and Y.

B Domain of the nullclines

Under the equilibrium assumption equation (12) can be solved for:

$$y_{1=2}(x) = \frac{k_{r}x^{2}}{k_{r}^{2}x^{4}} \frac{4k_{u}x(u-x)(sx-1-x^{2})}{2k_{u}(u-x)}$$
(B.1)

which describes a set of nullclines. There is an obvious singularity at x = u.

The solutions $y_{1=2}(x)$ are real for $0 < h(x) = k_r^2 x^4 - 4k_u x (u - x) (sx - 1 - x^2)$ with h(x) defined as the expression under the square root in the previous equation. For $k_r = 0$ the roots of h(x) are located at

$$x_1^h = 0$$
 (B.2)

$$x_2^h = \frac{s}{2} \qquad \frac{s^2}{4} \qquad 1$$
 (B.3)

$$x_3^h = u$$
 (B.4)

$$\mathbf{x}_{4}^{h} = \frac{s}{2} + \frac{s^{2}}{4} + 1$$
 (B.5)

Real roots at $x_{2=4}^{h}$ exist only for s 2. Fig. B.1 shows the function h(x) for s = 1.9 and s = 2.1. In the case s < 2 the parameter $u = x_{3}^{h}$ restricts the de nition space of the nullclines to $x \ge [0; x_{3}^{h}]$. For s > 2 three scenarios exist,



Figure B.1. The function $h(x) = k_r^2 x^4 - 4k_u x (u - x) (sx - 1 - x^2)$ is shown for $k_r = 0$, u = 1, $k_u = 1$ and s = 1.9 (a) and s = 2.1 (b).

where the singularity at $x = u = x_3^h$ m arks a boundary for distinct intervals in the dom ain: $x_3^h < x_2^h$ (with x 2 [0; x_3^h] [[k_2^h ; x_4^h]), $x_2^h < x_3^h < x_4^h$ (with x 2 [0; x_2^h][[k_3^h ; x_4^h]) shown in Fig.B 1 (b), and $x_4^h < x_3^h$ (with x 2 [0; x_2^h][[k_4^h ; x_3^h]).

C Derivation of bifurcation condition

The nullclines of the symmetric system derived in (12) and (13) are interpreted as functions of x and y:

$$\mathbf{x} = f(\mathbf{x}; \mathbf{y}) = \frac{\mathbf{s}\mathbf{x}^2 + \mathbf{u}\mathbf{k}_{\mathbf{u}}\mathbf{y}^2}{1 + \mathbf{x}^2 + \mathbf{k}_{\mathbf{u}}\mathbf{y}^2 + \mathbf{k}_{\mathbf{r}}\mathbf{x}\mathbf{y}}; \quad (C.1)$$

$$y = g(x;y) = \frac{sy^2 + uk_u x^2}{1 + k_u x^2 + y^2 + k_r xy}$$
 (C 2)

To derive bifurcation conditions one has to determ ine the point of tangency of the nullclines f(x;y), g(x;y) at a steady state (x;y), ie.

$$\frac{\mathrm{df}(\mathbf{x};\mathbf{y})}{\mathrm{dy}} = \frac{\mathrm{dg}(\mathbf{x};\mathbf{y})}{\mathrm{dx}} (\mathbf{x}_{\mathbf{y}}) = \frac{\mathrm{dg}(\mathbf{x};\mathbf{y})}{\mathrm{dx}} (\mathbf{x}_{\mathbf{y}})$$
(C.3)

Generally, it holds for inverse functions h and $k = h^{1}$ that $k^{0}(h(x)) = (h^{0}(x))^{1}$. Considering only points at the diagonal x = h(x) = y, it follows that $h^{0}(x) = (k^{0}(x))^{1}$. A sum ing identity of the rst order derivatives h^{0} and k^{0} at some point x on the diagonal yields, therefore, $(h^{0}(x))^{2} = (k^{0}(x))^{2} = 1$.

From these statem ents, it follows that we have to consider the following equal-

ities to nd the bifurcation conditions for the symmetric system, restricting to symmetric steady states of the form (x;x):

$$\frac{\mathrm{df}(\mathbf{x};\mathbf{y})}{\mathrm{dx}} = \frac{\mathrm{dg}(\mathbf{x};\mathbf{y})}{\mathrm{dx}} = \mathrm{jl}; \qquad (C.4)$$

References

- Akashi, K., 2005. Lineage promiscuity and plasticity in hem atopoietic development. Ann N Y Acad Sci 1044, 125{31.
- A kashi, K., He, X., Chen, J., Iwasaki, H., Niu, C., Steenhard, B., Zhang, J., Haug, J., Li, L., 2003. Transcriptional accessibility for genes of multiple tissues and hem atopoietic lineages is hierarchically controlled during early hem atopoiesis. B lood 101 (2), 383{9.
- Becskei, A., Seraphin, B., Serrano, L., 2001. Positive feedback in eukaryotic gene networks: cell di erentiation by graded to binary response conversion. EM BO J 20 (10), 2528{35.
- Berger, S., Felsenfeld, G., 2001. Chrom atin goes global. MolCell8 (2), 263{8.
- Bundschuh, R., Hayot, F., Jayaprakash, C., 2003. The role of dimerization in noise reduction of simple genetic networks. J. Theor. Biol. 220 (2), 261 {269.
- Cantor, A.B., Orkin, S.H., 2002. Transcriptional regulation of erythropoiesis: an a air involving multiple partners. Oncogene 21 (21), 3368{76.
- Chen, H., Ray-gallet, D., Zhang, P., Hetherington, C. J., Gonzalez, D. A., Zhang, D. E., Moreau-gachelin, F., Tenen, D. G., 1995. PU.1 (Spi-1) autoregulates its expression in myeloid œlls. Oncogene 11 (8), 1549{60.
- C inquin, O., Demongeot, J., 2002. Positive and negative feedback: striking a balance between necessary antagonists. J Theor Biol. 216 (2), 229{41.
- C inquin, O., Demongeot, J., 2005. High-dimensional switches and the modelling of cellular di erentiation. J Theor B iol 233 (3), 391 {411.
- Cross, M.A., Enver, T., 1997. The lineage commitment of haem opoietic progenitor cells. Curr Opin Genet Dev 7 (5), 609{13.
- Cross, M. A., Heyworth, C. M., Murrell, A. M., Bockamp, E. O., Dexter, T. M., Green, A. R., 1994. Expression of lineage restricted transcription factors precedes lineage speci c di erentiation in a multipotent haem opoietic progenitor cell line. Oncogene 9 (10), 3013{16.
- Du, J., Stankiewicz, M. J., Liu, Y., Xi, Q., Schmitz, J. E., Lekstrom H in es, J. A., A ckerm an, S. J., 2002. Novel combinatorial interactions of GATA -1, PU.1, and C/EBP epsilon isoform s regulate transcription of the gene encoding eosinophil granule major basic protein. J B iol Chem 277 (45), 43481 [94.
- Enver, T., Greaves, M., 1998. Loops, lineage, and leukem ia. Cell 4 (1), 9{12.
- Francois, P., Hakim, V., 2004. Design of genetic networks with speci ed functions by evolution in silico. Proc NatlA cad SciU SA 101 (2), 580{5.

- Gardner, T.S., Cantor, C.R., Collins, J.J., 2000. Construction of a genetic toggle switch in Escherichia coli. Nature 403 (6767), 339{42.
- G lass, L., K au man, S., 1973. The logical analysis of continuous, non-linear biochem ical control networks. J Theor Biol 39 (1), 103{29.
- Hu, M., Krause, D., Greaves, M., Sharkis, S., Dexter, M., Heyworth, C., Enver, T., 1997. Multilineage gene expression precedes commitment in the hem opoietic system. Genes Dev 11 (6), 774 [85.
- Hume, D., 2000. Probability in transcriptional regulation and its im plications for leukocyte di erentiation and inducible gene expression. Blood 96 (7), 2323{8.
- Kaem, M., Elston, T., Blake, W., Collins, J., 2005. Stochasticity in gene expression: from theories to phenotypes. Nat Rev Genet 6 (6), 451{64.
- Loe er, M., Roeder, I., 2002. Tissue stem cells: De nition, plasticity, heterogeneity, self-organization and models -a conceptual approach. Cells Tissues Organs 171 (1), 8{26.
- Lord, B. I., 1997. Biology of the haem opoietic stem cell. A cadem ic Press, C am bridge, pp. 401{422.
- M cA dam s, H ., A rkin, A ., 1998. Simulation of prokaryotic genetic circuits. A nnu R ev B iophys B iom ol Struct 27, 199{224.
- Nerby, C., Querfurth, E., Kulessa, H., Graf, T., 2000.GATA-1 interacts with the myeloid PU.1 transcription factor and represses PU.1-dependent transcription.Blood 95 (8), 2543{51.
- N ishim ura, S., Takahashi, S., Kuroha, T., Suwabe, N., Nagasawa, T., Trainor, C., Yam am oto, M., 2000. A GATA box in the GATA-1 gene hem atopoietic enhancer is a critical element in the network of GATA factors and sites that regulate this gene. Mol Cell Biol 20 (2), 713 {23.
- O ikawa, T., Yamada, T., Kihara-Negishi, F., Yamamoto, H., Kondoh, N., Hitomi, Y., Hashimoto, Y., 1999. The role of Ets family transcription factor PU.1 in hem atopoietic cell di erentiation, proliferation and apoptosis. Cell Death Dier 6 (7), 599{608.
- O kuno, Y., Huang, G., Rosenbauer, F., Evans, E. K., Radom ska, H. S., Iwasaki, H., Akashi, K., M oreau-G achelin, F., Li, Y., Zhang, P., Gottgens, B., Tenen, D.G., 2005. Potential autoregulation of transcription factor PU.1 by an upstream regulatory element. MolCellBiol25 (7), 2832 [45.
- Orkin, S.H., 1995. Hem atopoiesis: how does it happen? Curr Opin Cell Biol 7 (6), 870{7.
- Orkin, S. H., 2000. Diversi cation of haem atopoietic stem cells to speci c lineages. Nat Rev Genet 1 (1), 57{64.
- P irone, J., E lston, T., 2004. F luctuations in transcription factor binding can explain the graded and binary responses observed in inducible gene expression. J Theor B iol 226 (1), 111{21.
- Potten, C.S., Loe er, M., 1990. Stem cells: attributes, cycles, spirals, pitfalls and uncertainties. lessons for and from the crypt. Development 110 (4), 1001{1020.
- Rekhtman, N., Choe, K.S., Matushansky, I., Murray, S., Stopka, T., Skoultchi,

A.I., 2003.PU.1 and pRB interact and cooperate to repress GATA-1 and block erythroid di erentiation.MolCellBiol23 (21), 7460{74.

- Rekhtman, N., Radparvar, F., Evans, T., Skoultchi, A. I., 1999. Direct interaction of hem atopoietic transcription factors PU.1 and GATA-1: functional antagonism in erythroid cells. Genes Dev 13 (11), 1398{411.
- Rosmarin, A., Yang, Z., Resendes, K., 2005. Transcriptional regulation in myelopoiesis: Hem atopoietic fate choice, myeloid di erentiation, and leukem ogenesis. Exp Hem atol 33 (2), 131{43.
- Santillan, M., M. ackey, M. C., 1998. Dynam ic behavior in m. athem aticalm odels of the tryptophan operon. Chaos 11, 261 {8.
- Santillan, M., Mackey, M. C., 2001. Dynamic regulation of the tryptophan operon: A modeling study and comparison with experimental data. Proc NatlAcad SciU S A 98, 1364{9.
- Santillan, M., Mackey, M.C., 2004. Why the lysogenic state of phage lambda is so stable: a mathematical modeling approach. Biophys J 86 (1), 75{84.
- Sieweke, M.H., Graf, T., 1998. A transcription factor party during blood cell di erentiation. Curr Opin Genet Dev 8 (5), 545{51.
- Tenen, D.G., 2003. Disruption of di erentiation in hum an cancer: AM L shows the way. Nat Rev Cancer 3 (2), 89{101.
- T sai, S.F., Strauss, E., Orkin, S.H., 1991. Functional analysis and in vivo footprinting implicate the erythroid transcription factor GATA-1 as a positive regulator of its own promoter. Genes Dev 5 (6), 919{31.
- Vilar, J., Kueh, H., Barkai, N., Leibler, S., 2002. Mechanisms of noiseresistance in genetic oscillators. Proc NatlA cad SciU SA 99 (9), 5988 {92.
- Voso, M. T., Burn, T. C., Wulf, G., Lin, B., Leone, G., Tenen, D. G., 1994. Inhibition of hem atopoiesis by competitive binding of transcription factor PU.1. Proc NatlA cad SciU S A 91 (17), 7932 {36.
- Yam ada, T., Kihara-Negishi, F., Yam amoto, H., Yam amoto, M., Hashimoto, Y., Oikawa, T., 1998. Reduction of DNA binding activity of the GATA-1 transcription factor in the apoptotic process induced by overexpression of PU.1 in murine erythroleukem is cells. Exp Cell Res 245 (1), 186{94.
- Zhang, P., Behre, G., Pan, J., Iwama, A., Wara-Aswapati, N., Radomska, H.S., Auron, P.E., Tenen, D.G., Sun, Z., 1999. Negative cross-talk between hem atopoietic regulators: GATA proteins repress PU 1. Proc NatlA cad Sci U S A 96 (15), 8705{10.
- Zhang, P., Zhang, X., Iwama, A., Yu, C., Smith, K.A., Mueller, B.U., Narravula, S., Torbett, B.E., Orkin, S.H., Tenen, D.G., 2000.PU.l inhibits GATA-1 function and erythroid di erentiation by blocking GATA-1 DNA binding. Blood 96 (8), 2641{8.