Optim alD ynam icalR ange of Excitable N etworks at C riticality

O sam e Kinouchi

Departamento de F sica e Matematica, Faculdade de Filoso a, Ciências e Letras de Ribeirao Preto, Universidade de Sao Paulo, Av. dos Bandeirantes 3900, 14040-901, Ribeirao Preto, SP, Brazil

Mauro Copelli

Laboratorio de F sica Teorica e Computacional, Departamento de F sica, Universidade Federal de Pernambuco, 50670-901 Recife, PE, Brazil

A recurrent idea in the study of com plex systems is that optimal information processing is to be found near bifurcation points or phase transitions. However, this heuristic hypothesis has few (if any) concrete realizations where a standard and biologically relevant quantity is optim ized at criticality. Here we give a clear example of such a phenomenon: a network of excitable elements has its sensitivity and dynam ic range maxim ized at the critical point of a non-equilibrium phase transition. Our results are compatible with the essential role of gap junctions in olfactory glom enuli and retinal ganglionar cell output. Synchronization and global oscillations also appear in the network dynam ics. We propose that the main functional role of electrical coupling is to provide an enhancement of dynam ic range, therefore allow ing the coding of information spanning several orders of magnitude. The mechanism could provide a microscopic neural basis for psychophysical laws.

PACS num bers: 87.18 Sn, 87.19 La, 87.10 + e, 05.45.-a, 05.40.-a K eywords: G ap junction, Ephaptic interaction, O lfaction, R etina, Excitable m edia, N eural code, D ynam ic range, C ellular autom ata

P sychophysics is probably the rst experimental area in neuropsychology, having been founded by physiologists (Helm holtz and Weber) and physicists (Fechner, Plateau, Maxwell and Mach) in the middle of the nineteenth century [1]. Its aim is to study how physical stimuli transduce into psychological sensation, probably the most basic mind-brain problem. How to relate such high-level psychological phenomena to low-level neurophysiology is a task for the twenty-rst century brain sciences.

As the intensity of physical stimuli (light, sound, pressure, odorant concentration etc) varies by several orders of m agnitude, psychophysical laws must have a large dynam ical range. Although this has been extensively veried experimentally at the psychological [1] and neural [2, 3, 4] level, little work has been done regarding the mechanism that produces such psychophysical laws [5, 6, 7, 8]. However, two main ideas seem to be consensual: (1) nonlinear transduction must be done at the sensory periphery to prevent early saturation [1]; and (2) the broad dynam ic range is offen a collective phenom enon [1, 5, 6], because single cells usually respond in a linear saturating way with sm all ranges [9, 10].

Two nonlinear transfer functions have been widely used to texperimental data, both in psychophysics and in neural response: the bigarithm function F (S) = C bigS (W eber-Fechner law) and the power law function F (S) = C S^m (Stevens Law), where S is the stimulus level, C is a constant and m is the Stevens exponent. Later, other functions have been proposed to t data with more extended input range, and to account for sensory saturation, in particular the H ill function F (S) = $F_{m ax}S^m = (S^m + S_0^m)$ where $F_{m ax}$ is the saturation response and S_0 is the input level for half-m axim um response. Notice that, because both H ill and Stevens functions have a power law regime, it is natural to denote the exponents by the same parameter m.

Some authors have tried to derive such phenom enological laws from the structure of natural signals [11]. This type of work m ay furnish an evolutionary motivation for biological organism s to implement or approximate such laws. How ever, there is no consensual view on this theoretical aspect, and it does not provide a neural basis for implementing the psychophysical laws.

In contrast, our statistical physics approach to neural psychophysics [6, 7, 8, 12] shows how H ill-like transfer functions m ay arise in biological excitable systems: they are not put into the m odels by hand, nor m athem atically derived from a priori reasoning, but appear as a cooperative e ect in a network of excitable elements. At di erente biological levels, these elements m ay be interpreted as whole neurons, excitable dendrites, axons, or other subcellular excitable units. W e show that a network of excitable elements, each with sm all dynam ic range, presents a collective response with broad dynam ic range and high sensitivity. Even more interesting, we nd that the dynam ic range is m axim ized if the spontaneous activity of the network corresponds to a critical process. This is compatible with recent ndings of critical avalanches in in vitro neural networks [13, 14] and provides a clearcut exam ple of optim al inform ation processing at criticality [15, 16, 17]. The m odel also has other dynam ical features and permits us to m ake

a testable prediction.

In previous work we have introduced the idea that excitable waves in active media provide a mechanism for strong nonlinear ampli cation with large dynamic range. This has been shown in simulations with one-dimensional [6] and two-dimensional [7] deterministic cellular autom atom models, as well as with one-dimensional networks of coupled m aps and Hodgkin-Huxley elements [8]. A nalytical results have recently been obtained for the one-dimensional cellular autom atom models are model of the one-dimensional cellular autom atom models are the activity propagation is stochastic and the electrical synapses form a random network. This latter case seems to be a more realistic topology for, say, the olfactory intraglomerular network of excitable dendrites coupled by gap junctions [18, 19, 20]. We nd a whole new phenomenology related to a non-equilibrium phase transition to re-entrant activity present in this model.

In the present model, each excitable element i = 1; :::; N has n states: $s_i = 0$ is the resting state, $s_i = 1$ corresponds to excitation and the remaining $s_i = 2; :::; n$ 1 are refractory states. There are two ways for the ith element to go from state $s_i = 0$ to 1: (1) owing to an external stimulus, modelled here by a Poisson process with rate r (which implies a transition with probability = 1 exp(rt) per time step); (2) with probability p_{ij} , owing to a neighbour j being in the excited state in the previous time step. Time is discrete (we assume t = 1 m s) and the dynamics, after excitation, is deterministic: if $s_i = 1$, then in the next time step its state changes to $s_i = 2$ and so on until the state $s_i = n$ 1 leads to the $s_i = 0$ resting state, so the element is a cyclic cellular autom aton [21]. The Poisson rate r will be assumed to be proportional to the stimulus level S (for example, the odorant concentration in olfactory processing). Notice that each element receives external signals independently, that is, we have a Poisson process for each element (m odelling the arrival of axonal inputs from di erent receivors).

The network with N elements is an Erdøs-Renyi undirected random graph, with N K =2 links being assigned to random ly chosen pairs of elements. This produces an average connectivity K where each element i (i = 1;:::;N) is random ly connected to K_i neighbours. The distribution P (K_i) of neighbours is a binom ial distribution with average K (see Fig 1A). The probability that an active neighbour j excites element i is given by p_{ij} , a random variable with uniform distribution in the interval $[0;p_{m ax}]$. The weights are symmetrical $(p_{ij} = p_{ji})$ and are kept xed throughout each simulation (\quenched disorder"). This kind of coupling models electric gap junctions instead of chemical synapses because it is fast and bi-directional. However, symmetry is not a necessary ingredient, because sim ilar results are obtained in asymmetrical networks. Note that we are not assuming that gap junctions have a stochastic dynamics, but only that, because other internal factors and noise are also present, a probabilistic account of the excitation process may be more realistic.

The local branching ratio $j = \begin{bmatrix} r & K \\ i & p \end{bmatrix}$ corresponds to the average num ber of excitations created in the next time step by the j-th element [14]. The distribution P ($_j$) of local branching ratios (a bell-shaped distribution with average = h $_i$ i) is shown in Fig. 1A. The average branching ratio is the relevant control parameter. In the simulations, we set by choosing $p_{max} = 2 = K$ and keeping < K = 2.

The network instantaneous activity is the density t of active (s = 1) sites at a given time t. We also de ne the average activity $F = T^{1} P_{t=1}^{T} t$ where T is a large time window (of the order of 10^{3} time steps). Typical t curves in the absence of stimulus (r = 0) are shown in Fig.1B.N otwithstanding the large variance of P ($_{j}$), only supercritical networks (that is, with > c = 1) have self-sustained activity, F > 0. As expected, critical networks have a larger variance in the distribution of extinction times and present a power law behaviour in the distribution of avalanche sizes with exponent 3=2 (not shown), in agreement with notings in biological networks [13].

Two types of oscillations are observed in this system. Under su ciently strong stimulation, all networks present transient collective oscillations, with frequencies of the order of the inverse refractory period (Fig. 1C). They are a simple consequence of the excitable dynam ics and the sudden synchronous activation by stimulus initiation. This transient behaviour is rem iniscent of oscillations widely observed in experiments [2]. Networks with $>_{\rm osc} >_{\rm c}$ also present self-sustained oscillations in the absence of stimulus (Fig. 1D), where $_{\rm osc}$ is a bifurcation threshold. The frequency depends on the network parameters, but remain in the gam marange 30 60 Hz. The oscillations are similar to reentrant activity found in spatially extended models of electrically coupled networks [23], from which analytical techniques for calculating the frequency could perhaps be borrowed [24]. We remark that the large time window used in the denition of F is only chosen for convenience, as it can be seen from Figs. 1C, D and 200 m s is su cient to give reliable averages.

As a function of the stimulus intensity r, networks have a minimum response F_0 (= 0 for the subcritical and critical cases) and a maximum response F_{max} . We dene the dynamic range = 10 log ($r_{0:9}=r_{0:1}$) as the stimulus interval (measured in dB) where variations in r can be robustly coded by variations in F, discarding stimuli which are too weak to be distinguished from F_0 or too close to saturation. The range $[r_{0:1};r_{0:9}]$ is found from its corresponding response interval $F_{0:1};F_{0:9}$, where $F_x = F_0 + x(F_{max} - F_0)$ (see Fig. 2C). This choice of a 10% -90% interval is




FIG.1: Network characterization and density of active sites. Simulations with $N = 10^5$ sites, K = 10 and n = 10 states. (A) Probability density function of local branching ratio and (inset) connectivity (\degree") distribution; (B) Instantaneous density of active sites for subcritical (black), critical (red) and supercritical (blue) branching parameters as functions of time (three di erent runs for each case); (C), (D) Instantaneous density (of all sites; upper panels) and raster plot (of 10^3 random ly chosen sites; low er panels) in response to a square pulse of stimulus ($r = 0.5 \text{ m s}^1$ for 100 m s t 300 m s, null otherwise) for critical (C) and supercritical (D) branching parameters.

arbitrary, but is standard in the literature and does not a ect our results.

As can be seen in Figs. 2A and 2B, the response curves F (r) of the networks present a strong enhancement of dynamic range compared with the uncoupled case = 0. In the subcritical regime, sensitivity is enlarged because weak stimuli are amplied due to activity propagation among neighbours. As a result, the dynamic range () increases monotonically with . In the supercritical regime, the spontaneous activity F₀ masks the presence of weak stimuli, therefore () decreases. The optimal regime e occurs precisely at the critical point (see Fig. 2D). This is a new and important result, because it is perhaps the rst clear example of signal processing optimization at a phase transition, making use of a standard and easily measurable perform ance index.

The curves F (r) could be tted by a Hill function, but are not exactly Hill. The theoretical curves in Fig.2 are obtained from a simple mean-ekd calculation (see below) that provides a very good t to the simulation data without free parameters, and correctly predicts the exponents governing the low-stimulus response F / r^m . An important point is that the Stevens-Hill exponent m changes from m = 1 in the subcritical regime to m = 0.5 at criticality. If we assume that biological networks work in the optimal regime, the critical value m = 0.5 suggests how exponents less than one could emerge in psychophysics [1] and neural responses [2, 4]. We note that apparent exponents between 0.5 and 1.0 are observed [12] if nite size e ects are present, that is, if N is small.

In the simple mean-eld approximation, we take K $_{i}$ = K and p_{ij} as the average value =K. The probability p_{t} that an inactive site at time t will be activated in the next time step by at least one of its K neighbours (a fraction




FIG. 2: Response curves and dynam ic range. Points represent simulation results with $N = 10^5$ sites, K = 10, n = 5 states and $T = 10^3$ m s, whereas lines correspond to the mean-eld model described in the text. (A) Response curves (mean ring rate vs. stimulus rate) from = 0 to = 2 (in intervals of 0.2). Line segments are power laws F / r^m with m = 1 (subcritical) and m = 1=2 (critical). Inset: spontaneous activity F_0 vs. branching ratio . (B) The same as in (A), but with a linear vertical scale. (C) Response curve for = 1.2 and relevant parameters for calculating the dynamic range . (D) Dynamic range vs. branching ratio is optimized at the critical point = 1.

 F_t of which is currently active) is simply $p_t = 1$ (1 $F_t = K$)^K. This leads to the following mean-eld map:

$$F_{t+1} = P_t(0) + P_t(0) (1) p_t;$$
(1)

where

$$P_t(0) = 1$$
 (n 1) F_t (2)

is the approximate probability of nding a site in the resting state and $(r) = 1 \exp(r t)$. The rst term on the right-hand side of equation 1 corresponds to activation due to external input, and the second term corresponds to activation due to neighbour propagation. In the stationary regime, the response function F (r) is given by the solution of

$$F = (1 (n 1)F) 1 (1 F = K)^{K} (1 (r)) :$$
(3)

Note that equation 2 is only exact in the stationary state, but the resulting map is consistent because the result of its iterations coincides with the solution of equation 3 (in particular, F = 1=n, so that P(0) = 0).

As is usual in the statistical physics of phase transitions [21], we analyse two limits: the critical behavior without an external eld and the eld containing eld at the critical point. In the absence of external stimulus (= 0), in

the limit $!_{c} = 1 \text{ (F } ! 0)$, the order parameter behaviour is F ()' (1)=C where C = (n 1) + (K 1)=2K, which gives, from the de nition F () / (c), a critical exponent = 1. On the other hand, at the critical point = 1 we have F (r) r=C which gives the Stevens exponent m = 1=2. Notice that the rate r plays the role of an external eld h and, from the usual statistical-physics de nition [21] F / h^{1= h} we get the critical exponent h = 2. As expected, these are the classical mean-eld exponents for branching processes, which seem to be valid for our model even with the presence of quenched disorder. An important conceptual point is that the Stevens-H ill exponent is indeed found to be indeed a statistical-physics critical exponent m = 1= h.

Optim ization of dynam ic range at criticality is very robust; the shape of the plot in Fig. 2D does not depend on param eters such as the average number of neighbours K or refractory period n. The random -network case seems to be a lower bound for the dynam ic-range enhancement. In future works, we will report that () is even more enhanced in low-dimensional networks, and presents non-rational Stevens exponents. Perhaps a better compromise between larger dynam ic range and biological realism would be a small-world or scale-free network.

As an example of the robustness of the results, if we put H odgk in H ux by elements with coupling with values $p_{ij} = 0$ or 1, which would correspond to a determ inistic case with the presence of absence of bonds, we obtain a (p) curve, where p is the probability that a bond exists between the elements. This (p) curve is very similar to the () curve of the present m odel, that is, (p) has a strong peak at the percolation phase transition $p = p_c$ (to be reported elsew here). However, the present m odel has the virtue of enabling analytical results that provide a benchm ark for the perform ance of networks with other topologies.

Now we discuss the possible relevance of our results to biological sensory processing. Recent notings show that projection cells in sensory systems are coupled via dendro-dendritic electrical synapses, for example -ganglionar cells in the retina [25, 26] and m itral cells in the olfactory bulb [18, 19, 20]. In most of these notings the electrical coupling is mediated by connexins, but pannexins could also be present, and could even be more important than connexins for providing electrical coupling between excitatory cells [27]. How ever, the functional role of this electrical coupling is largely unknown. We propose that the electrically coupled dendritic trees in these systems form an excitable network (where each element of our model represents an excitable dendritic patch). Our results are consistent with the reduction in sensitivity, dynam ical range and synchronization recently observed in retinal ganglion cell response of connexin-36 knockout mice [28].

In the case of olfactory system, we identify the excitable random network with the dendro-dendritic network in the glom enuli [18], and each element is interpreted as an active dendritic compartment containing ion channels. It is known that relevant electrical coupling between m itral cells is done at the glom enular level [19, 20] because only cells that have their apical dendrite tufts in the same glom enulus show synchronized activity. In connexin-36 knockout m ice, the synchronized activity of m itral cells is absent.

O ur hypothesis could be tested in the following way. The dynamic range of glom enuli is of the order of 30 dB as measured recently [2, 4] (in contrast to 10 dB of single olfactory receptor neurons). We predict that in connexin-36 knockout mice, this dynamic range shall be strongly reduced. Of course, for a decisive test, we need to exam ine if other electrical synapses based on connexin-45 [29] and pannexins [27] are irrelevant in olfactory glom enuli.

The text-book account of large dynam ic range in intensity coding is the recruitment model or its variants [5] where di erent elements, with diverse activation thresholds, are sequentially recruited. Our mechanism is not incompatible with this scenario, and we expect that threshold variability in our model enlarges the dynam ic range even more. How ever, we must remember that recruitment is a linear mechanism where, to explain each order of magnitude in dynam ic range, we must postulate a corresponding order of magnitude in activation thresholds, which is not a plausible assumption [5].

We may ask how the network could self-organize to the critical point $_{\rm c}$ = 1. It is not hard to conceive that hom eostatic mechanisms, acting on the number and conductance of gap junctions, could tune the system. It is well known that extensive pruning of gap junctions occurs during development and maturation. This could represent the initial self-organization process towards criticality. Spontaneous activity in the absence of input furnishes a signature of supercriticality that could be used as a feedback signal to control the system.

This self-tuning criticality has recently been proposed in a model of sound nonlinear amplication by H opfoscillators in the cochlea [30]. In the cochlear model, it is assumed that each oscillator is poised at its H opf bifurcation point, producing enhanced sensitivity and enlarged dynam ic range. The principal difference from our mechanism is that it is a model for individual cells, that is, it is not based on a collective phenom enon. This implies that the dynam icrange exponent must be classical (rational). In our case, the rational exponent m = 1=2 is a particularity of the random network, and, similarly to other statistical mechanism odels, more structured network topologies may have non-classical exponents.

A lthough in this work we restricted our attention to sensory processing, we note that the dynam ic range of more central networks could also be improved by the sam emechanism. Similarly to excitatory networks, inhibitory networks

must also work robustly in the presence of large variations in input. So, the presence of electrical synapses in cortical inhibitory networks [31] could re ect the same principle.

The mechanism for amplied nonlinear response due to wave creation and annihilation is a basic property of excitable media. We found in this work that, if active media are tuned at the critical point of activity propagation, the response is optimized. We proposed that this principle is present in electrically coupled excitable dendritic networks in projection neurons of sensory systems and is a generative mechanism for psychophysical laws. This computational principle based in critical activity could also be present in other brain regions and could be implemented in arti cial sensors by using excitable media as detectors.

This research is supported by CNPq, FACEPE, CAPES and PRONEX. The authors are grateful for discussions with A.C.Roque, R.F.Oliveira, D.Restrepo, T.Cleland, V.R.Vitorino de Assis and for encouragement from N. Caticha.

E lectronic address: osam e@ clrp.usp.br; Corresponding author

- ^Y E lectronic address: m copelli@ df.ufpe.br
- [1] S.S. Stevens, Psychophysics: Introduction to its Perceptual, Neural and Social Prospects (Wiley, New York, 1975).
- [2] M.W. achowiak and L.B.Cohen, Neuron 32, 723 (2001).
- [3] A.M. Angioy, A.D esogus, I.T. Barbarossa, P.Anderson, and B.S.Hansson, Chem. Senses 28, 279 (2003).
- [4] H.U.Fried, S.H.Fuss, and S.I.Korsching, Proc. Natl. A cad. Sci. USA 99, 3222 (2002).
- [5] T.A.C leland and C.Linster, Neural Computation 11, 1673 (1999).
- [6] M. Copelli, A. C. Roque, R. F. Oliveira, and O. Kinouchi, Phys. Rev. E 65, 060901 (2002).
- [7] M. Copelli and O. K inouchi, Physica A 349, 431 (2005).
- [8] M. Copelli, R. F. Oliveira, A. C. Roque, and O. Kinouchi, Neurocomputing 65–66, 691 (2005).
- [9] J.Reiser and H.M atthews, J.Physiol. 530 (2001).
- [10] A. Tom aru and T. Kurahashi, J. Neurophysiol. 93, 1880 (2005).
- [11] N.Chater and G.D.Brown, Cognition 69, B17 (1999).
- [12] L.S.Furtado and M.Copelli, Phys. Rev. E 73, 011907 (2006).
- [13] J.M. Beggs and D. Plenz, J. Neurosci. 23 11167 (2003).
- [14] C.Haldem an and J.M. Beggs, Phys. Rev. Lett. 94, 058101 (2005).
- [15] C.G.Langton, Physica D 42, 12 (1990).
- [16] P.Bak, How Nature Works: The Science of Self-Organized Criticality (Oxford Univ. Press, New York, 1997).
- [17] D.R. Chialvo, Physica A 340, 756 (2004).

[18] T.Kosaka, M.R.Deans, D.L.Paul, and K.Kosaka, Neurosci. 134, 757 (2005).

- [19] M.Migliore, M.L.Hines, and G.M.Shepherd, J.Comput. Neurosci. 18, 151 (2005).
- [20] J. Christie, C. Bark, S. G. Hormuzdi, I. Helbig, H. Monyer, and G. L. Westbrook, Neuron 46, 761 (2005).
- [21] J.M arro and R.D ickm an, Nonequilibrium Phase Transitions in Lattice Models (Cambridge University Press, 1999).
- [22] G.Laurent, Nat. Rev. Neurosci. 3, 884 (2002).
- [23] T.J.Lew is and J.R inzel, Neurocom puting 38-40, 763 (2001).
- [24] T.J.Lew is and J.R inzel, Network Comput. Neural Syst. 11, 299 (2000).
- [25] T.Schubert, J.Degen, K.W illecke, S.G.Hormuzdi, H.Monyer, and R.Weiler, J.Comp.Neurol. 485, 191 (2005).
- [26] S. Hidaka, Y. Akahori, and Y. Kurosawa, J. Neurosci. 24, 10553 (2005).
- [27] A. Vogt, S.G. Horm uzdi, and H. Monyer, Brain Res. Mol. Brain Res. 141, 113 (2005).
- [28] M.R.Deans, B.Volgyi, D.A.Goodenough, S.A.Bloom eld, and D.L.Paul, Neuron 36, 703 (2002).
- [29] C. Zhang and D. Restrepo, Brain Res. 929, 37 (2002).
- [30] S.Camalet, T.Duke, F.Jullicher, J.Prost, Proc. NatlAcad. Sci.USA 97, 3183 (2000).

6

[31] G. Sohl, S. Maxeiner, and K. Willecke, Nat. Rev. Neurosci. 6, 191 (2005).