Quantum mechanical calculation of the e ects of sti and rigid constraints in the conform ational equilibrium of the A lanine dipeptide

Pablo Echenique^{1,2} , Ivan Calvo^{1,2} and J.L.A $lonso^{1,2}$

¹ Departamento de F sica Teorica, Facultad de Ciencias, Universidad de Zaragoza, Pedro Cerbuna 12, 50009, Zaragoza, Spain.

² Instituto de Biocom putacion y F sica de los Sistem as Com plejos (BIFI), Edicio Cervantes, Corona de Aragon 42, 50009, Zaragoza, Spain.

M arch 31, 2022

A bstract

If constraints are in posed on a macrom olecule, two inequivalent classicalm odels may be used: the sti and the rigid one. This work studies the e ects of such constraints on the C onform ational Equilibrium D istribution (CED) of the model dipeptide HCO-L-A la-NH2 without any sim plifying assumption. We use ab initio Quantum Mechanics calculations including electron correlation at the MP2 level to describe the system , and we measure the conform ational dependence of all the correcting term s to the naive CED based in the Potential Energy Surface (PES) that appear when the constraints are considered. These terms are related to mass-metric tensors determ inants and also occur in the Fixm an's compensating potential. We show that some of the corrections are non-negligible if one is interested in the whole R am achandran space. On the other hand, if only the energetically lower region, containing the principal secondary structure elements, is assumed to be relevant, then, all correcting term sm ay be neglected up to peptides of considerable length. This is the rst time, as far as we know, that the analysis of the conform ational dependence of these correcting terms is performed in a relevant biom olecule with a realistic potential energy function.

PACS: 87.14 Ee, 87.15.-v, 87.15 Aa, 87.15 Cc, 89.75.-k

Corresponding author. E-m ail address: pnique@unizar.es

1 Introduction

In computer simulations of large complex systems, such as macromolecules and, specially, proteins [1{6], one of the main bottlenecks to design e cient algorithms is the necessity to sample an astronom ically large conformational space [3,7]. In addition, being the typical timescales of the dimension movements in a wide range, demandingly small timesteps must be used in Molecular Dynamics simulations in order to properly account for the fastest modes, which lie in the fem to second range. However, most of the biological interesting behaviour (allosteric transitions, protein folding, enzymatic catalysis) is related to the slow est conformational changes, which occur in the timescale of milliseconds or even seconds [4,8{11]. Fortunately, the fastest modes are also the most energetic ones and are rarely activated at room temperature. Therefore, in order to alleviate the computational problems and also simplify the images used to think about these elusive systems, one may naturally consider the reduction of the number of degrees of freedom describing macromolecules via the imposition of constraints [12].

How to study the conform ational equilibrium of these constrained systems has been an object of much debate [13{17]. Two di erent classical models exist in the literature which are conceptually [13{16,18,19}] and practically [6,13,20{24}] inequivalent. In the classical rigid model, the constraints are assumed to be exact and all the velocities that are orthogonal to the hypersurface de ned by them vanish. In the classical sti¹ model, on the other hand, the constraints are assumed to be approximate and they are implemented by a steep potential that drives the system to the constrained hypersurface. In this case, the orthogonal velocities are activated and may act as heat containers".

In this work, we do not address the question of which model is a better approximation of physical reality. A lthough, in the literature, it is commonly assumed (often implicitly) that the classical stim odel should be taken as a reference [6,9,16,19,20,22,26], we believe that this opinion is much in uenced by the use of popular classical force elds $[6,27\{37\}]$ (which are still by construction) and by the goal of reproducing their results at a lower computational cost, i.e., using rigid M olecular D ynamics simulations $[4,5,8,9,14,19,21\{23,25,26,38\{42\}]$. In our opinion, the question whether the rigid or the still model should be used to approximate the real quantum mechanical statistics of an arbitrary organic molecule has not been satisfactorily answered yet. For discussions about the topic, see references $[13\{15,17,18,43\{45\}]$. In this work, we adopt the cautious position that any of the two models may be useful in certain cases or for certain purposes and we study them both on equal footing. Our concern is, then, to study the elects that either way of imposing constraints causes in the conformational equilibrium of macromolecules.

In the Born-Oppenheim er approximation [46] custom arily used in Quantum Mechanics and in the majority of the classical force elds, the relevant degrees

¹Som e authors use the word exible to refer to this model [15,21,22,25]. W e, how ever, prefer to term it sti [18] and keep the name exible to refer to the case in which no constraints are in posed.

of freedom are the Euclidean (also called Cartesian by som e authors) 3n coordinates of the n nuclei. However, it is frequent to de ne a di erent set of coordinates in which the overall translation and rotation of the system are distinguished and the remaining 3n 6 degrees of freedom are chosen (according to di erent prescriptions as internal coordinates, which are simple geom etrical param eters (typically consisting of bond lengths, bond angles and dihedral angles) that describe the internal structure of the system [47].

In macrom olecules, the natural constraints are those derived from the relative rigidity of the internal covalent structure of groups of atoms that share a common center (and also from the rigidity of rotation around double or triple bonds) compared to the energetically \cheaper" rotation around single bonds. In internal coordinates, these chemical constraints may be directly implemented by asking that some conveniently selected hard coordinates (norm ally, bond lengths, bond angles and some dihedrals) have constant values or values that depend on the remaining soft coordinates (see ref. [15] for a de nition). In Euclidean coordinates, on the other hand, the expression of the constraints is more cumbersome e and complicated procedures [25,26,40,48{50] must be used at each timestep to implement them in Molecular Dynamics simulations. This is why, in the classical stimmedel, as well as in the rigid one, it is common to use internal coordinates and they are also the choice throughout this work.

In the equilibrium Statistical M echanics of both the sti and rigid models, the marginal probability density in the coordinate part of the phase space in these internal coordinates is not proportional to the naive $\exp[-V_{(q^i)}]$, where $V_{(q^i)}$ denotes the potential energy on the constrained hypersurface². Instead, some correcting terms that come from di erent sources must be added to the potential energy $V_{(q^i)}$ [13,15,18,19,39,51,52]. These terms involve determ inants of massmetric tensors and also of the H essian matrix of the constraining part of the potential (see sec. 2). If M onte C arlo simulations in the coordinate space are to be performed [5,53{57}] and the probability densities that correspond to any of these two models sam pled, the corrections should be included or, otherwise, show ed to be negligible.

A dditionally, the three di erent correcting term s are involved in the de nition of the so-called Fixm an's compensating potential [16], which is frequently used to reproduce the sti equilibrium distribution using rigid M olecular Dynam ics simulations $[9,14,19,21\{23,38,39,42,51]$.

Custom arily in the literature, some of these corrections to the potential energy are assumed to be independent of the conformation and thus dropped from the basic expressions. A lso, subtly entangled to the assumptions underlying m any classical results, a second type of approximation is made that consists of assuming that the equilibrium values of the hard coordinates do not depend on the soft coordinates.

In this work, we measure the conform ational dependence of all correcting terms and of the Fixm an's compensating potential in the model dipeptide HCO –

 $^{^2\}text{By}\,q^i$, we denote the soft internal coordinates of the system . See sec. 2 and the Appendix for a precise de nition .

L-A la-N H₂ without any simplifying assumption. The potential energy function is considered to be the electrice B om-O ppenheim erpotential for the nuclei derived from ab initio quantum mechanical calculations including electron correlation at the M P 2 level. We also repeat the calculations, with the same basis set (6-31++G (d,p)) and at the H artree-Fock level of the theory in order to investigate if this less demanding method without electron correlation may be used in further studies. It is the strate of study is performed in a relevant biom olecule with a realistic potential energy function.

In sec. 2, we introduce the notation to be used and derive the Statistical M echanics form ulae of the rigid and sti m odels in the general case. In sec. 3, we describe the computational methods used and we sum marize the factorization of the external coordinates presented in ref. [58]. Sec. 4 is devoted to the presentation and discussion of the assessment of the approximation that consists of neglecting the di erent corrections to the potential energy in the m odel dipeptide $H C O - L - A la - N H_2$, without any simplifying assumption, which is the central aim of this work. The conclusions are summarized in sec. 5. Finally, in the appendix, we discuss the use of the di erent approximations in the literature and we give a precise de nition of exactly and approximately separable hard and soft coordinates which will shed some light on the relation between the di erent types of sim pli cations aforem entioned.

2 Theory

F irst of all, it is convenient to introduce certain notational conventions that will be used extensively in the rest of the work:

The system under scrutiny will be a set of n mass points term ed atom s. The Euclidean coordinates of the atom in a set of axes xed in space are denoted by x. The subscript runs from 1 to n.

The curvilinear coordinates suitable to describe the system will be denoted by q; = 1;:::;3n and the set of Euclidean coordinates by x when no explicit reference to the atom s index needs to be made. We shall offen use N = 3n for the total number of degrees of freedom.

The coordinates q are split into $(q^A;q^a)$; $a = 7; \ldots; N$. The rst six are term ed external coordinates and are denoted by q^A . They describe the overall position and orientation of the system with respect to a fram e xed in space (see ref. 58 for further details). The coordinates q^a are said internal coordinates and determ ine the positions of the atom s in the fram e xed in the system. They parameterize what we shall call the internal subspace or conform ational space, denoted by I and the coordinates q^A parameterize the external subspace, denoted by E.

The general set-up of the problem may be described as follows: Instead of us being interested on the conformational equilibrium of the system in

the external subspace E plus the whole internal subspace I (i.e., the whole space, denoted by E I), we wish to nd the probability density on a hypersurface I of dimension M (plus the external subspace E), i.e., on E \cdot .

In typical internal coordinates \hat{q} , norm ally consisting of bond lengths, bond angles and dihedral angles (see ref. 59 and references therein), the hypersurface is described via $L := N \quad M \quad 6$ constraints:

$$q^{I} = f^{I}(q^{i})$$
 $I = M + 7; ...; N$; (2.1)

where the q^a are split into q^a $(q^{i};q^{I})$, and the qⁱ, i = 7; :::;M + 6, which parameterize , are called internal soft coordinates, whereas the q^I are term ed hard coordinates. The external coordinates q^A, together with the qⁱ, form the whole set of soft coordinates, denoted by q^{II} $(q^{A};q^{i})$, u = 1; :::;M + 6.

In table 1, a sum m ary of the indices used is given.

Indiæs	Range	N um ber	D escription	
; ; ;::: ; ; ;:::	1;:::;n 1;:::;N	n N = 3n	A tom s A 11 coordinates	
A;B;C;:::	1 ;:::; 6	6	External coordinates	
a;b;c;:::	7 ;:::; N	N 6	Internal coordinates	
i;j;k;:::	7 ;:::; M + 6	М	Soft internal coordinates	
I;J;K ;:::	M + 7;:::;N	L = N M 6	Hard internal coordinates	
u;v;w;:::	1 ;:::; M + 6	M + 6	All soft coordinates	

Table 1: De nition of the indices used.

2.1 Classical sti model

In the classical sti model, the constraints in eq.2.1 are implemented by imposing an strong energy penalization when the internal conformation of the system, described by q^a , departs from the constrained hypersurface. To ensure this, we must have that the potential energy function in I satisfies exertain conditions. First, we write the potential V (q^a) as follow s^3 :

$$\nabla (q^{i};q^{I}) = \bigvee_{\substack{q^{i};f^{I}(q^{i})\\ \forall (q^{i})}} q^{i};q^{I}) = \bigvee_{\substack{q^{i};f^{I}(q^{i})\\ \forall (q^{i})}} q^{i};q^{I}) = \bigvee_{\substack{q^{i};q^{I}\\ \forall (q^{i})}} q^{i};q^{I}) = \bigvee_{\substack{q^{i};q^{I}\\ \forall (q^{i})}} q^{i};q^{I})$$
(2.2)

Next, we impose the following conditions on the constraining potential $V_c \ (q^i; q^I)$ de ned above:

³N ote that we have simply added and subtracted from the total potential energy V $(q^i;q^I)$ V (q^a) of the system the same quantity, V $(q^i;f^I(q^i))$.

- (i) That V_c qⁱ;f^I (qⁱ) V_c (qⁱ;q^I) 8qⁱ;q^I, i.e., that be the global minimum of V_c (and, henceforth, a local one too) with respect to variations of the hard coordinates.
- (ii) That, for small variations q^{I} on the hard coordinates (i.e., for changes q^{I} considered as physically irrelevant), the associated changes in $V_{c}(q^{i};q^{I})$ are much larger than the therm all energy RT.

The advantages of this form ulation, much sim ilar to that on [15], are many. First, it sets a convenient fram ework for the derivation of the Statistical M echanics form ulae of the classical stim odel relating it to the fully exible model in the whole space E I. Second, it clearly separates the potential energy on

from the part that is responsible of in plementing the constraints. Third, contrarily to the form ulation based on delta functions [51], it allows to clearly understand the necessity of including the correcting term associated to the determ inant of the Hessian of V_c (see the derivation that follows). Finally, and m ore in portantly for us, it provides a direct prescription for calculating V (qⁱ) and (the Potential Energy Surface (PES), frequently used in Q uantum Chem - istry calculations [60{64}) via geometry optimization at xed values of the soft coordinates.

We also remark that, in order to satisfy point (ii) above and to allow the derivation of the di erent correcting terms that follows and the validity of the nalexpressions, the hard coordinates q^{I} must be indeed hard, how ever, the soft coordinates q^{i} do not have to be soft (in the sense that they produce energetic changes much smaller than R T when varied). They may be interesting for some other reason and hence voluntarily picked to describe the system studied, without altering the form use presented in this section. Despite this quali cations, the terms soft and hard will be kept in this work for consistence with most of the existing literature [15,18,52,65,66], although, in some cases, the labels important and unimportant (for q^{i} and q^{I} respectively), proposed by K arplus and K ushick [67], may be more appropriate.

In the case of the m odel dipeptide H C O -L -A la-N H $_2$ investigated in this work, for example, the barriers in the R am achandran angles and m ay be as large as

40 RT, how ever, the study of sm all dipeptides is norm ally aim ed to the design of e ective potentials for polypeptides [68{70}], where long-range interactions in the sequencem ay compensate these local energy penalizations. This and the fact that the R am achandran angles are the relevant degrees of freedom to describe the conform ation of the backbone of these system s, m ake it convenient to choose them as soft coordinates qⁱ despite the fact that they m ay be energetically hard in the case of the dipeptide HCO-L-A la-NH₂. As rem arked above, this does not a ect the calculations.

Now, due to condition (ii) above, the statistical weights of the conform ations which lie far away from the constrained hypersurface are negligible and, therefore, it su ces to describe the system in the vicinity of the equilibrium values of the q^{I} . In this region, for each value of the internal soft coordinates q^{i} , we may expand V_{c} ($q^{i}; q^{I}$) in eq. 2.2 up to second order in the hard coordinates around (i.e., around $q^{I} = f^{I}$ (q^{i})) and drop the higher order term s:

$$V_{c}(q^{i};q^{I}) ' V_{c} q^{i};f^{I}(q^{i}) + \frac{\varrho V_{c}}{\varrho q^{J}} q^{J} f^{J}(q^{i}) + \frac{1}{2} | \frac{\varrho^{2} V_{c}}{|\frac{\varrho q^{J}}{2} | \frac{q^{2}}{q^{I}} | \frac{q^{J}}{q^{I}} f^{J}(q^{i}) q^{K} f^{K}(q^{i}) ; (2.3)$$

where the subindex indicates evaluation on the constrained hypersurface and a more compact notation, H $(\!q^i)$, has been introduced for the Hessian matrix of V_c with respect to the hard variables evaluated on $\$. Also, the Einstein's sum convention is assumed on repeated indices.

In this expression, the zeroth order term $V_c q^i; f^I(q^i)$ is zero by de nition of V_c (see eq.2.2) and the linear term is also zero, because of the condition (i) above. Hence, the rst non-zero term of the expansion in eq.2.3 is the second order one. U sing this, together with eq.2.2, we may write the sti H am iltonian

$$H_{s}(q;p) \coloneqq \frac{1}{2}p G (q^{u};q^{I})p + V (q^{i}) + \frac{k}{2}H_{JK}(q^{i}) q^{J} f^{J}(q^{i}) q^{K} f^{K}(q^{i}); \quad (2.4)$$

the mass-metric tensor G being

$$G \quad (q^{\mu}; q^{I}) := \frac{X^{N}}{e_{1}} \frac{(e_{1} \times e_{1})}{e_{1}} m \frac{(e_{1} \times e_{1})}{e_{1}} m (2.5)$$

and G its inverse, de ned by

$$G (q^{\mu}; q^{I}) G (q^{\mu}; q^{I}) = ; \qquad (2.6)$$

where denotes the K ronecker's delta.

Therefore, the sti partition function of the system is⁴

$$Z_{s} = \frac{QM}{h^{N}} dq dp exp H_{s}(q;p); \qquad (2.7)$$

where h is Planck's constant, we denote = 1=R T (permole energy units are used throughout the article, so R T is preferred over $k_B T$) and $_{QM}$ is a combinatorial number that accounts for quantum indistinguishability and that must be specified in each particular case (e.g., for a gas of N indistinguishable particles, $_{QM} = 1=N$!).

Now, using the condition (ii) again, the q^{I} appearing in the mass-metric tensor G in H_s (in eq.2.7) can be approximately evaluated at their equilibrium values $f^{I}(q^{i})$, yielding, for the stipartition function,

 $^{^4}$ No Jacobian appears in the integral measure because q and p are obtained from the Euclidean coordinates via a canonical transform ation [71].

$$Z_{s} = \frac{QM}{h^{N}} dq^{\mu} dq^{I} dp \exp \frac{1}{2} p G q^{\mu}; f^{I}(q^{i}) p + V(q^{i}) + \frac{1}{2} H_{JK}(q^{i}) q^{J} f^{J}(q^{i}) q^{K} f^{K}(q^{i}) ;$$
(2.8)

If we now integrate over the hard coordinates q^{I} , we have

$$Z_{s} = \frac{2}{h^{N}} \frac{\frac{1}{2}}{h^{N}} \frac{\sum_{q \in M} Z}{dq^{u} dp exp} = \frac{1}{2} p G q^{u}; f^{I}(q^{i}) p + Q^{i} + V (q^{i}) + T \frac{R}{2} \ln \det H (q^{i}) :$$

$$(2.9)$$

where the part of the result of the G aussian integral consisting of det $^{\rm 1=2}{\rm H}$ has been taken to the exponent.

Note that the H essian m atrix H $_{\rm JK}$ involves only derivatives with respect to the hard coordinates (see eq.2.3), so that the m inimization protocolem bodied in eq.2.2 (which is identical to the procedure followed in Q uantum C hem istry for computing the PES along reaction coordinates) guarantees that H $_{\rm JK}$ is positive de ned and, hence, detH is positive, allowing to take its logarithm as in the previous expression. The fact that it is only this 'partial H essian' that m akes sense in the computation of equilibrium properties along soft (or reaction) coordinates, has been recently pointed out in ref. 72.

It is also frequent to integrate over the m om enta in the partition function. Doing this in eq.2.9 and taking the determ inant of the mass-metric tensor that show sup⁵ to the exponent, we may write the partition function as an integral only on the coordinates:

$$Z_{s} = {}_{s}(T) dq^{u} \exp V (q^{i}) + T \frac{R}{2} \ln \det H (q^{i})$$

$$T \frac{R}{2} \ln \det G q^{u}; f^{I}(q^{i}) ; \qquad (2.10)$$

where the multiplicative factor that depends on T has been de ned as follows:

$$_{s}(T) \coloneqq \frac{2}{h^{N}} \stackrel{\frac{N+L}{2}}{\frac{QM}{h^{N}}} :$$
 (2.11)

If the exponent in eq. 2.10 is seen as a free energy, then, V (q^i) may be regarded as the internal energy and the two conform ation-dependent correcting

 $^{^5\}mathrm{N}$ ote that, by G , we denote the m atrix that corresponds to the m ass-m etric tensor with two covariant indices G $\,$. The same convention has been followed for the H essian m atrix H in eq.2.9 and for the reduced m ass-m etric tensor g in eq.2.21.

terms that are added to it as elective entropies (which is compatible with their being linear in RT). The second one comes only from the desire to write the marginal probabilities in the coordinate space (i.e., averaging the momenta) and may be called a kinetic entropy [17], the rst term, on the other hand, is truly an entropic term that comes from the averaging out of certain degrees of freedom and it is reminiscent of the conformational or con gurational entropies appearing in quasiharm onic analysis [6,67,73].

In this spirit, we de ne

$$F_{s}(q^{u}) \coloneqq V(q^{i}) T S_{s}^{c}(q^{i}) + S_{s}^{k}(q^{u}) ; \qquad (2.12a)$$

$$S_s^c(q^i) \coloneqq \frac{R}{2} \ln \det H(q^i)$$
; (2.12b)

$$S_{s}^{k}(q^{\mu}) \coloneqq \frac{R}{2} \ln \det G q^{\mu}; f^{I}(q^{i})^{L} : \qquad (2.12c)$$

In such a way that the sti equilibrium probability in the soft subspace ${\rm E}$ is given by

$$P_{s}(q^{\mu}) = \frac{\exp F_{s}(q^{\mu})}{Z_{s}^{0}}; \text{ with } Z_{s}^{0} \coloneqq dq^{\mu} \exp F_{s}(q^{\mu}) : (2.13)$$

Now, it is worth remarking that, although the kinetic entropy S_s^k depends on the external coordinates q^A , we have recently shown [58] that the determ inant of the mass-metric tensor G may be written, for any molecule, general internal coordinates and arbitrary constraints, as a product of two functions; one depending only on the external coordinates, and the other only on the internal ones q^A . Hence the externals-dependent factor in eq. 2.12c may be integrated out independently to yield an elective free energy and a probability density P_s that depend only on the soft internals q^i (see sec. 3.1).

2.2 Classical rigid model

If the relations in eq.2.1 are considered to hold exactly and are treated as holonom ic constraints, the H am iltonian function that describes the C lassical M echanics in the subspace (E) (E), spanned by the coordinates q^u , may be written as follows:

$$H_{r}(q^{u}; u) \coloneqq \frac{1}{2} vg^{vw}(q^{u}) + V(q^{i}); \qquad (2.14)$$

where the reduced mass-metric tensor $g_{vw}\;(q^n)$ in E , that appears in the kinetic energy, is (see what follows)

$$g_{vw} (q^{u}) = G_{vw} q^{u}; f^{I} (q^{i}) + \frac{\partial f^{J} (q^{i})}{\partial q^{v}} G_{JK} q^{u}; f^{I} (q^{i}) \frac{\partial f^{K} (q^{i})}{\partial q^{w}} + G_{vK} q^{u}; f^{I} (q^{i}) \frac{\partial f^{K} (q^{i})}{\partial q^{w}} + \frac{\partial f^{J} (q^{i})}{\partial q^{v}} G_{Jw} q^{u}; f^{I} (q^{i}) \rightleftharpoons (q^{i}) \rightleftharpoons (q^{i}) = \frac{\partial f^{I}}{\partial q^{v}} G_{Jw} q^{u}; f^{I} (q^{i}) \rightleftharpoons (q^{i}) \rightleftharpoons (q^{i}) = \frac{\partial f^{I}}{\partial q^{v}} G_{Jw} q^{u}; f^{I} (q^{i}) \land (q^{i}) \models (2.15)$$

and $g^{\nu w}~(q^{\mu})$ is de ned to be its inverse in the sense of eq. 2.6. Also, the notation

$$f' := \begin{array}{c} (\\ q^{11} \\ f' := \end{array} \begin{array}{c} if \ u := \\ f^{I} (q^{i}) \\ f' := \end{array} \begin{array}{c} M + 6 \\ f' := \\ f^{I} (q^{i}) \end{array} \begin{array}{c} (2.16) \\ f' := \\$$

has been introduced for convenience.

Note that eq.2.15 m ay derived from the unconstrained H am iltonian in (E I),

$$H (q;p) := \frac{1}{2}p G (q)p + V (q^{a});$$
 (2.17)

using the constraints in eq.2.1, together with its time derivatives (denoted by an overdot: as in A-)

$$q^{I} \coloneqq \frac{\theta f^{I}(q^{i})}{\theta q^{j}} q^{j}$$
(2.18)

and de ning the momenta $\ _{v}$ as

$$v \coloneqq g_{vw} (q^{\mu}) q^{\mu} = g_{vw} (q^{\mu}) G^{w} q^{\mu}; f^{I} (q^{i}) p :$$
 (2.19)

Hence, the rigid partition function is

$$Z_{r} = \frac{QM}{h^{M+6}} dq^{\mu} d_{\mu} \exp \frac{1}{2} v g^{\nu w} (q^{\mu}) v + V (q^{i}) :$$
 (2.20)

Integrating over the m om enta, we obtain the m arginal probability density in the coordinate space analogous to eq. 2.10:

$$Z_{r} = {}_{r}(T) dq^{\mu} exp \qquad V (q^{i}) \qquad T \frac{R}{2} \ln \det g(q^{\mu}) ; \qquad (2.21)$$

where

$$r(T) := \frac{2}{h^{\frac{M+6}{2}}} \frac{QM}{h^{\frac{M+6}{2}}} :$$
 (2.22)

Repeating the analogy with free energies and entropies in the last paragraphs of the previous subsection, we dene

$$F_{r}(q^{\mu}) \coloneqq V(q^{i}) \quad TS_{r}^{k}(q^{\mu}); \qquad (2.23a)$$

$$S_{r}^{k}(q^{u}) \coloneqq \frac{R}{2} \ln^{11} \det g(q^{u})^{T};$$
 (2.23b)

being the rigid equilibrium probability in the soft subspace E

$$P_r(q^{\mu}) = \frac{\exp \left(F_r(q^{\mu})\right)}{Z_r^0}; \quad \text{with} \quad Z_r^0 \coloneqq dq^{\mu} \exp \left(F_r(q^{\mu})\right): \quad (2.24)$$

As in the case of G, we have shown in ref. [58] that the determ inant of the reduced mass-metric tensor g may be written, for any molecule, general internal coordinates and arbitrary constraints, as a product of two functions; one depending only on the external coordinates, and the other only on the internal ones q^i . Hence the externals-dependent factor in detg(q^i) may be integrated out independently to yield a free energy and a probability density P_r that depend only on the soft internals q^i (see sec 3.1).

To end this subsection, we rem ark that it is frequent in the literature [9,18,19, 21{23,38,42,51,57} to de ne the so-called Fixm an's compensating potential [16] as the di erence between $F_s(q^u)$, in eq. 2.12, and $F_r(q^u)$, de ned above, i.e.,

$$V_{F}(q^{u}) \coloneqq TS_{r}^{k}(q^{u}_{H}) TS_{s}^{c}(q^{i}) TS_{s}^{k}(q^{u}_{H}) =$$
$$= \frac{RT}{2} \ln \frac{\det G(q^{u})}{\det H(q^{i}) \det g(q^{u})} : \qquad (2.25)$$

Hence, perform ing rigid M olecular D ynam ics simulations, which would yield an equilibrium distribution proportional to exp[$F_r(q^\mu)$], and adding $V_F(q^\mu)$ to the potential energy V (q^i) one can reproduce instead the sti probability density P_s / exp[$F_s(q^\mu)$] [14,18,19,21{23,38,39,42,51}]. This allows to obtain at a lower computational cost (due to the tim escale problem s discussed in the introduction) equilibrium averages that otherwise must be extracted from expensive fully exible whole-space simulations. In fact, it seems that this particular application of the theoretical tools herein described, and not the search for the correct probability density to sam ple in M onte C arb simulations, was what prom pted the interest in the study of mass-metric tensors e ects.

Finally, in table 2 we summarize the equilibrium probability densities and the di erent correcting terms derived in this section.

Classical Sti Model	ClassicalRigid Model			
$P_{s}(q^{\mu}) = \frac{\exp \left(F_{s}(q^{\mu})\right)}{Z_{s}^{0}}$	$\mathbb{P}_{r}(q^{u}) = \frac{\exp \left(\frac{F_{r}(q^{u})}{Z_{r}^{0}} \right)}{\mathbb{Z}_{r}^{0}}$			
$\mathbb{F}_{s}(q^{u}) \coloneqq \mathbb{V}(q^{i}) = \mathbb{T} S_{s}^{c}(q^{i}) + S_{s}^{k}(q^{u})$	$F_r(q^u) \coloneqq V(q^i) TS_r^k(q^u)$			
$S_{s}^{k}(q^{\mu}) \coloneqq \frac{R}{2} \ln \det G q^{\mu}; f^{I}(q^{i})$	$S_r^k(q^u) \coloneqq \frac{R}{2} \ln \det(q^u)^i$			
$S_{s}^{c}(q^{i}) \coloneqq \frac{R}{2} \ln \det H(q^{i})$				

Table 2: Equilibrium probability densities and correcting term s to the potential energy $V(q^i)$ in the classical sti and rigid models of constraints.

3 M ethods

3.1 Factorization of the external coordinates

In the recent work [58], we have shown that the determ inant of the massm etric tensor G in eq.2.12c can be written as follows if the SASM IC [59] coordinates for general branched molecules are used:

det G =
$$m^{3} \sin^{2} r^{4} \sin^{2} r^{3}$$
 (3.1)

where the r are bond lengths and the bond angles.

Note that this expression, whose validity was proved for the more particular case of serial polymers by Go and Scheraga [15] and, before, by Volkenstein [74], does not explicitly depend on the dihedral angles. However, it may depend on them via the hard coordinates if the constraints in the form presented in eq.2.1 are used.

The term depending on the masses of the atom s in the expression above may be dropped from eq.2.12c, because it does not depend on the conformation, and the only part of detG that depend on the external coordinates, \sin^2 , may be integrated out in eq.2.10 (is one of the externals $\frac{1}{4}$ that describe the overall orientation of the molecule; see ref. 58 for further details). Hence, the kinetic entropy due to the mass-metric tensor G in the sti case, may be written, up to additive constants, as

$$S_{s}^{k}(q^{i}) = \frac{R}{2} \prod_{i=2}^{W} \ln r^{4} + \prod_{i=3}^{W} \ln sin^{2} ; \qquad (3.2)$$

where the individual contributions of each degree of freedom have been factorized.

A lso in reference [58], we have shown that the determ inant of the reduced mass-metric tensor g in eq.2.23b can be written as follows:

$$detg = \sin^2 \quad detg_i (q^i); \tag{3.3}$$

being the matrix g₂

$$g_{2} = \begin{bmatrix} 0 & & & 1 \\ m_{tot} \mathbf{I}^{(3)} & m_{tot} \mathbf{v}^{(\mathcal{R})} & & & \\ m_{tot} \mathbf{v}^{T} (\mathcal{R}) & J & & & \\ \hline \\ m_{tot} \mathbf{v}^{T} (\mathcal{R}) & J & & & \\ \hline \\ m_{tot} \frac{\mathcal{Q}\mathcal{R}}{\mathcal{Q}_{1}^{i}} & m & \frac{\mathcal{Q}\mathcal{R}^{0}}{\mathcal{Q}_{1}^{i}} \mathbf{x}^{0} & \\ \hline \\ & & & \\ \hline \\ m_{tot} \frac{\mathcal{Q}\mathcal{R}}{\mathcal{Q}_{1}^{i}} & m & \frac{\mathcal{Q}\mathcal{R}^{0}}{\mathcal{Q}_{1}^{i}} \mathbf{x}^{0} & \\ \hline \\ & & & \\ \hline \\ & & & \\ \hline \\ \end{array} \end{bmatrix} \begin{bmatrix} 1 \\ t^{(3)} \mathcal{R}^{(3)} \\ m & \frac{\mathcal{Q}\mathcal{R}^{0}}{\mathcal{Q}_{1}^{i}} & \\ m & \frac{\mathcal{Q}\mathcal{R}^{0}}{\mathcal{Q}_{1}^{i}} & \\ \hline \\ & & \\ \hline \\ & & \\ \hline \\ \hline \\ \end{bmatrix} \begin{bmatrix} 1 \\ m \\ \mathcal{Q}\mathcal{R} \\ \mathcal{Q}^{i} \\ \mathcal{Q}^{i} \\ \mathcal{Q}^{j} \\ \mathcal{Q}^$$

where the superindex T indicates matrix transposition, $I^{(3)}$ denotes the 3 3 identity matrix and x^0 is the position of atom in the reference frame exed in the system (the primed' reference frame).

Additionally, we denote the total mass of the system by $m_{tot} = m_{tot}^{P} m_{tot}$, the position of the center of mass of the system in the primed reference frame by $R = m_{tot}^{-1} m_{tot}^{P} m_{tot}^{A}$ and the inertia tensor of the system, also in the primed reference frame, by

$$J \coloneqq \sum_{i=1}^{n} m_{i} (x^{02})^{2} + (x^{03})^{2}) \sum_{i=1}^{n} m_{i} x^{01} x^{02} \sum_{i=1}^{n} m_{i} x^{01} x^{03} \sum_{i=1}^{n} m_{i} (x^{01})^{2} + (x^{03})^{2}) \sum_{i=1}^{n} m_{i} x^{02} x^{03} \sum_{i=1}^{n} m_{i} (x^{01})^{2} + (x^{02})^{2})$$

$$(3.5)$$

The matrix $v(\mathbf{\tilde{R}})$ is de ned as:

and denotes the usual vector cross product.

Then, since \sin^2 may be integrated out in eq.2.21, we can write, om itting additive constants, the kinetic entropy associated to the reduced mass-metric tensor g depending only on the soft internals q^i :

$$S_{r}^{k}(q^{i}) = \frac{R}{2} \ln \det g_{2}(q^{i})$$
 : (3.7)

Finally, one may note that, since \sin^2 divides out in the second line of eq. 2.25 or, otherwise stated, eqs. 3.2 and 3.7 may be introduced in the rst line, then the Fixm an's potential is independent of the external coordinates as well.

3.2 Computational M ethods

In the particular molecule treated in this work (the model dipeptide HCO-L-Ala-NH₂ in g. 1), the formulae in the preceding sections must be used with M = 2, being the internal soft coordinates qⁱ (;) the typical R am achandran angles [75] (see table 3), the total number of coordinates N = 48 and the number of hard internals L = 40.

Regarding the side chain angle , it has been argued elsew here [59] that it is soft with the same right as the angles and , i.e., the barriers that hinder the rotation on this dihedral are comparable to the ones existing in the Ram achandran surface. However, the height of these barriers is su cient (6-12 RT, see ref. [59]) for the condition (ii) in sec. 2.1 to hold and, therefore, its inclusion in the set of hard coordinates is convenient due to its unim portant character (see discussion in sec. 2.1). Moreover, to describe the behaviour associated to with a probability density di erent from a Gaussian distribution (i.e., its potential energy di erent from an harm onic oscillator), for example with the tools used in the eld of circular statistics [76{78], would severely complicate the derivation of the classical stim odel without adding any conceptual insight to the problem. In addition, although is a periodic coordinate with threefold symmetry, the considerable height of the barriers between consecutive m in im a allow s to m ake the quadratic assumption in eq.2.3 at each equivalent valley and perm its the approximation of the integral on by three times a Gaussian integral. The multiplicative factor 3 simply adds a tem perature- and conform ation-independent reference to the con gurational entropy S_s^c in eq.2.12b.

The same considerations are applied to the dihedral angles, $!_0$ and $!_1$ (see table 3), that describe the rotation around the peptide bond, and the quadratic approximation described above can also be used, since the heights of the rotation barriers around these degrees of freedom are even larger than the ones in the case of .




Figure 1: A tom num eration of the protected dipeptide $HCO - L-A la - NH_2$.

The ab initio quantum mechanical calculations have been done with the

package GAMESS [79] under Linux and in 3.20 GHz PIV machines. The coordinates used for the HCO-L-A la-NH₂ dipeptide in the GAMESS input les and the ones used to generate them with automatic Perl scripts are the SAS-MIC coordinates introduced in ref. 59. They are presented in table 3 indicating the name of the conventional dihedral angles (see also g. 1 for reference). To perform the energy optimizations, how ever, they have been converted to Delocalized Coordinates [80] in order to accelerate convergence.

Atom name	Bond length	Bond angle	Dihedralangle
H ₁			
C 2	(2,1)		
N ₃	(3,2)	(3,2,1)	
O 4	(4,2)	(4 , 2 , 1)	(4,2,1,3)
C 5	(5,3)	(5,3,2)	! ₀ ≒ (5,3,2,1)
H ₆	(6,3)	(6,3,2)	(6,3,2,5)
C 7	(7,5)	(7 , 5 , 3)	: (7,5,3,2)
C 8	(8,5)	(8,5,3)	(8,5,3,7)
H 9	(9,5)	(9 , 5 , 3)	(9,5,3,7)
H 10	(10,8)	(10,8,5)	; (10,8,5,3)
H 11	(11,8)	(11,8,5)	(11,8,5,10)
H ₁₂	(12,8)	(12,8,5)	(12,8,5,10)
N 13	(13,7)	(13,7,5)	; (13,7,5,3)
O 14	(14,7)	(14,7,5)	(14,7,5,13)
H ₁₅	(15,13)	(15,13,7)	! ₁ ≔ (15,13,7,5)
H 16	(16,13)	(16,13,7)	(16,13,7,15)

Table 3: SA SM IC internal coordinates (E chenique P.and A lonso J.L., To be published in J. Comp. Chem., arXiv:q-bio.BM/0511004) in Z-m atrix form of the protected dipeptide $H CO - L - A = N H_2$. Principal dihedrals are indicated in bold face and their typical biochem ical name is given.

First, we have calculated the typical Potential Energy Surface (PES) in a regular 12x12 grid of the bidim ensional space spanned by the Ram achandran angles and , with both angles ranging from 165° to 165° in steps of 30°. This has been done by running constrained energy optimizations at the MP2/6-31++G (d,p) level of the theory, freezing the two Ram achandran angles at each value of the grid, starting from geom etnies previously optimized at a lower level of the theory and setting the gradient convergence criterium to OPTTOL=10⁵ and the self-consistent Hartree-Fock convergence criterium to CONV=10⁶.

The results of these calculations (which took 100 days of CPU time) are 144 conform ations that de ne and the values of V (;) at these points (the PES itself).

Then, at each optim ized point of $\$, we have calculated the Hessian m atrix in the coordinates of table 3 removing the rows and columns corresponding to

the soft angles and , the result being the matrix H (;) in eq.2.12b. This has been done, again, at the M P2/6-31++G (d,p) level of the theory, taking 140 days of CPU time.

Eqs. 32 and 3.7 in sec. 3.1 have been used to calculate the kinetic entropy terms associated to the determ inants of the mass-metric tensors G and g, respectively. The quantities in eq. 32, being simply internal coordinates, have been directly extracted from the GAMESS output les via autom ated Perl scripts. On the other hand, in order to calculate the matrix g_2 in eq. 3.4 that appears in the kinetic entropy of the classical rigid m odel, the Euclidean coordinates \mathbf{x}^0 of the 16 atom s in the reference frame xed in the system, as well as their derivatives with respect to qⁱ (;), must be computed. For this, two additional 12x12 grids as the one described above have been computed; one of them displaced 2° in the positive -direction and the other one displaced 2° in the positive -direction. This has been done, again, at the M P2/6-31++G(d,p)level of the theory, starting from the optim ized structures found in the com putation of the PES described above and taking 75 days of CPU time each grid. Using the values of the positions \mathbf{x}^0 in these two new grids and also in the original one, the derivatives of these quantities with respect to the angles and , appearing in g_2 , have been num erically obtained as nite di erences.

The three calculations have been repeated for six special points in the Ram achandran space that correspond to important elements of secondary structure (see sec. 4), the total CPU time needed for computing all correcting terms at these points has been 16 days. A total of 406 days of CPU time has been needed to perform the whole study at the MP2/6-31++G (d,p) level of the theory.

F inally, we have repeated all the calculations at the HF/6-31++G (d,p) level of the theory in order to investigate if this less dem anding m ethod (10 days for the PES, 8 days for the H essians, 10 days for each displaced grid, 2 days for the special secondary structure points, being a total of 40 days of CPU time) may be used instead of MP2 in further studies.

4 Results

In table 4, the maximum variation, the average and the standard deviation in the 12x12 grid de ned in the Ramachandran space of the protected dipeptide HCO-L-A la-NH₂ are shown for the three energy surfaces, V , F_s and F_r (see eqs.2.12 and 2.23), for the three correcting terms, TS_s^k , TS_s^c , and TS_r^k and for the Fixm an's compensating potential V_F (see eq.2.25). All the functions have been referenced to zero in the grid.

In g.2, the Potential Energy Surface V , at the M P2/6-31++G (d,p) level of the theory, is depicted with the reference set to zero for visual convenience⁶. Neither the surfaces de ned by F_s and F_r at the M P2/6-31++G (d,p) level of

 $^{^{6}\}mathrm{At}$ the level of the theory used in the calculations, the m inim um of V (;) in the grid is -416.0733418995 hartree.

	MP2/6-31++G(d,p)			HF/	HF/6-31++G (d,p)		
	M ax.ª	Ave. ^b	Std.°	M ax.ª	Ave. ^b	Std.°	
V	21.64	6.76	3.88	23.62	6.92	4.35	
Fs	21.43	6.47	3.93	23.78	7.17	4.38	
Fr	21.09	6.46	3.82	23.09	6.76	4.31	
ΤS ^k _s	0.24	0.09	0.05	0.23	0.09	0.04	
ΤSsc	1.67	0.98	0.32	1.34	0.63	0.30	
ΤSr	0.81	0.37	0.12	0.75	0.38	0.12	
$V_{\rm F}$	1.68	0.89	0.30	1.35	0.55	0.27	

Table 4: ^aM aximum variation, ^baverage and ^cstandard deviation in the 12x12 grid de ned in the Ram achandran space of the protected dipeptide HCO-L-A la-NH₂ for the three energy surfaces, V , F_s and F_r , the three correcting term s, TS_s^k , TS_s^c , and TS_r^k and the Fixm an's compensating potential V_F . The results at both M P2/6-31++G (d,p) and HF/6-31++G (d,p) levels of the theory are presented and all the functions have been referenced to zero in the grid. The units used are kcal/m ol.

the theory nor the three energy surfaces V , F_s and F_r at HF/6-31++G (d,p) are shown graphically since they are visually very similar to the surface in g.2.

In g.3, the three correcting term s, TS_s^k , TS_s^c and TS_r^k and the Fixman's compensating potential V_F , at the M P2/6-31++G (d,p) level of the theory, are depicted with the reference set to zero. The analogous surfaces at the HF/6-31++G (d,p) level of the theory are visually very similar to the ones in g.3 and have been therefore om itted.

From the results presented, one may conclude that, although the conform ational dependence of the correcting terms TS_s^k , TS_s^c and TS_r^k is more than an order of magnitude smaller than the conform ational dependence of the P otential Energy Surface V in the worst case, if chemical accuracy (typically de ned in the eld of ab initio quantum chemistry as 1 kcal/m ol [81]) is sought, they may be relevant. In fact, they are of the order of magnitude of the di erences between the energy surface V , F_s and F_r calculated at M P2/6-31++G (d_rp) and the ones calculated at HF/6-31++G (d_rp).

For the same reasons, we may conclude that, if ab initio derived potentials are used to carry out M olecular D ynam ics simulations of peptides, the Fixm an's compensating potential V_F should be included. Finally, regarding the relative importance of the dimension correcting terms TS_s^k , TS_s^c and TS_r^k , the results in table 4 suggest that the less important one is the kinetic entropy TS_s^k of the stimulation case (related to the determ inant of the mass-metric tensor G) and that the most important one is the one related to the determ inant of the H essian matrix H of the constraining part of the potential, i.e., the conformational entropy

 TS_s^k . The second usion is in agreement with the approximations typically made in the literature, the second one, however, is not (see the Appendix).





Figure 2: Potential Energy Surface (PES) of the model dipeptide $HCO-L-A la-N H_2$, computed at the M P2/6-31+ + G (d,p) level of the theory. The surface has been referenced to zero and sm oothed with bicubic splines for visual convenience. The units in the z-axis are kcal/m ol.



F igure 3: R am achandran plots of the correcting term s appearing in eqs.2.12 and 2.23, together with the F ixm an's compensating potential de ned in eq.2.25, computed at the M P2/6-31++G (d,p) level of the theory in the model dipeptide H CO -L-A la-N H₂. The surfaces have been referenced to zero and sm oothed with bicubic splines for visual convenience. The units in the z-axes are kcal/m ol.

Now, although the relative sizes of the conform ational dependence of the di erent term s m ay be indicative of their in portance, the degree of correlation am ong the surfaces is also relevant (see table 6). Hence, in order to arrive to m ore precise conclusions, we reexam ine here the results using a physically m eaningful criterium to com pare potential energy functions that has been introduced in ref. 82. The distance, denoted by d_{12} , between any two di erent potential energy functions, V_1 and V_2 , is an statistical quantity that, from a working set of conform ations (in this case, the 144 points of the grid), m easures the typical error that one m akes in the energy di erences if V_2 is used instead of V_1 , adm itting a linear rescaling.

In table 5, which contains the central results of this work, the distances between some of the energy surfaces that play a role in the problem are shown. We present the result in units of RT (at 300° K, where RT ' 0.6 kcal/mol) because it has been argued in ref. 82 that, if the distance between two di erent approxim ations of the energy of the same system is less than RT, one may safely substitute one by the other without altering the relevant physical properties. Moreover, if one assumes that the elective energies compared will be used to construct a polypeptide potential and that it will be designed as simply the sum of mono-residue ones (making each term suitably depend on di erent pairs of R am achandran angles), then, the number N res of residues up to which one may go keeping the distance between the two approximations of the the N-residue potential below R T is (see eq. 23 in ref. 82):

$$N_{res} = \frac{RT}{d_{12}}^2$$
: (4.1)

This number is also shown in table 5, together with the slope b_{12} of the linear rescaling between V_1 and V_2 and the Pearson's correlation ∞ cient [83], denoted by r_{12} .

The results at both MP2/6-31++G (d,p) and HF/6-31++G (d,p) levels of the theory are presented. The rst three rows in each of the rst two blocks are related to the classical still model, the next row to the classical rigid model and the last one in each block to the comparison between the two models. The third block in the table is associated to the comparison between the two dilerent levels of the theory used.

The F_s vs. V row (in the rst two blocks) assess the in portance of the two correcting terms, TS_s^k and TS_s^c , in the sticase. The result $d_{12} = 0.74R T$ indicates that, for the alanine dipeptide, V may be used as an approximation of F_s with caution if accurate results are sought. In fact, the low value of $N_{res} = 1.82 < 2$ shows that, if we wanted to describe a 2-residue peptide om itting the sticorrecting terms, we would typically make an error greater than the therm allows in the energy differences. The next two rows investigate the e ect of each one of the individual correcting terms. The conclusion that can be extracted from them (as the relative sizes in table 4 already suggested) is that the conformational entropy associated to the determinant of the H essian matrix H is much more relevant than the correcting term TS_s^k , related to the

Corr.ª	V_1^{b}	V2 ^c	d_{12}^{d}	N res ^e	b_{12}^{f}	r ₁₂ g		
	MP2/6-31++G(d,p)							
TS ^k TS ^c TS ^c TS ^k s	Fs Fs Fs	V V TS ^k s V TS ^c s		1.82 1.83 80.45	0.98 0.98 1.00	0.9967 0.9967 0.9999		
TSr	Fr	V	0.29 R T	11.62	1.01	0.9995		
V _F	Fs	Fr	0.67 R T	2,24	0.97	0.9972		
	Н F / 6-31+ + G (d,р)							
TS ^k TS ^c TS ^c TS ^k s	Fs Fs Fs	V V TS ^k _s V TS ^c _s		1.90 2.00 90.99	0.99 0.99 1.00	0.9975 0.9976 0.9999		
TSr	Fr	V	0.26 R T	14.83	1.01	0.9997		
V _F	Fs	Fr	0 . 61 R T	2.69	0.98	0.9982		
MP2/6-31++G (d,p) vs.HF/6-31++G (d,p)								
	V F _s F _r	V Fs Fr	1.25 R T 1.18 R T 1.18 R T	0.64 0.72 0.72	1.12 1.11 1.12	0.9925 0.9934 0.9932		

Table 5: C om parison of di erent energy surfaces involved in the study of the constrained equilibrium of the protected dipeptide $HCO-L-A la-NH_2$. ^aC orrecting term whose in portance is measured in the corresponding row, ^breference potential energy V₁ (the \correct" one, the one containing the correcting term), ^capproxim ated potential energy V₂ (i.e, V₁ m inus the correcting term in colum n a), ^d statistical distance between V₁ and V₂ (see A lonso J. L. and Echenique P., J. C om p. Chem. 27 (2006) 238{252}, ^em axim um number of residues in a polypeptide potential up to which the correcting term in colum n a may be om itted, ^f slope of the linear rescaling between V₁ and V₂ and ^gP earson's correlation coe cient. A ll quantities are dimensionless, except for d₁₂ which is given in units of the therm al energy RT at 300° K.

m ass-m etric tensor G, allowing to drop the latter up to 80 residues (according to M P 2/6-31++G (d_p) calculations). As has been already remarked, this second conclusion is in agreement with the approximations frequently done in the literature; how ever, it turns out that the importance of the Hessian-related term has been persistently underestimated (see the Appendix for a discussion).

The F_r vs. V row, in turn, shows the data associated to the kinetic entropy term TS_r^k , which is related to the determ inant of the reduced mass-metric tensor g in the classical rigid model. From the results there ($d_{12} = 0.29$ RT and N res = 11:62 at the M P 2/6-31++G (d_{2p}) level), we can conclude that the only

V_1^{a}		V_2^{b}	r ₁₂ °				
	MP2/6-31++G(d,p)						
V	vs.	TSsc	0.1572				
V	vs.	TSsk	-0.0008				
V	vs.	TSr	-0.3831				
V	VS.	V _F	0.3334				
	HF/6-31++G (dp)						
V	VS.	TSsc	0.0682				
V	vs.	TSsk	0.0897				
V	vs.	ΤSr	-0.3544				
V	VS.	V _F	0,2404				
MP2/6-31++G (d,p) vs. HF/6-31++G (d,p)							
ΤS _s ^c	vs.	ΤS ^c	0.9136				
ΤS ^k s	vs.	ΤS ^k s	0.9808				
ΤSr	vs.	ΤSr	0.9316				
V _F	VS.	V _F	0.9217				

Table 6: Correlation between the di erent correcting terms involved in the study of the constrained equilibrium of the protected dipeptide $HCO-L-A la-N H_2$. ^aR efference potential energy, ^bapproxim ated potential energy, ^cP earson's correlation coe cient.

correction term in the rigid case is less in portant than the ones in the sti case and that V $\,$ m ay be used as an approximation of F $_r$ for oligopeptides of up to 12 residues.

The last row in each of the rst two blocks in table 5 is related to the interesting question in M olecularD ynam ics of whether or not one should include the Fixm an's compensating potential V_F (see eq. 2.25) in rigid simulations in order to obtain the sti equilibrium distribution, exp (F_s), instead of the rigid one, exp (F_r). This question is equivalent to asking whether or not F_r is a good approximation of F_s . From the results in the table, we can conclude that the Fixm an's potential is relevant for peptides of more than 2 residues and its om ission may cause an error greater than the thermal noise in the energy di erences.

The appreciable sizes of the di erent correcting terms, shown in table 4, together with their low correlation with the Potential Energy Surface V, presented in the rst two blocks of table 6, explain their considerable relevance discussed in the preceding paragraphs.

M oreover, from the comparison of the M P2/6-31++G (d,p) and the HF/6-31++G (d,p) blocks, one can tell that the study herein perform ed m ay well have been done at the lower level of the theory (if we had known) with a tenth of the computationale ort (see sec. 3). This fact, explained by the high correlation, presented in the third block of table 6, between the correcting terms calculated at the two levels, is very relevant for further studies on more complicated dipeptides or longer chains and it indicates that the di erences in size between the di erent correcting terms s at M P 2/6-31++G (d,p) and HF/6-31++G (d,p), which are presented in table 4, are mostly due to a harm less linear scaling effect sim ilar to the well-known empirical scale factor frequently used in ab initio vibrational analysis [84{86]. This view is supported by the data in the third block of table 5, related to the comparison between the energy surfaces calculated at M P 2/6-31++G (d,p) and HF/6-31++G (d,p), where the slopes b_{12} are consistently larger than unity.

A last conclusion that may be extracted from the block labeled $MP2/6-31++G(d_p)$ vs. $HF/6-31++G(d_p)$ " in table 5 is that the typical error in the energy di erences (given by the distances d_{12}) produced when one reduces the level of the theory from $MP2/6-31++G(d_p)$ to $HF/6-31++G(d_p)$ is comparable (less than twice) to the error made if the most important correcting terms of the classical constrained models studied in this work are dropped. This is a useful hint for researchers interested in the conform ational analysis of peptides with quantum chem istry methods [60{64,70,87} and also to those whose aim is the design and parametrization of classical force elds from ab initio quantum mechanical calculations [68{70}].

-helix	-57	-47
3 ₁₀ -helix	-49	-26
-helix	-57	-70
polyproline II	-79	149
parallel -sheet	-119	113
antiparallel -sheet	-139	135

Table 7: Ram achandran angles (in degrees) of som e important secondary structure elements in polypeptides. Data taken from Lesk A.M., Introduction to Protein Architecture, Oxford University Press, Oxford, 2001.

F inally, in order to enrich and qualify the analysis, a new working set of conformations, di erent from the 144 points of the grid in the Ram achandran space, have been selected and the whole study has been repeated on them. These new conformations are six important secondary structure elements which form repetitive patterns stabilized by hydrogen bonds in polypeptides. Their conventional names and the corresponding values of the and angles have been taken from ref. 88 and are shown in table 7.

In g.4, the relative energies of these conform ations are shown for the three relevant potentials, V , F_s and F_r , at both MP2/6-31++G (d,p) and HF/6-31++G (d,p) levels of the theory. Since the antiparallel -sheet is the structure




Figure 4: Relative energies of som e important elements of secondary structure for the three potentials V , F_s and F_r , in the model dipeptide HCO-L-A la-NH₂ and at both MP2/6-31++G (d,p) and HF/6-31++G (d,p) levels of the theory. The energy of the antiparallel -sheet has been taken as reference. The units are kcal/m ol.

with the minimum energy in all the cases, it has been set as the reference and the rest of energies in the gure should be regarded as relative to it.

The meaningful assessment, using the statistical distance described above, of the typical error made in the energy di errences has been also performed on this new working set of conformations. The results are presented in table 8.

The distances between the free energies, $F_{\rm s}$ and $F_{\rm r}$, and their corresponding approximations obtained dropping the correcting entropies, $T\,S_{\rm s}^{\rm k}$, $T\,S_{\rm s}^{\rm c}$ and

T S_r^k, or the Fixm an's compensating potential V_F, in the rst two blocks of the table, are consistently sm aller than the ones found in the study of the grid de ned in the whole Ram achandran space (cf. table 5). And so are the distances between the three relevant potentials, V , F_s and F_r, calculated at the M P 2/6-31++G (d_xp) and H F/6-31++G (d_xp) levels of the theory.

A lthough the distance d_{12} used is a statistical quantity and, therefore, one must be cautious when working with such a small set of conformations (of size six, in this case), the conclusion drawn from this second part of the study is that, if one is interested only in the \bwer region" of the R am achandran surface, where the typical secondary structure elements lie, then, one may safely neglect the conformational dependence of the di erent correcting terms appearing in the study of the constrained equilibrium of peptides. At least, up to oligopeptides (poly-alanines) of 10 residues in the worst case (the neglect of the F ixm an's

Corr.ª	V_1^{b}	V2 ^c	d_{12}^{d}	N res e	b_{12}^{f}	r ₁₂ g		
	MP2/6-31++G(d ₄ p)							
TS ^k TS ^c TS ^c TS ^k s	Fs Fs Fs	V V TS ^k s V TS ^c s		19.72 14.07 298.13	0.99 0.98 1.01	0.9990 0.9985 0.9999		
TSr	Fr	V	0.20 R T	25.64	0.99	0.9992		
V _F	Fs	Fr	0.34 R T	8.73	0.99	0.9977		
		HF/6-31+	+G (d,p)					
TS ^k TS ^c TS ^c TS ^k s	Fs Fs Fs	V V TS ^k s V TS ^c s		47.94 46.12 380.30	1.00 1.00 1.00	0.9997 0.9997 0.9999		
TSr	Fr	V	0.15 R T	41.85	0.99	0.9997		
V _F	Fs	Fr	0.18 R T	30.12	1.01	0.9996		
MP2/6-31++G(d,p)vs.HF/6-31++G(d,p)								
	V Fs Fr	V Fs Fr	0.77 R T 0.77 R T 0.71 R T	1.68 1.69 1.96	128 126 128	0.9929 0.9928 0.9939		

Table 8: C om parison of di erent approximations to the energies of some important elements of secondary structure (see table 7) in the study of the constrained equilibrium of the protected dipeptide $H C O - L - A la - N H_2$. See the caption of table 5 for an explanation of the keys in the di erent columns.

com pensating potential V $_{\rm F}$ in the F $_{\rm s}$ vs. F $_{\rm r}$ com parison at M P 2/6–31+ + G (d,p)).

This di erence between the two working set of conform ations may be explained looking at one of the ways of expressing the statistical distance used (see eq. 12a in ref. 82):

$$d_{12} = {\stackrel{p}{\underline{-}}}_{2} {}_{2} (1 { r}_{12}^{2})^{1=2}; \qquad (4.2)$$

where r_{12} is the Pearson's correlation coe-cient between the potential energies denoted by V_1 and V_2 and $_2$ is the standard deviation in the values of V_2 on the relevant working set of conform ations.

This last quantity, $_2$, is the responsible of the di erences between tables 5 and 8, since the set of conform ations comprised by the six secondary structure elements in table 7 spans a smaller energy range than the whole P otential E nergy Surface in g.2 (or F_s , or F_r , which have very similar variations). A coordingly, the dispersion in the energy values is smaller: $_2$ ' 2 kcal/m ol in the case of the secondary structure elements and $_2$ ' 4 kcal/m ol for the grid in the whole

R am achandran space (see table 4). Since the correlation coe cient in both cases are of sim ilar magnitude, the di erences in $_2$ produce a smaller distance d₁₂ for the second set of conformations studied, i.e., a smaller typical error made in the energy di erences when om itting the correcting terms derived from the consideration of constraints.

To end this section, we remark that, although this \lower region" of the R am achandran space contains the most relevant secondary structure elements (which are also the most commonly found in experimentally resolved native structures of proteins [89{92]) and may be the only region explored in the dynamical or thermodynamical study of small peptides, if the aim is the design of e ective potentials for computer simulation of polypeptides [68{70}], then, some caution is recommended, since long-range interactions in the sequence may tem porarily compensate local energy penalizations and the higher regions of the energy surfaces studied could be important in transition states or in some relevant dynam ical paths of the system.

In the following section, the many results discussed in the preceding paragraphs are summarized.

5 Conclusions

In this work, the theory of classical constrained equilibrium has been collected for the sti and rigid models. The pertinent correcting terms, which may be regarded as elective entropies, as well as the Fixm an's compensating potential, have been derived and theoretically discussed (see eqs. 2.12, 2.23 and 2.25, together with the form ulae in sec. 3.1). In addition, the common approximation of considering that, for typical internals, the equilibrium values of the hard coordinates do not depend on the soft ones, has also been discussed and related to the rest of simplications. The treatment of the assumptions in the literature is thoroughly reviewed and discussed in the Appendix.

In the central part of the work (sec. 4), the relevance of the di erent correcting term s has been assessed in the case of the model dipeptide HCO-L-A la- NH_2 , with quantum mechanical calculations including electron correlation. A lso, the possibility of perform ing analogous studies at the less dem anding H artree Fock level of the the theory has been investigated. The results found are sum marized in the follow ing points:

In M onte Carb simulations of the classical stim odel at room temperature, the elective entropy TS_s^k , associated to the determinant of the massmetric tensor G, may be neglected for peptides of up to 80 residues. Its maximum variation in the Ramachandran space is 0.24 kcal/mol.

In M onte C arb simulations of the classical stim odel at room temperature, the elective entropy TS_s^c , associated to the determinant of the Hessian H of the constraining part of the potential, should be included for peptides of m ore than 2 residues. Its maximum variation in the Ra-machandran space is 1.67 kcal/m ol.

In M onte C arb simulations of the classical rigid m odel at room tem – perature, the elective entropy TS_r^k , associated to the determinant of the reduced m ass-metric tensor g, m ay be neglected for peptides of up to 12 residues. Its maximum variation in the R am achandran space is 0.81 kcal/mol.

In rigid M olecular D ynam ics simulations intended to yield the sti equilibrium distribution at room temperature, the Fixman's compensating potential V_F should be included for peptides of more than 2 residues. Its maximum variation in the Ram achandran space is 1.68 kcal/m ol.

If the assumption that only the more stable region of the Ramachandran space, where the principal elements of secondary structure lie, is relevant, then, the importance of the correcting terms decreases and the limiting number of residues in a polypeptide potential up to which they may be om itted is approximately four times larger in each of the previous points.

In both cases (i.e., either if the whole R am achandran space is considered relevant, or only the lower region), the errors m ade if the most important correcting terms are neglected are of the same order of m agnitude as the errors due to a decrease in the level of theory from $M P 2/6-31++G (d_{a}p)$ to $H F / 6-31++G (d_{a}p)$.

The whole study of the relevance of the di erent correcting term s (or future analogous investigations) m ay be performed at the HF/6-31++G (d,p) level of the theory, yielding very similar results to the ones obtained at MP2/6-31++G (d,p) and using a tenth of the computational e ort.

To end this discussion, some quali cations should be made. On one hand, the conclusions above refer to the case in which a classical potential directly extracted from the quantum mechanical (B om-O ppenheimer) one is used; for the considerably simpler force elds typically used for macrom olecular simulations, the study should be repeated and di erent results may be obtained. On the other hand, the investigation performed in this work has been done in one of the simplest dipeptides; both its isolated character and the relatively sm all size of its side chain play a role in the results obtained. Hence, for bulkier residues included in polypeptides, these conclusions should be approached with caution and much interesting work remains to be done.

A cknow ledgm ents

W e would like to thank F.Falceto and V.Laliena for illum inating discussions and also the reviewers of the manuscript for much useful suggestions. The num erical calculations have been performed at the BIFI com puting facilities. We thank I.Campos, for the invaluable CPU time and the e ciency at solving the problem s encountered.

This work has been supported by the A ragon G overnm ent (B iocom putacion y F sica de Sistem as C om plejos" group) and by the research grants M EC (Spain) F IS2004-05073 and FPA 2003-02948, and M CYT (Spain) BFM 2003-08532. P. E chenique and I.Calvo are supported by M EC (Spain) FPU grants.

Appendix

M any approximations may be done to simplify the calculation of the di erent correcting terms introduced in the previous subsections. The most frequently found in the literature are the following three:

- (i) To neglect the conform ational dependence of detG.
- (ii) To neglect the conform ational dependence of detH .
- (iii) To assume that the hard coordinates are constant, i.e., that the $f^{I}(q^{i})$ in eq.2.1 do not depend on the soft coordinates q^{i} .

The conform ational dependence of detg is custom arily regarded as in portant since it was shown to be non-negligible even for simple systems some decades ago [13,21{23} (norm ally in an indirect way, while studying the in uence of the F ixm an's compensating potential in eq. 2.25; see discussion below). W ith this same aim, P atriciu et al. [20] have very recently measured the conform ational dependence of detg for a serial polymer with xed bond lengths and bond angles (in the approximation (iii)), show ing that it is non-negligible and suggesting that it may be so also for more general system s.

Note that, if approximations (i) and (ii) are assumed, then the Fixman's potential depends only on detg. In fact, whereas in the general case the Fixman's compensating potential cannot be simplied beyond the expression in eq. 2.25, if one assumes approximation (iii), then the reduced mass-metric tensor g turns out to be the subblock of G with soft indices and, in this case, the quotient detG = detg has been shown to be equal to 1=deth by Fixman [16], where h denotes the subblock of G ¹ with hard indices, i.e.,

$$h^{IJ}(q) \coloneqq \frac{X^{N}}{e} \frac{\partial q^{I}}{\partial x} \frac{1}{m} \frac{\partial q^{J}}{\partial x} : \qquad (5.1)$$

This result has been extensively used in the literature $[21\{23,38,42,57]$, since each of the internal coordinates q^a typically used in m acrom olecular simulations only involves a sm all num ber of atom s, thus rendering the m atrix h above sparse and allow ing for e cient algorithm s to be used in order to nd its determ inant.

Now, although detg is custom arily regarded as important, the conform ational variations of detG are alm ost unanim ously neglected (approximation (i)) in the literature [15,55] and may only be said to be indirectly included in h by the authors that use the expression above $[20\{23,38,57]$. This is mainly due to the fact, reported by G o and Scheraga [15] and, before, by Volkenstein [74], that detG in a serial polymer may be expressed as in eq.3.1, being independent of the dihedral angles (which are custom arily taken as the soft coordinates). If one also assumes approximation (iii), which, as will be discussed later, is very common, then detG is a constant for every conformation of the molecule.

P robably due to com putational considerations, but also som etim es to the use of a form ulation of the sti case based on delta functions [51], the conform ational dependence of det H is alm ost unanim ously neglected (approxim ation (ii)) in the literature [15,16,19,20,41,55,65,66]. Only a few authors include this term in di erent stages of the reasoning [13{15,18,19,26,39},m ost of them only to argue later that it is negligible.

A lthough for som e simple ad hoc designed potentials that lack long-range term s [21,22,57], the aforem entioned simplifying assumptions and the ones that will be discussed in the following paragraphs may be exactly fulled, in the case of the potential energies used in force elds for macrom olecular simulation $[6,27\{37]$, they are not. The typical energy function in this case, has the form

$$V (q^{a}) := \frac{1}{2} \sum_{i=1}^{M'r} K_{r} (r r^{0})^{2} + \frac{1}{2} \sum_{i=1}^{M} K_{r} (r^{0})^{2} + \frac{1}{2} \sum_{i=1}^{M} K_{r} ($$

where r are bond lengths, are bond angles, are dihedral angles and, for the sake of sim plicity, no harm onic terms have been assumed for out-of-plane angles or for hard dihedrals (such as the peptide bond !). N_r is the number of bond lengths, N the number of bond angles and the quantities K_r, K, r⁰ and ⁰ are constants. The term denoted by V^{tors}() is a commonly included torsional potential that depends only on the dihedral angles and V^{long range} (q^a) norm ally comprises long-range interactions such as C oulom b or van der W aals; hence, it depends on the atom ic positions x⁰ which, in turn, depend on all the internal coordinates q^a.

O ne of the reasons given for neglecting detH, when classical force elds are used with potential energy functions such as the one in eq. 52, is that the harm onic constraining terms dominate over the rest of interactions and, since the constants appearing on these terms (the K $_{\rm r}$, K in eq. 52) are independent of the conform ation by construction, so is detH [15,19,26]. Here, we analyze a more realistic quantum -mechanical potential and these considerations are not applicable, however, they also should be checked in the case of classical force elds, since, for a potential energy such as the one in eq. 52), the quantities K $_{\rm r}$ and K are nite and the long-range terms will also affect the H essian at each point of the constrained hypersurface , rendering its determ inant conform ation-dependent.

For the same reason, even in classical force elds, the equilibrium values of the hard coordinates are not the constant quantities r^0 and 0 in eq. 5.2 but some functions $f^{I}(q^{i})$ of the soft coordinates (see eq. 2.1). This fact, recognized

by som e authors [15,25,45,93], provokes that, if one chooses to assum e approximation (iii) and the constants r^0 and 0 appearing in eq.5.2 are designated as the equilibrium values, the potential energy in may be heavily distorted, the cause being simply that the long-range interactions between atoms separated by three covalent bonds are not fully relaxed [93]. This e ect is probably larger ifbond angles, and not only bond lengths, are also constrained, which may partially explain the di erent dynam ical behaviour found in ref. 6 when com paring these types of constraints in M olecular D ynam ics sim ulations. In quantum mechanical calculations of sm all dipeptides, on the other hand, the fact that the bond lengths and bond angles depend on the R am achandran angles (;) has been pointed out by Scha er et al. [94]. Therefore, approximation (iii), which is very com m on in the literature [6,13,15,16,18 {20,26,38 {42,51,52,55,65,66,95,96], should be critically analyzed in each particular case.

Apart from the typical internal coordinates q^a used until now, in terms of which the constrained hypersurface is described by the relations $q^{\rm I} = f^{\rm I} \, (q^i)$ in eq.2.1, with I = M + 7;:::;N, one may de nead i erent set Q ^a such that, on , the corresponding hard coordinates are arbitrary constants Q $^{\rm I} = C^{\rm I}$ (the external coordinates $q^{\rm A}$ and Q $^{\rm A}$ are irrelevant for this part of the discussion). To do this, for example, let

$$Q^{i} \coloneqq q^{i} \qquad i = 7; \dots; M + 6 \quad \text{and} \\ Q^{I} \coloneqq q^{I} \quad f^{I}(q^{i}) + C^{I} \quad I = M + 7; \dots; N :$$

$$(5.3)$$

W ellthen, while the relation between bond lengths, bond angles and dihedral angles (the typical q^a [59]) and the Euclidean coordinates is straightforward and simple, the expression of the transform ation functions $Q^a(x)$ needs the know ledge of the f^I, which must be calculated num erically in most real cases. This drastically reduce the practical use of the Q^a, how ever, it is also true that they are conceptually appealing, since they have a property that closely m atch our intuition about what the soft and hard coordinates should be (nam ely, that the hard coordinates Q^{I} are constant on the relevant hypersurface); and this is why we term them exactly separable hard and soft coordinates. Now, we must also point out that, although the real internal coordinates q^a do not have this property, they are usually close to it. The custom ary labeling of soft and hard coordinates in the literature is based on this circum stance. Som ehow, the dihedral angles are the \softest" of the internal coordinates, i.e., the ones that \vary the most" when the system visits di erent regions of the hypersurface ; and this is why we term the real q^a approximately separable hard and soft coordinates, considering approximation (iii) as a useful reference case.

To sum up, the three sim plifying assumptions (i), (ii) and (iii) in the beginning of this section should be regarded as approximations in the case of classical force elds, as well as in the case of the more realistic quantum -mechanical potential investigated in this work, and they should be critically assessed in the system s of interest. Here, while studying the model dipeptide $HCO - L - A la - N H_2$, no sim plifying assumptions of this type have been made.

References

- [1] J.L.A lonso, G.A.Chass, I.G.Csizm adia, P.E chenique, and A.Tarancon. Do theoretical physicists care about the protein folding problem? In R.F. A lvarez-E strada et al., editors, M eeting on Fundam ental Physics A lberto G alindo'. A ula D ocum ental, M adrid, 2004. (arX iv q-bio BM /0407024).
- [2] C.M. Dobson. Protein folding and m isfolding. Nature, 426:884 (890, 2003.
- [3] K.A.D ill. Polym erprinciples and protein folding. Prot. Sci., 8:1166{1180, 1999.
- [4] S.He and H.A. Scheraga. Brownian dynamics simulations of protein folding. J. Chem. Phys., 108:287, 1998.
- [5] R.A.Abagyan, M.M. Totrov, and D.A.Kuznetsov. ICM: A new method for protein modeling and design: Applications to docking and structure prediction from the distorted native conformation. J.Comp.Chem., 15:488{ 506, 1994.
- [6] W.F.Van Gunsteren and M.Karplus. E ects of constraints on the dynamics of macrom olecules. Macrom olecules, 15:1528 (1544, 1982.
- [7] C. Levinthal. How to fold graciously. In J.T.P.DeB runner and E.M unck, editors, M ossbauer Spectroscopy in Biological Systems, pages 22{24, A llerton House, M onticello, Illinois, 1969. University of Illinois Press.
- [8] H.M.Chun, C.E.Padilla, D.N.Chin, M.W atanabe, V.I.Karlov, H.E. Alper, K.Soosaar, K.B.Blair, O.M.Becker, L.S.D.Caves, R.Nagle, D.N.Haney, and B.L.Farmer. MBO (N)D: A multibody method for long-time M olecular D ynam ics simulations. J.Comp.Chem., 21:159{184, 2000.
- [9] S.Reich.Sm oothed Langevin dynam ics of highly oscillatory system s. Physica D, 118:210{224,2000.
- [10] S.Reich. Multiple tim e scales in classical and quantum -classical molecular dynamics. J. Comput. Phys., 151:49{73, 1999.
- [11] T. Schlick, E. Barth, and M. Mandziuk. Biom olecular dynamics at long tim esteps: Bridging the tim escale gap between simulation and experim entation. Annu. Rev. Biophys. Biom ol. Struct., 26:181{222,1997.
- [12] N.G. Van Kampen and J.J.Lodder. Constraints. Am. J. Phys., 52:419 424, 1984.
- [13] J.M. Rallison. The role of rigidity constraints in the rheology of dilute polymer solutions. J. Fluid Mech., 93:251 (279, 1979.
- [14] E.Helfand.Flexible vs. rigid constraints in Statistical Mechanics. J. Chem. Phys., 71:5000, 1979.

- [15] N.Go and H.A. Scheraga. On the use of classical statistical mechanics in the treatment of polymer chain conformation. Macromolecules, 9:535, 1976.
- [16] M. Fixman. Classical Statistical Mechanics of constraints: A theorem and application to polymers. Proc. Natl. Acad. Sci. USA, 71:3050{3053,1974.
- [17] N. Go and H. A. Scheraga. A nalysis of the contributions of internal vibrations to the statistical weights of equilibrium conformations of macromolecules. J. Chem. Phys., 51:4751, 1969.
- [18] D.C.Morse. Theory of constrained B rownian motion. Adv. Chem. Phys., 128:65{189,2004.
- [19] W.K.Den Otter and W.J.Briels. Free energy from molecular dynamics with multiple constraints. Mol. Phys., 98:773 (781, 2000.
- [20] A. Patriciu, G. S. Chirik jian, and R. V. Pappu. Analysis of the conform ational dependence of m ass-m etric tensor determ inants in serial polymers with constraints. J. Chem. Phys., 121:12708 {12720, 2004.
- [21] D. Perchak, J. Skolnick, and R. Yaris. Dynamics of rigid and exible constraints for polymers. E ect of the Fixm an potential. M acrom olecules, 18:519{525, 1985.
- [22] M.R.Pear and J.H.W einer. Brownian dynam ics study of a polymer chain of linked rigid bodies. J. Chem. Phys., 71:212, 1979.
- [23] D. Chandler and B. J. Berne. Comment on the role of constraints on the conform ational structure of n-butane in liquid solvent. J. Chem. Phys., 71:5386{5387,1979.
- [24] M. Gottlieb and R. B. Bird. A Molecular Dynamics calculation to conm the incorrectness of the random -walk distribution for describing the K ram ers freely jointed bead-rod chain. J. Chem. Phys., 65:2467, 1976.
- [25] J. Zhou, S. Reich, and B. R. Brooks. Elastic molecular dynamics with self-consistent exible constraints. J. Chem. Phys., 111:7919, 2000.
- [26] H.J.C.Berendsen and W.F.Van Gunsteren. Molecular Dynamics with constraints. In J.W. Perram, editor, The Physics of Superionic Conductors and Electrode Materials, volume NATO ASI Series B92, pages 221{240. Plenum Press, 1983.
- [27] A. D. MacKerell Jr., B. Brooks, C. L. Brooks III, L. Nilsson, B. Roux, Y. W on, and M. Karplus. CHARMM: The energy function and its param – eterization with an overview of the program. In P.v.R. Schleyer et al., editors, The Encyclopedia of Computational Chemistry, pages 217{277. John Wiley & Sons, Chichester, 1998.

- [28] B.R.Brooks, R.E.Bruccoleri, B.D.Olafson, D.J.States, S.Swam inathan, and M.Karplus. CHARMM: A program form acrom olecular energy, m inim ization, and dynam ics calculations. J. Comp. Chem., 4:187{217, 1983.
- [29] W. D. Cornell, P. Cieplak, C. I. Bayly, I.R. Gould, Jr. Merz, K. M., D. M. Ferguson, D. C. Spellmeyer, T. Fox, J.W. Caldwell, and P.A. Kollman. A second generation force eld for the simulation of proteins, nucleic acids, and organic molecules. J. Am. Chem. Soc., 117:5179{5197, 1995.
- [30] D. A. Pearlman, D. A. Case, J. W. Caldwell, W. R. Ross, T. E. Cheatham III, S. DeBolt, D. Ferguson, G. Seibel, and P. Kollman. AM – BER, a computer program for applying molecular mechanics, norm alm ode analysis, molecular dynamics and free energy calculations to elucidate the structures and energies of molecules. Comp. Phys. Commun., 91:1{41, 1995.
- [31] W . L. Jorgensen and J. Tirado-Rives. The OPLS potential functions for proteins. Energy m inimization for crystals of cyclic peptides and Crambin. J. Am. Chem. Soc., 110:1657{1666, 1988.
- [32] W. L. Jorgensen, D. S. Maxwell, and J. Tirado-Rives. Development and testing of the OPLS all-atom force eld on conformational energetics and properties of organic liquids. J. Am. Chem. Soc., 118:11225{11236, 1996.
- [33] T. A. Halgren. Merck molecular force eld. I. Basis, form, scope, parametrization, and perform ance of MMFF94. J. Comp. Chem., 17:490 519, 1996.
- [34] T.A.Halgren.Merckmolecular force eld.II.MMFF94 van derWaals and electrostatica parameters for intermolecular interactions. J. Comp. Chem., 17:520{552,1996.
- [35] T.A.Halgren. M erck m olecular force eld. III. M olecular geom etrics and vibrational frequencies for M M FF 94. J. Com p. Chem., 17:553 (586, 1996.
- [36] T.A.Halgren. Merck molecular force eld. IV. Conformational energies and geometries for MMFF94. J. Comp. Chem., 17:587{615, 1996.
- [37] T.A.Halgren.M erck m olecular force eld.V.Extension of MMFF94 using experim ental data, additional computational data, and empirical rules. J. Comp.Chem., 17:616{641,1996.
- [38] M. Pasquali and D. C. Morse. An e-cient algorithm for metric correction forces in simulations of linear polymers with constrained bond lengths. J. Chem. Phys., 116:1834, 2002.
- [39] W.K.Den Otter and W.J.Briels. The calculation of free-energy di erences by constrained m olecular-dynam ics simulations. J. Chem. Phys., 109:4139, 1998.

- [40] G. Ciccotti and J. P. Ryckaert. M olecular dynamics simulation of rigid m olecules. Comput. Phys. Rep., 4:345{392, 1986.
- [41] H.J.C.Berendsen and W.F.Van Gunsteren. Molecular Dynamics sin ulations: Techniques and approaches. In A.J. et al. Barnes, editor, Molecular Liquids-Dynamics and Interactions, pages 475 (500. Reidel Publishing Company, 1984.
- [42] M. Fixm an. Simulation of polymer dynamics. I. General theory. J. Chem. Phys., 69:1527, 1978.
- [43] R.F.A lvarez-E strada and G.F.Calvo. M odels for biopolymers based on quantum mechanics. M ol. Phys., 100:2957{2970,2002.
- [44] R.F.A lvarez-Estrada.M odels of macrom olecular chains based on C lassical and Q uantum M echanics: comparison with G aussian models. M acromol. Theory Simul, 9:83{114,2000.
- [45] B. Hess, H. Saint-Martin, and H. J. C. Berendsen. Flexible constraints: An adiabatic treatment of quantum degrees of freedom, with application to the exible and polarizable mobile charge densities in harmonic oscillators model for water. J. Chem. Phys., 116:9602, 2002.
- [46] M.Bom and J.R.Oppenheim er. ZurQuantentheorie der Molekeln. Ann. Phys. Leipzig, 84:457 (484, 1927.
- [47] E.B.W ilson Jr., J.C.Decius, and P.C.Cross. M olecular V ibrations: The Theory of Infrared and Ram an V ibrational Spectra. Dover Publications, New York, 1980.
- [48] E. Barth, K. Kuczera, B. Leim kuhler, and R. D. Skeel. A lgorithm s for constrained M olecular D ynam ics. J. Comp. Chem., 16:1192{1209, 1995.
- [49] H.C.Andersen. Rattle: A \velocity" version of the Shake algorithm for m olecular dynam ics calculations. J. Com put. Phys., 52:24{34, 1983.
- [50] J. P. Ryckaert, G. Ciccotti, and H. J. C. Berendsen. Numerical integration of the Cartesian equations of motion of a system with constraints: M olecular dynamics of n-alkanes. J. Comput. Phys., 23:327 [341, 1977.
- [51] J. Schlitter and M. K lan. The free energy of a reaction coordinate at multiple constraints: a concise formulation. Mol. Phys., 101:3439{3443, 2003.
- [52] W.F.Van Gunsteren.M ethods for calculation of free energies and binding constants: Successes and problem s. In W.F.Van Gunsteren and P.K. W einer, editors, C om puter Sim ulations of B iom olecular System s, pages 27{ 59.E scom science publishers, N etherlands, 1989.
- [53] A.R.D inner. Local deform ations of polymers with nonplanar rigid mainchain coordinates. J.Comp.Chem., 21:1132{1144,2000.

- [54] J.Scho eld and M.A.Ratner.M onte Carlom ethods for short polypeptides. J.Chem. Phys., 109:9177, 1998.
- [55] A.J.Pertsin, J.Hahn, and H.P.G rossm ann. Incorporation of bond-lengths constraints in M onte C arb simulations of cyclic and linear m olecules: conform ational sampling for cyclic alkanes as test system s. J. C om p. C hem., 15:1121{1126,1994.
- [56] E.W. Knapp and A. Irgens-D efregger. O -lattice M onte C arlo m ethod w ith constraints: Long-tim e dynam ics of a protein m odel w ithout nonbonded interactions. J. F luid M ech., 14:19{29, 1993.
- [57] N.G.Almarza, E.Enciso, J.Alonso, F.J.Bermejo, and M.Alvarez. Monte Carlo simulations of liquid n-butane. Mol. Phys., 70:485{504, 1990.
- [58] P.Echenique and I.Calvo.Explicit factorization of external coordinates in constrained Statistical Mechanics models. To appear in J.Comp.Chem., 2006. (arX iv q-bioQM /0512033).
- [59] P.Echenique and J.L.Alonso. De nition of System atic, Approximately Separable and Modular Internal Coordinates (SASMIC) for macromolecular simulation. To be published in J.Comp.Chem., 2006. (arXiv:qbioBM /0511004).
- [60] A. Lang, I. G. C sizm adia, and A. Perczel. Peptide models. XLV: Conform ational properties of N-form yl-L-m ethioninam ide ant its relevance to m ethionine in proteins. PROTEINS: Struct. Funct. Bioinf., 58:571{588, 2005.
- [61] A. Perczel, O. Farkas, I. Jakli, I. A. Topol, and I. G. C sizm adia. Peptide m odels. XXX III. Extrapolation of low-level Hartree Fock data of peptide conform ation to large basis set SCF, MP2, DFT and CCSD (T) results. The R am achandran surface of alanine dipeptide computed at various levels of theory. J. Comp. Chem., 24:1026{1042, 2003.
- [62] R. Vargas, J. Garza, B. P. Hay, and D. A. Dixon. Conform ational study of the alanine dipeptide at the MP2 and DFT levels. J. Phys. Chem. A, 106:3213{3218,2002.
- [63] C.H.Yu, M.A.Norman, L.Schafer, M.Ramek, A.Peeters, and C.van Alsenoy. Ab initio conformational analysis of N-form yl L-alanine am ide including electron correlation. J. Mol. Struct., 567{568:361{374,2001.
- [64] A.G.C saszar and A.Perczel. Ab initio characterization of building units in peptides and proteins. Prog. B jophys. M ol. B jol., 71:243{309, 1999.
- [65] M.P.Allen and D.J.Tildesley. Computer simulation of liquids. Clarendon Press, Oxford, 2005.

- [66] D.Frenkel and Sm it B. Understanding molecular simulations: From algorithm s to applications. A cadem ic Press, O rlando FL, 2nd edition, 2002.
- [67] M.Kamplus and J.N.Kushick. M ethod for estimating the congurational entropy of macrom olecules. Macrom olecules, 14:325{332, 1981.
- [68] A.R.M acK erellJr., M.Feig, and C.L.B rooks III. Extending the treatment of backbone energetics in protein force elds: Limitations of gas-phase quantum mechanics in reproducing protein conformational distributions in molecular dynamics simulations. J.Comp.Chem., 25:1400{1415,2004.
- [69] A.J.Bordner, C.N.Cavasotto, and R.A.Abagyan.Direct derivation of van derW aals force eldsparameters from quantum mechanical interaction energies.J.Phys.Chem.B, 107:9601 {9609, 2003.
- [70] M. Beachy, D. Chasman, R. Murphy, T. Halgren, and R. Friesner. A ccurate ab initio quantum chemical determination of the relative energetics of peptide conformations and assessment of empirical force elds. J. Am. Chem. Soc., 119:5908 (5920, 1997.
- [71] V. I. A mold. M athem atical M ethods of C lassical M echanics. G raduate Texts in M athem atics. Springer, New York, 2nd edition, 1989.
- [72] B. Viskolcz, S. N. Fejer, and I. G. Csizmadia. Thermodynamic functions of conformational changes. 2. Conformational entropy as a measure of information accumulation. To be published in J. Phys. Chem. A., 2006. (ASAP article http://pubs.acs.org/cgibin/abstract.cgi/jcafh/asap/abs/jp058219t.htm l).
- [73] I. Andricioaei and M. Karplus. On the calculation of entropy from covariance matrices of the atom ic uctuations. J. Chem. Phys., 115:6289{6292, 2001.
- [74] M.V.Volkenstein.Con gurational Statistical of Polymeric Chains. Interscience, New York, 1959.
- [75] G.N.Ramachandran and C.Ramakrishnan. Stereochem istry of polypeptide chain con gurations. J.M ol. Biol., 7:95{99, 1963.
- [76] V. Hnizdo, A. Fedorowicz, H. Singh, and E. Dem chuk. Statistical therm odynamics of internal rotation in a hindering potential of mean force obtained from computer simulations. J. Comp. Chem., 24:1172 (1183, 2003.
- [77] E.Dem chuk and H.Singh. Statistical therm odynam ics of hindered rotation from computer simulations. Mol. Phys., 99:627{636, 2001.
- [78] K.V.M ardia and P.E.Jupp. D irectional Statistics. John W iley & Sons, Chichester, 2000.

- [79] M.W. Schmidt, K.K. Baldridge, J.A. Boatz, S.T. Elbert, M.S.Gordon, H.J. Jensen, S.Koseki, N.Matsunaga, K.A. Nguyen, S.Su, T.L.W indus, M. Dupuis, and J.A. Montgomery. General Atom ic and Molecular Electronic Structure System. J. Comp. Chem., 14:1347{1363, 1993.
- [80] J. Baker, A. Kessi, and B. Delley. The generation and use of delocalized internal coordinates in geometry optimization. J. Chem. Phys., 105:192{ 212, 1996.
- [81] I. P. Daykov, T. A. Arias, and T. D. Engeness. Robust ab initio calculation of condensed matter: Transparent convergence through sem icardinal multiresolution analysis. Phys. Rev. Lett., 90:216402, 2003.
- [82] J. L. A lonso and P. E chenique. A physically meaningful method for the comparison of potential energy functions. J. Comp. Chem., 27:238{252, 2006.
- [83] J. D. Dobson. Applied multivariate data analysis, volume I. Springer-Verlag, New York, 1991.
- [84] I.N. Levine. Quantum Chemistry. Prentice Hall, Upper Saddle River, 5th edition, 1999.
- [85] M.D.Halls, J.Velkovski, and H.B.Schlegel. Harmonic Frequency Scaling Factors for Hartree-Fock, S-VW N, B-LYP, B3-LYP, B3-PW 91 and MP2 and the Sadlej pVTZ Electric Property Basis Set. Theo. Chem. Acc., 105:413, 2001.
- [86] A.P.Scott and L.R adom .H arm onic vibrational frequencies: A n evaluation of Hartree-Fock, M ller-Plesset, Quadratic C on guration Interaction, D ensity Functional Theory, and sem iem pirical scale factors. J. Phys. Chem., 100:16502 {16513, 1996.
- [87] M. Elstner, K. J. Jalkanen, M. Knapp-M oham m ady, and S. Suhai. Energetics and structure of glycine and alanine based m odel peptides: A pproxim ate SCC-DFTB, AM 1 and PM 3 m ethods in comparison with DFT, HF and MP2 calculations. Chem. Phys., 263:203(219, 2001.
- [88] A.M. Lesk. Introduction to Protein Architecture. Oxford University Press, Oxford, 2001.
- [89] P. Chakrabarti and D. Pal. The interrelationships of side-chain and mainchain conformations in proteins. Prog. Biophys. Mol. Biol., 76:1{102,2001.
- [90] H.M. Berman, J.W estbrook, Z.Feng, G.G illiland, T.N. Bhat, H.W eissig, I.N. Shindyalov, and P.E.Bourne. The protein data bank. Nucleic Acids Research, 28:235{242, 2000.
- [91] K. Gunasekaran, C. Ramakrishnan, and P. Balaram. Disallowed Ramachandran conformations of amino acid residues in protein structures. J. M ol. Biol., 264:191{198,1996.

- [92] T.E.Creighton.Proteins: Structures and M olecular Properties. Freem an, W.H., New York, 2nd edition, 1992.
- [93] J.Chen, W. Im, and C.L.Brooks III. Application of torsion angle molecular dynam ics for e cient sampling of protein conformations. J. Comp. Chem., 26:1565{1578,2005.
- [94] L. Schafer and C. Ming. Predictions of protein backbone bond distances and angles from rst principles. J. Mol. Struct., 333:201 (208, 1995.
- [95] M. Mazars. Canonical partition function of freely jointed chains. J. Phys. A: Math. Gen., 31:1949{1964, 1998.
- [96] M. Mazars. Statistical physics of the freely jointed chain. Phys. Rev. E, 53:6297, 1996.