

# STRUCTURED THALAMOCORTICAL CONNECTIVITY REVEALED BY RANDOM WALKS ON COMPLEX NETWORKS

Luciano da Fontoura Costa

Instituto de Física de São Carlos, Universidade de São Paulo, Av. Trabalhador São Carlense 400,  
Caixa Postal 369, CEP 13560-970, São Carlos, São Paulo, Brazil

Olaf Sporns<sup>y</sup>

Department of Psychological and Brain Sciences,  
Indiana University, Bloomington, IN 47405, USA

(Dated: 5th Feb 2006)

The segregated regions of the mammalian cerebral cortex and thalamus form an extensive and complex network, whose structure and function are still only incompletely understood. The present article describes an application of the concepts of complex networks and random walks that allows the identification of non-random, highly structured features of thalamocortical connections, and their potential effects on dynamic interactions between cortical areas in the cat brain. Utilizing large-scale anatomical data sets of this thalamocortical system, we investigate uniform random walks in such a network by considering the steady state eigenvector of the respective stochastic matrix. It is shown that thalamocortical connections are organized in such a way as to guarantee strong correlation between the outdegree and occupancy rate (a stochastic measure potentially related to activation) of each cortical area. Possible organizational principles underlying this effect are identified and discussed.

The relationship between the topology of structural connection patterns and cortical dynamics currently represents a significant challenge to brain theory. Numerous neuroanatomical studies have revealed that the pathways of the mammalian thalamocortical system exhibit specific patterns ranging in scale from interconnections linking whole brain regions to intra-areal patterns of connections between cell populations or individual cortical neurons. Comprehensive descriptions of large-scale anatomical patterns of cortical connectivity have been collated for several mammalian species (e.g. [1, 2]). Quantitative analysis has revealed that these patterns are neither completely regular nor completely random [3], but exhibit specific attributes, such as short path lengths combined with high clustering [4, 5], short wiring and distinctive hierarchical features [6].

The current article proposes a novel macroscopic approach to studying neuronal dynamics in cortical maps based on concepts from random walks (e.g. [7, 8]) and complex networks (e.g. [9, 10, 11]). Specifically, we examine the interactions between cortex and thalamus in the mammalian central nervous system. The cerebral cortex consists of a network of interconnected functionally specialized regions, each of which maintains reciprocal connections with a specific set of thalamic nuclei. These nuclei relay cortical inputs and outputs, and are thought to play a role in coordinating cortical interactions. The thalamocortical architecture is represented in terms of a complex network, obtained by assigning a node to each cortical region and thalamic nucleus (nodes) and

distributing anatomical connections as links (or edges) between such nodes. The flow of activation of each node is then modeled by an agent engaged in a random walk on the edges of the network. The movement of the agent expresses the exchange or flow of information between adjacent nodes, analogous to a diffusive process. After a long number of time steps  $T$ , the occupancy rate, i.e. the number of times each node was visited divided by  $T$ , captures the rate of participation of each node in network-wide information flow, a dynamic measure analogous to a cortical activation level. In order to relate this dynamical measure to the underlying network topology, we examine the outdegree of each network node.

The anatomical connection matrix of cat cortex and thalamus was obtained from the study by Scannell et al. [2] and contained 53 cortical regions and 42 thalamic regions (Fig. 1a). Most corticocortical connections and all thalamocortical connections are reciprocal (i.e. the matrix is nearly symmetric), and no connections exist between any of the thalamic nodes. Following the analysis in [4] the 53 cortical areas can be divided into four functionally distinct clusters: visual (17, 18, 19, PLLS, PMLS, ALLS, AMLS, VLS, DLS, 21a, 21b, 20a, 20b, 7, AES, PS), auditory (AI, AII, AAF, P, VPC, EPP, Tem), somatomotor (31, 3b, 1, 2, SII, SIV, 4g, 4, 6l, 6m, 5Am, 5Al, 5Bm, 5Bl, SSAi, SSAo), and frontolimbic (PFCM il, PFCM d, PFCL, Ia, Ig, CGa, CGp, RS, 35, 36, pSb, Sb, Enr, Hipp). Visual and auditory clusters are combined into a 'posterior' cluster, while somatomotor and frontolimbic clusters are combined into an 'anterior' cluster. The adjacency matrix is henceforth represented as  $K$ , with the presence of a directed connection from region  $j$  to  $i$  indicated as  $K(i; j) = 1$ .

Because of the relatively small number of involved regions and the fact that any node can be reached from

<sup>E</sup>lectronic address: luciano@ifsc.usp.br

<sup>y</sup>Electronic address: osporns@indiana.edu

any other node in the network (i.e. it forms a single connected component), it is possible to characterize the steady state node occupation effectively in terms of the dominant eigenvector of the respective stochastic matrix (the random walk can be understood as a random driven Markov chain) [12]. In order to do so, a stochastic version  $S$  of the adjacency matrix  $K$  is obtained by making

$$O(j) = \sum_{i=1}^N K(i;j) \quad (1)$$

$$S(i;j) = K(i;j)/O(j) \quad (2)$$

where  $O(j)$  is the outdegree of node  $j$ , i.e. the number of edges emanating from that node. Note that  $S$  is a stochastic matrix because  $S(i;j) \geq 0$  for any  $i$  and  $j$  and  $\sum_{i=1}^N S(i;j) = 1$  for any  $j$ . At steady state, the occupation rate of the random walk movements can be obtained from the eigenvector equation  $S\mathbf{v} = \mathbf{v}$ , where  $\mathbf{v}$  is the eigenvector of the stochastic matrix  $S$  associated to the unit eigenvalue [12]. The eigenvector  $\mathbf{v}$  is normalized so that  $\sum_{i=1}^N v_i = 1$ . The average occupancy rate for each cortical region  $i$  is therefore given by the eigenvector element  $v(i)$ .

Figure 1b shows the outdegree ( $O$ ) versus the occupancy rate ( $R$ ) for the cortical regions obtained by the eigenvalue analysis considering all thalamo-cortical interconnections. Posterior and anterior regions are identified by ' ' and ' ' respectively. This network exhibits a pronounced positive correlation (Pearson correlation  $r = 0.84$ ) between the topological and dynamical measures. We examined the hypothesis that this strong correlation is due to the specific organizational pattern of thalamo-cortical connections. First, we repeat the analysis considering only the cortico-cortical connections. Now, a markedly weaker correlation ( $r = 0.54$ ) was observed between the topological (i.e. outdegree) and dynamical (i.e. occupancy rate) measures, which are illustrated in Figure 1c. When replacing the thalamo-cortical connections by the same number of randomly distributed edges (maintaining reciprocity), we again obtained a weaker correlation ( $r = 0.65$   $0.044$  considering 20 random realizations) between outdegree and occupancy ratio. Randomization of the cortico-cortical connections while keeping the original thalamo-cortical edges results in a decrease, though slightly less marked, of the correlation between outdegree and occupancy rates ( $r = 0.67$   $0.041$  for 20 random realizations). Similar figures were obtained regarding the correlation between indegree and activation rate. Such results seem to indicate that the specific pattern of thalamo-cortical connections promotes a strong correlation between topological and dynamical features of cortical interactions.

In order to try to infer which structural features of the thalamo-cortical interconnections are principally responsible for generating the correlation between topology and dynamics, each thalamic node was isolated together with the respective in and outbound connections { defining

a V-structure, as allowed by the complete lack of inter-connectivity between these nodes. The shortest paths between each of the cortical regions at the destination of the outbound edges of thalamic node  $i$  and all the cortical regions at the origin of the inbound edges are computed. The respective average ( $A_v$ ) and standard deviation values ( $St$ ) are calculated and shown in Figure 1d in decreasing order. The plot shows that the additional edges (and therefore outdegree) imposed by the thalamic V-structures tend to implement cycles of just a few edges among the group of cortical regions to which it is attached. In other words, if the shortest path between two cortical areas attached to a V-structure involves  $k$  edges, the connection to the thalamic structure will define a cycle with  $k + 2$  edges. This effect contributes to a directly proportional increase of the occupancy rate, being potentially related to the observed correlation between the considered topological and dynamical features. Other elements of the thalamo-cortical structure also contributing to the strong correlation include the marked similarity between the in and outdegree at each node as well as the fact that the whole network defines a connected component.

In conclusion, we have shown in this article that a novel combination of concepts from complex networks and random walks/Markov chains can reveal important structural properties underlying thalamo-cortical architecture in the cat brain. Our analysis demonstrates the presence of a strong correlation between network topology (in and outdegree) and dynamical features of cortical activation (occupancy rate) as modeled by random walks. The basic feature of the thalamic connections that is responsible for establishing such a correlation has also been investigated in terms of shortest path analysis between cortical nodes participating in each of the thalamic V-structures. A possible critical feature of such circuits was identified as the relatively short paths between subgroups of cortical regions. In a naive analogy, it is as if the thalamic V-structures provided a feedback mirror to groups of closely connected cortical regions. As for the possible implications of the correlation between outdegree and occupancy rate, it has the two following interesting effects: (1) hubs of connectivity are likely to be hubs of neuronal information flow and activity; and (2) in the case of scale free topology (degrees), the dynamical participation or cortical activation at each node will also become scale free.

The reported findings have potentially important implications for the understanding of cortical functional architecture and pave the way to a number of subsequent studies, including the comparison of cortical connectivity of other species and the use of other kinds of random walks and dynamics. At the same time, several other problems in complex networks can be addressed in terms of the correlation measurement proposed in this article.

# Acknowledgments

Luciano da F. Costa thanks CNPq (308231/03-1) for sponsorship. Olaf Sporns was supported by the James S. McDonnell Foundation.

Adjacency matrix of the directed complex network defined by the cat thalamo-cortical connections (a). The scatterplot of outdegree ( $O$ ) and occupancy rate ( $R$ ) considering all thalamo-cortical connections (b) and only cortical connections (c). The average standard deviation of the shortest paths between the cortical nodes connecting to thalamic nodes, considering only cortical interconnections (d).

## List of Captions [3]:

- 
- [1] D. J. Fellman and D. C. V. Essen, *Cereb. Cortex* 1, 1 (1991).
  - [2] J. W. Scannell, G. A. P. C. Burns, C. C. Hilgetag, M. A. O'Neill, and M. P. Young, *Cereb. Cortex* 9, 277 (1999).
  - [3] O. Sporns, D. Chialvo, M. Kaiser, and C. C. Hilgetag, *Trends Cogn. Sci.* 8, 418 (2000).
  - [4] C. C. Hilgetag, G. A. Burns, M. A. O'Neill, J. W. Scannell, and M. P. Young, *Phil. Trans. R. Soc. Lond. B Biol. Sci.* 355, 91 (2000).
  - [5] O. Sporns, G. Tononi, and G. M. Edelman, *Cereb. Cortex* 10, 127 (2000).
  - [6] L. da F. Costa and O. Sporns, *Eur. Phys. J. B* 48, 567 (2006), q-bio/NC/0508007.
  - [7] F. Spitzer, *Principles of Random Walk* (Springer, 2001).
  - [8] D. ben Avraham and S. Havlin, *Diffusion and reactions in fractals and disordered systems* (Cambridge Univ. Press, 2000).
  - [9] M. E. J. Newman, *SIAM Review* 45, 167 (2003), cond-mat/0303516.
  - [10] R. Albert and A. L. Barabasi, *Rev. Mod. Phys.* 74, 47 (2002).
  - [11] S. Boccaletti, V. Latora, Y. Moreno, M. Chavez, and D.-U. Hwang, *Physics Reports* (2006), accepted.
  - [12] P. Brémaud, *Markov Chains: Gibbs Fields, Monte Carlo Simulation, and Queues* (Springer Verlag, 2001).




FIG .1: Figure 1.