

Adiabatic Theory of Nonlinear Evolution of Quantum States

Jie Liu,¹ Biao Wu,^{1,2} and Qian Niu¹

¹*Department of Physics, The University of Texas, Austin, Texas 78712*

²*Solid State Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831-6032*

(Dated: August 1st, 2002)

We present a general theory for adiabatic evolution of quantum states as governed by the nonlinear Schrödinger equation, and provide examples of applications with a nonlinear tunneling model for Bose-Einstein condensates. Our theory not only spells out conditions for adiabatic evolution of eigenstates, but also characterizes the motion of non-eigenstates which cannot be obtained from the former in the absence of the superposition principle. We find that in the adiabatic evolution of non-eigenstates, the Aharonov-Anandan phases play the role of classical canonical actions.

PACS numbers: 03.65.-w, 03.65.Vf, 03.75.Fi, 71.35.Lk

Adiabatic evolution has been an important method of preparation and control of quantum states[1, 2, 3]. The main guidance comes from the adiabatic theorem of quantum mechanics [4], which dictates that an initial nondegenerate eigenstate remains to be an instantaneous eigenstate when the Hamiltonian changes slowly compared to the level spacings. More precisely, the quantum eigenstate evolves only in its phase, given by the time integral of the eigenenergy (known as the dynamical phase) and a quantity independent of the time duration (known as the geometric phase). The linearity of quantum mechanics then immediately allows a precise statement about the adiabatic evolution of non-eigenstates through the superposition principle.

Our concern here is how the adiabatic theorem gets modified in nonlinear evolution of quantum states. Nonlinearity has been introduced in various forms as possible modifications of quantum mechanics on the fundamental level[5]. Our motivation, however, derives from practical applications in current pursuits of adiabatic control of Bose Einstein condensates (BECs)[6], which can often be accurately described by the nonlinear Schrödinger equation. Here the nonlinearity stems from a mean field treatment of the interactions between atoms. Difficulties in theoretical study of adiabatic control of the condensate arise not only from the lack of unitarity in the evolution of the states but also from the absence of the superposition principle. This problem was recently addressed for BECs in some specific cases[7], and a similar problem was discussed in the past for soliton dynamics[8].

In this Letter, we present a general adiabatic theory for the nonlinear evolution of quantum states (eigenstates or noneigenstates). Our study is conducted by transforming the nonlinear Schrödinger equation into a mathematically equivalent classical Hamiltonian problem, where nonlinearity is no longer a peculiar issue but rather a common character. We can thus apply the adiabatic theory for classical systems[9, 10, 11] to the study of adiabatic evolution of quantum states. The eigenstates become fixed points in the classical problem, whose adiabatic evolution can be understood from a stability anal-

ysis of such points. Aharonov-Anandan phases[12] of cyclic or quasi-cyclic quantum states play the role of canonical action in the classical problem, and are therefore conserved during adiabatic changes of the control parameters. In the linear case, i.e., standard quantum mechanics, these conserved actions are just the occupation probabilities on the eigenenergy levels and the quantum adiabatic theorem becomes a special case of classical adiabatic theorem. Our results are illustrated with a nonlinear two-level model[13], which describes the tunneling of BECs.

General formalism — The quantum system we consider follows the nonlinear Schrödinger equation,

$$i\frac{d|\Psi(t)\rangle}{dt} = H(|\Psi\rangle, \langle\Psi|, \mathbf{R})|\Psi(t)\rangle, \quad (1)$$

where \hbar has been set to one and \mathbf{R} denotes all the system parameters subject to adiabatic change. For clarity and simplicity, we focus in this Letter on quantum systems of finite number of levels, that is, $|\Psi\rangle = \text{col}(\psi_1, \psi_2, \dots, \psi_N)$ and $\langle\Psi| = (\psi_1^*, \psi_2^*, \dots, \psi_N^*)$. We also assume that the system is invariant under the global phase transformation since it is the case of most interest in physics. The total probability $\langle\Psi|\Psi\rangle$ is then conserved and is set to unity hereafter. As shown below, it is possible to separate out the dynamics of the overall phase, and to describe the motion of the remaining variables exactly in terms of an $(N-1)$ dimensional classical Hamiltonian dynamics.

First, we write $|\Psi\rangle = e^{i\lambda(t)}|\Phi\rangle$, where the overall phase λ may be chosen to be the phase of any component, say, ψ_N . Time evolution of the overall phase is given in terms of $|\Phi\rangle$ as derived from Eq.(1)

$$\frac{d\lambda(t)}{dt} = \langle\Phi(t)|i\frac{\partial}{\partial t}|\Phi(t)\rangle - \langle\Phi(t)|H|\Phi(t)\rangle. \quad (2)$$

The first term is related to a geometric phase, called the Aharonov-Anandan phase [12], which will be a key component of our theory.

Second, without the total phase λ , the state $|\Phi\rangle$ becomes a member of the so-called projective Hilbert space

and has $2(N - 1)$ independent real variables, which can be conveniently chosen as

$$\mathbf{Q} = (|\Phi_1|^2, |\Phi_2|^2, \dots, |\Phi_{N-1}|^2), \quad (3)$$

$$\mathbf{P} = (\arg(\Phi_1), \arg(\Phi_2), \dots, \arg(\Phi_{N-1})). \quad (4)$$

They turn out to form a canonical set for an effective classical Hamiltonian dynamics as shown below.

Finally, the dynamics of these variables may be obtained from the time-dependent variational principle of the original nonlinear Schrödinger equation (1) with the Lagrangian

$$\begin{aligned} \mathcal{L} &= \langle \Psi | i \frac{\partial}{\partial t} | \Psi \rangle - \mathcal{H}(|\Psi\rangle, \langle \Psi|, \mathbf{R}) \\ &= \langle \Phi | i \frac{\partial}{\partial t} | \Phi \rangle - \mathcal{H}(|\Phi\rangle, \langle \Phi|, \mathbf{R}) - \frac{d}{dt} \lambda(t), \end{aligned} \quad (5)$$

where \mathcal{H} is the total energy of the system and we have used the gauge symmetry of the Hamiltonian with respect to the overall phase. Note that $\mathcal{H} = \langle \Phi | H | \Phi \rangle$ is true only for a linear system; for a nonlinear system, we generally have $H|\Phi\rangle = \delta \mathcal{H} / \delta (\langle \Phi |)$. Because the overall phase enters in the Lagrangian only through a term of total time derivative, the dynamics of the $2(N - 1)$ independent variables of $|\Phi\rangle$ are closed. With the generalized coordinates introduced in Eqs.(3,4) and the relation $\langle \Phi | i \frac{\partial}{\partial t} | \Phi \rangle = -\mathbf{Q} \cdot \frac{d\mathbf{P}}{dt}$, in virtue of the variational principle of the Lagrangian, we find that motions of the generalized coordinates are governed by a classical Hamiltonian $\mathcal{H}_{cl}(\mathbf{Q}, \mathbf{P}, \mathbf{R}) = \mathcal{H}(|\Phi(\mathbf{Q}, \mathbf{P})\rangle, \langle \Phi(\mathbf{Q}, \mathbf{P})|, \mathbf{R})$, with the equations of motion

$$\frac{d\mathbf{Q}}{dt} = \frac{\partial \mathcal{H}_{cl}}{\partial \mathbf{P}}, \quad \frac{d\mathbf{P}}{dt} = -\frac{\partial \mathcal{H}_{cl}}{\partial \mathbf{Q}}. \quad (6)$$

In this way, we have cast the nonlinear quantum problem (1) into a formalism of classical dynamics, which will allow us to use some important results on the adiabaticity of classical mechanics.

Eigenstates — We first consider adiabatic evolution of quantum eigenstates, which can be defined as extremum states of the system energy by $H(|\Psi^j(\mathbf{R})\rangle, \langle \Psi^j(\mathbf{R})|, \mathbf{R})|\Psi^j(\mathbf{R})\rangle = E^j(\mathbf{R})|\Psi^j(\mathbf{R})\rangle$, where the eigenenergies are Lagrange multipliers for imposing the normalization condition of the states. The eigenstates correspond to fixed points in the classical dynamics (6) at a given \mathbf{R} [14]. For an elliptic fixed point, we expect it to be able to follow adiabatically the control parameter provided the latter changes slowly compared with the fundamental frequencies of periodic orbits around the fixed point. The frequencies can be evaluated by linearizing Eq.(6) about the fixed point [9, 10, 11]. In standard linear quantum mechanics, these frequencies are just the level spacings, so that breakdown of adiabaticity occurs by level crossing. In the nonlinear quantum problem, the fundamental frequencies are in general different from the level spacings, so that adiabaticity can often

be maintained even if the energy levels cross as demonstrated later with a two-level model.

Nonlinearity in our quantum problem not only makes different eigenstates non-orthogonal to each other, but also can produce more eigenstates than the dimension N of the Hilbert space. Some of these additional eigenstates correspond to hyperbolic points in the classical dynamics, characterized by complex fundamental frequencies and strong sensitivity to small perturbations. We thus expect that such quantum eigenstates not to be able to follow adiabatically the control parameter \mathbf{R} . In addition, quantum states corresponding to elliptic fixed points can collide and annihilate with a hyperbolic point during a change of the parameters, leading to the breakdown of adiabaticity.

As an illustration, we consider a nonlinear two-level model,

$$i \frac{\partial}{\partial t} |\Psi\rangle = \left(\frac{R}{2} \sigma_z - \frac{c}{2} \langle \Psi | \sigma_z | \Psi \rangle \sigma_z + \frac{v}{2} \sigma_x \right) |\Psi\rangle, \quad (7)$$

where $|\Psi\rangle = \text{col}(a, b)$ and $\sigma_{z,x}$ is Pauli matrix. This model was proposed to describe the tunneling of Bose-Einstein condensates in optical lattices[13, 15] or in a double-well potential[16]. In the model, the parameter c characterizes the interaction strength between atoms; v is the coupling strength between the two modes. The parameter R can be the Bloch wave number[13] or energy difference between the two wells [16]. We are interested in the tunneling between the energy levels shown in the top panels of Fig.1 when R is increased slowly from the far negative end to the far positive end.

Following the scheme in our general theory, we choose the total phase as $\lambda = \varphi_b = \arg(b)$ and introduce a pair of canonical variables, $q = \varphi_a - \varphi_b$ and $p = |a|^2$. The total energy of the system is $\mathcal{H} = \frac{v}{2} \langle \Psi | \sigma_x | \Psi \rangle + \frac{R}{2} \langle \Psi | \sigma_z | \Psi \rangle - \frac{c}{4} \langle \Psi | \sigma_z | \Psi \rangle \langle \Psi | \sigma_z | \Psi \rangle$. Then we have the equivalent classical Hamiltonian as in Eq.(6),

$$\mathcal{H}_{cl} = v \sqrt{p(1-p)} \cos(q) + \frac{R}{2} (2p-1) - \frac{c}{4} (2p-1)^2. \quad (8)$$

Fig.1 shows in the top panels the structure of eigenenergy levels of Eq.(7) while the phase space orbits of the corresponding classical system (8) is shown in the bottom panels. When $c < v$, there are only two eigenstates and two fixed points. Both fixed points are surrounded by periodic orbits and hence elliptic. The frequencies of the surrounding orbits approach the fundamental frequencies of the fixed points as they get closer to the fixed points. Since both f_1 and f_2 are elliptic, the corresponding quantum states are expected to be able to follow the adiabatically changing R . This has been corroborated by our numerical simulations.

When $c > v$, the situation becomes very different. In the energy band, there are two more eigenstates, forming a loop structure; in the phase space portrait, there appear two more fixed points with one of them, f_3 , being




FIG. 1: The two top panels show eigen-energies as a function of R for two typical cases $c < v$ and $c > v$. The two bottom panels show the corresponding phase space portraits at a given value of R ($=-0.05$). The arrows on the fixed points indicate the directions of their movements as R increases.

hyperbolic. Due to this structure change, the adiabatic evolution becomes very different here. First, the eigenstate corresponding to f_3 will not be able to follow the adiabatic change of R since f_3 is hyperbolic. This has been checked by our numerical integration of Eq.(7). Second, even elliptic point may not be able to follow adiabatically. The fixed point f_1 can annihilate itself by colliding with f_3 as R changes slowly, leading to the breakdown of adiabaticity of the tunneling as reported numerically in Ref.[13]. We also notice that there is a level crossing between f_1 and f_4 at $R = 0$; however, our calculation shows that their fundamental frequencies are $v((\frac{c}{v})^2 - 1)^{1/2}$, not equal to zero. This clearly illustrates our statement in the general theory that the fundamental frequencies are not related to the level spacing in the nonlinear case.

Cyclic and quasi-cyclic states Compared with eigenstates, adiabatic evolution of non-eigenstates is in general very complicated as the motions given by Eq. (6) may be chaotic. We choose to focus on the quantum states around an elliptic point, where the classical orbits are regular. In particular, for the nonlinear two-level problem, the classical dynamics is completely integrable. The non-eigenstates in such cases are cyclic or quasicyclic states, in which the system returns or almost returns to its original state after an evolution. On the $(N - 1)$ -dimensional torus of the regular region, we may introduce a set of action-angle variables, with $\mathbf{I} = (I_1, I_2, \dots, I_{N-1})$, $\boldsymbol{\Theta} = (\Theta_1, \Theta_2, \dots, \Theta_{N-1})$ [9, 10, 11]. The angular variables change with time with the frequencies $\boldsymbol{\omega} = (\omega_1, \omega_2, \dots, \omega_{N-1})$ while the actions \mathbf{I} are constants. More importantly, according to the classical adiabatic theorem [10, 11], the actions \mathbf{I} are adiabatic invariants in the sense that they remain constant even if the control parameter \mathbf{R} changes (slowly) in time. The ex-

istence of these adiabatic invariants presents strong constraint on the motion, and guarantees a quantum state initially close to an eigenstate (elliptic point) to stay near it as the system is changed adiabatically.

Interestingly, we can attach a physical meaning to these adiabatic invariants in the effective classical description by making connection to the Aharonov-Anandan (AA) phase of the quantum states. The AA phase is defined as the time integral of the first term in Eq.(2) for a periodic orbit or a quasi-periodic state[17],

$$\gamma_{AA}(\mathbf{R}) = \int_0^\tau \langle \Phi(t) | i \frac{\partial}{\partial t} | \Phi(t) \rangle dt \quad (9)$$

We can rewrite it with the canonical variables (\mathbf{Q}, \mathbf{P}) and further with the action-angle variables,

$$\gamma_{AA} = \oint \mathbf{P} d\mathbf{Q} = \mathbf{I} \cdot \boldsymbol{\Omega}. \quad (10)$$

where $\boldsymbol{\Omega} = (\omega_1\tau, \omega_2\tau, \dots, \omega_{N-1}\tau)$ and τ is time period. Therefore, the actions are related to the AA phase γ_{AA} , which is an observable physical quantity [18]. In the special case of $N = 2$, there is only one independent action, so the AA phase is simply $\gamma_{AA} = 2\pi I$. This simple connection can be expanded to the general case of $N > 2$, where one can single out a particular cyclic state that involves with only one action I_n . For this cyclic state, we again have the simple relation $\gamma_{AA n} = 2\pi I_n$. As a result, one can identify these AA phases $\gamma_{AA n}$ as the adiabatic invariants in place of the actions I_n .

How do the above adiabatic invariants connect to the familiar notions in the standard linear quantum mechanics? Consider the time evolution of a general state in a linear quantum system for a given \mathbf{R} , $|\Psi\rangle = \sum_n c_n e^{iE^n t} |n\rangle$, where E^n 's are the eigenenergies. This is a (quasi-)cyclic state with the projective wave function given by $|\Phi\rangle = \sum_{n=1}^{N-1} c_n e^{i(E^n - E^N)t} |n\rangle + c_N |N\rangle$. Its AA phase can be computed with Eq.(9); after comparing with Eq.(10), we immediately find that $I_n = |c_n|^2$. Therefore, in the linear quantum mechanics, these adiabatically invariant actions I_n are nothing but the probabilities on the energy levels. In this way, we have rederived the adiabatic theorem of linear quantum mechanics which states that the probability on each energy level is conserved in adiabatic processes. Note that this derivation of the quantum adiabatic theorem through our effective classical description is distinct from the usual semiclassical relation discussed in Ref.[19].

The conservation of probabilities on the energy levels can be generalized to the case where the system is nonlinear only in an intermediate range of the parameter \mathbf{R} (see Fig.2). When the actions (or AA phases) are conserved during the entire process, the initial and final probabilities on the energy levels must remain the same, $I_n = I'_n$, because the system is linear at the beginning and end of the process so the actions are just the occupation probabilities. On the other hand, in the intermediate range




FIG. 2: Nonlinear tunneling of a system that is nonlinear in an intermediate range of the parameter R . I 's and I' 's are the occupation probabilities on different eigenstates, at the beginning and the end of the tunneling process, respectively.

where the nonlinearity dominates, the probabilities will change (even oscillate greatly) since the conserved actions are not probabilities on the energy levels.




FIG. 3: Left panels: Change of probabilities on the two levels with R , which changes with rate $\alpha = 0.0001$, for three different cases. The right two panels show how the AA phases and the fundamental frequencies change with R in these three cases, respectively.

We now illustrate this important result using our two-level model, where, at the two infinite ends with $|R| \gg c$, the nonlinear term can be ignored and the system is effectively linear. For $c < v$, where all the fixed points are elliptic, the fundamental frequency ω for the periodic orbit remains finite and the AA phase (action) is conserved (see lines (a) in the right panels of Fig.3). The initial and final probabilities on each level are indeed the

same (Fig.3(a)), although they oscillate in the intermediate range of the parameter where the system is nonlinear. As the nonlinearity gets strong, the occurrence of tunneling begins to depend on the choice of the initial state. In case (b) of Fig.3, where one starts with probability $I = 0.1$ on level two, tunneling happens; however, in case (c) where one starts with probability $I = 0.8$ on level two, there is no tunneling. The difference is whether there is collision with the hyperbolic point f_3 . In case (b), the initial noneigenstate falls on a periodic orbit surrounding the fixed point f_1 , which will later collide with the hyperbolic point f_3 , where the fundamental frequency drops to zero and the AA phase has a finite jump (see lines (b) in the right panels). The jump height is proportional to the tunneling probability. In case (c), the initial state falls on a periodic orbit around the fixed point f_2 , which will not collide with f_3 .

In conclusion, we have generalized the standard quantum adiabatic theorem to nonlinear quantum systems by transforming them into mathematically equivalent classical Hamiltonians. In the classical systems, the eigenstates become fixed points and their adiabatic evolutions are determined by whether these fixed points are elliptic or hyperbolic. Furthermore, we have found that the adiabatic evolutions of non-eigenstates are controlled by AA phases, which play the role of classical actions. Within the same framework we have also considered the case of closed-loop parameter change as pioneered by Berry[20], we find that Hannay's angles[21] are accumulated at the return of the quantum state (details will be published in the future).

We acknowledge the support of NSF, R.A. Welch Foundation, and the International Center of Quantum Structures in Beijing.

-
- [1] K. Bergmann *et al.*, Rev. Mod. Phys. **70**, 1003 (1998).
 - [2] M.B. Dahan *et al.*, Phys. Rev. Lett. **76**, 4508 (1996); S.R. Wilkinson *et al.*, Phys. Rev. Lett. **76**, 4512 (1996).
 - [3] S.Das, *et al* Phys. Rev. A **65**, 062310 (2002); N.F.Bell, *et al* Phys. Rev. A **65**, 042328 (2002); A.M.Childs, *et al* Phys. Rev. A **65**, 012322 (2002); R.G.Unanyan, *et al* Phys. Rev. Lett. **87**, 137902 (2001).
 - [4] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics* (Pergamon Press, New York, 1977).
 - [5] S. Weinberg, Phys. Rev. Lett. **62**, 485 (1989); S. Weinberg, Ann. Phys. **194**, 336 (1989).
 - [6] F. Dalfovo *et al.*, Rev. Mod. Phys. **71**, 463 (1999); A.J. Leggett, Rev. Mod. Phys. **73**, 307 (2001); R. Dum *et al.*, Phys. Rev. Lett. **80**, 2972 (1998); Z.P. Karkuszewski, K. Sacha, and J. Zakrzewski, Phys. Rev. A **63**, 061601(R) (2001); T. L. Gustavson, *et al*, Phys. Rev. Lett. **88**, 020401 (2002); J. Williams, *et al* Phys. Rev. A **61**, 033612 (2000); Matt Mackie, *et al* Phys. Rev. Lett. **84**, 3803 (2000); Roberto B. Diener, Biao Wu, Mark G. Raizen, and Qian Niu, Phys. Rev. Lett. **89**, 070401 (2002).
 - [7] Y.B. Band, B. Malomed, and M. Trippenbach, Phys.

- Rev. A **65**, 033607 (2002); Y.B. Band and M. Trippenbach, Phys. Rev. A **65**, 053602 (2002); G.J. de Valcárcel, cond-mat/0204406.
- [8] Y.S. Kivshar and B.A. Malomed, Rev. Mod. Phys. **61**, 763 (1989).
- [9] V.I. Arnol'd, *Mathematical Methods of Classical Mechanics* (Springer-Verlag, New York, 1978).
- [10] A.J. Lichtenberg and M.A. Lieberman, *Regular and Stochastic Motion* (Springer-Verlag, 1983); W. Dittrich and M. Reuter, *Classical and Quantum Dynamics: from classical paths to path integrals* (Springer-Verlag, 1992).
- [11] L.D. Landau and E.M. Lifshitz, *Mechanics* (Pergamon Press, New York, 1976).
- [12] For quantum mechanics, see Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593, (1987). For nonlinear systems, see J.C. Garrison and R.Y. Chiao, Phys. Rev. Lett. **60**, 165 (1988); J. Anandan, Phys. Rev. Lett. **60**, 2555 (1988).
- [13] Biao Wu and Qian Niu, Phys. Rev. A **61**, 023402 (2000).
- [14] For an eigenstate $|\Psi^j\rangle$, its time evolution involves only change of phase, $|\Psi^j\rangle(t) = e^{i\phi(t)}|\Psi^j\rangle(0)$. With the erasure of the overall phase λ , the corresponding state $|\Phi^j\rangle$ in the projective space does not change with time, and thus is a fixed point.
- [15] O. Zobay and B.M. Garraway, Phys. Rev. A **61**, 033603 (2000).
- [16] G.J. Milburn *et al.*, Phys. Rev. A **55**, 4318 (1997); A. Smerzi *et al.*, Phys. Rev. Lett. **79**, 4950 (1997); S. Raghuvaran *et al.*, Phys. Rev. A **59**, 620 (1999); I. Marino *et al.*, Phys. Rev. A **60**, 487 (1999); A. Vardi and J.R. Anglin, Phys. Rev. Lett. **86** 568 (2001); I. Zapata, F. Sols and A.J. Leggett, Phys. Rev. A **57**, R28 (1998); S. Kohler and F. Sols Phys. Rev. Lett. **89**, 060403 (2002) .
- [17] For quasiperiodic motions, where the frequencies ω_i are not commensurate with each other, one can use rational numbers n/m to approach the ratios ω_i/ω_j . The accuracy of the approximation can be improved indefinitely with more complex rationals. In this spirit, Eq.(10) also holds for quasi-cyclic motions.
- [18] D. Suter, K.T. Mueller, and A. Pines, Phys. Rev. Lett. **60**, 1218 (1988); H. Weinfurter, Phys. Rev. Lett. **64** 1318 (1990).
- [19] M.V. Berry, J. Phys. A: **17** 1225 (1984).
- [20] M.V. Berry, Proc. R. Soc. London A **392**, 45 (1984).
- [21] J.H. Hannay, J. Phys. A **18**, 221 (1985); M.V. Berry and J.H. Hannay, J. Phys. A. **21**, L325 (1988).