Shedding (red and green) light on "time related hidden parameters"

N. David Mermin

Laboratory of Atomic and Solid State Physics Cornell University, Ithaca, NY 14853-2501

I explain in elementary terms why the critique of Hess and Philipp in Section 3.2 of quant-ph/0103028 fails to invalidate the nontechnical version of Bell's theorem I gave twenty years ago, involving two detectors with 3-pole switches and red and green lights.

In 1985¹ I described a very simple special case of Bell's theorem in which two farapart detectors each have three possible settings (labeled 1, 2, and 3), randomly and independently selected after two appropriately correlated particles have left their common source, and before either particle has arrived at its detector. When a particle does arrive its detector flashes a red (R) or green (G) light. The data accumulated in many runs of this experiment have two important features:²

(i) In those runs in which the detectors happen to have been given the same settings, the light always flash the same color.

(ii) If all runs are examined without reference to the settings of the detectors, the pattern of flashes is completely random; in particular the colors flashed are equally likely to be the same or different.

To account for feature (i) in the absence of any communication between the two wings of the experiment it is plausible to entertain the hypothesis that in each run both particles carry to their detectors identical sets of instructions specifying what color the detector is to flash for each of the three possible settings. The instruction set GGR, for example, means flash G for settings 1 and 2, R for setting 3. An instruction is required for every one of the three possible settings, because the settings are not chosen until after the particles have separated, and each of the pairs of settings 11, 22, and 33 has 1/9 of a chance of being chosen in any given run. The particles must carry such identical instruction sets in

¹ N. David Mermin, Physics Today, April 1985, 38-47.

² Although it is irrelevant to any of the points made below except for footnotes 4 and 13, I note that such data are produced by two spin-1/2 particles in the singlet state, when the detectors are Stern-Gerlach magnets, the three settings are associated with measuring the spin along three coplanar directions 120° apart, and R and G signal spin-up and spin-down at one detector, while signalling spin-down and spin-up at the other.

every single run of the experiment, since each run has a 1/3 chance of ending up one in which the settings are the same.

But this obvious explanation of feature (i) is incompatible with feature (ii) of the data. Since each of the nine possible pairs of settings is equally likely in any run, any of the six instruction sets in which both colors appear (for example GGR) will result in the same color flashing 5/9 of the time (in 11, 22, 33, 12, and 21 runs). The remaining two instruction sets, RRR and GGG, result in the same colors flashing every time. Consequently if such instruction sets were the correct explanation for feature (i), then when the data was examined without regard to the settings, the same colors would be found to flash at least 5/9 of the time, contradicting feature (ii).³ One is left with the puzzle of how to explain feature (i), in the absence of such instruction sets.⁴

I first put forth this special case of Bell's theorem over twenty years ago⁵ to demonstrate to nonscientists in a simple but rigorous way precisely what was so extraordinary about quantum correlations. In the intervening years I have found that its transparency also makes it a good testing ground for claims of conceptual error in the formulation or proof of Bell's theorem. Confusion buried deep in the formalism of very general critiques tends to rise to the surface and reveal itself when such critiques are reduced to the language of my very elementary example.

Such claims of error have recently been made on $arXiv^6$ and $elsewhere^7$ by Karl Hess and Walter Philipp. Conveniently, in Section 3.2 of quant-ph/0103028 they bring their elaborate general argument to bear on my 1985 version of Bell's theorem. In that context their criticism of Bell's theorem is so simple that it is easy to explain why it fails to undermine my argument that no local classical instruction sets can account for the data.

According to Hess and Philipp my argument overlooks the fact that associated with each of the three randomly and independently chosen settings of each detector in each run

³ Bell's inequality in this simple example is "same at least 5/9" of the time". It is violated by the data: "equally likely to be the same or different". Note that this argument is independent of the specific form of the distribution of instruction sets, which can vary from one run to the next (and can even depend on all the data collected in all earlier runs). This is not, however, the kind of time dependence that Hess and Philipp want to pin their local hidden-variables model on.

 $^{^4}$ Quantum mechanics does, of course, provide an explanation, if you want to call it that — namely the one given in footnote 1 above.

⁵ N. David Mermin, American Journal of Physics **49**, 940-943 (1981).

⁶ Karl Hess and Walter Philipp, quant-ph/0103028, quant-ph/0206046.

⁷ Proc. Nat. Academy Sci. (USA) **98**, 14224-27, 14227-34 (2001); Europhys. Lett. **57**, 775-781 (2002).

of the experiment can be an enormous number of different possible microscopic settings, one of which, varying from one run to the next, describes the actual condition of the detector in any given run. Even though the *settings* are randomly and independently chosen in each run, the underlying *microsettings* for a given pair of settings can be correlated between the two detectors because, for example, of time-dependent but common conditions prevailing at the two detectors at the moment⁸ they are triggered.⁹ My simple three-entry instruction sets can therefore be expanded to much larger sets that tell each detector how to flash for each of the many possible microsettings that may be underlying each setting. Because the microsettings that underlie two settings can be correlated even though the settings themselves are independently random, Hess and Philipp assert that my argument "cannot proceed."

They do not, however, spell out why it cannot proceed in the face of this complication. Nor do they produce a local classical hidden-variables model, tailored to my simple special case, that exploits correlations of the microsettings to produce the data I describe.¹⁰ The reason for the latter shortcoming is that my argument can, in fact, proceed, with very little complication. Expanded instruction sets cannot do the job.

What Hess and Philipp fail to note is that if instruction sets are to be based on the microsettings that underly the settings, then the requirement (i), that the lights flash the same colors when the settings are the same, enormously constrains both the possible correlations between the microsettings at the two detectors, and the possible forms of the expanded instruction sets.

To see this note first that if every pair of microsettings associated with the same two settings (11, 22, or 33) has a non-zero probability of occurring in a run, then no matter how strongly correlated the microsettings may otherwise be, the expanded instruction sets for that run must assign the same color to every microsetting, to guarantee that the lights flash the same colors when the settings are the same. Evidently it is enough to specify what that common color is for each of the three entire collections of microsettings — i.e. for each

 $^{^{8}\,}$ All temporal statements may be interpreted in a given inertial frame of reference — for example the laboratory frame.

⁹ From this point onwards the reader must take care always to distinguish between the settings (1, 2, or 3) and the microsettings that underly them.

¹⁰ They do present a very general hidden-variables model intended to reproduce the quantum correlations in the singlet state for the whole continuum of possible settings. Myrvold (quant-ph/0205032) has adapted their very elaborate construction to the special case of only two settings at each detector where he shows that it is explicitly nonlocal. The same conclusion is reached more generally by R. D. Gill, G. Weihs, A. Zeilinger, and M. Zukowski (private communication).

of the three settings. The expanded instruction sets of Hess and Philipp then collapse back to the instruction sets of my example. As before, these must exist in every run, whether or not the detectors do end up with the same setting, because every run has probability 1/3 of being either a 11, 22, or 33 run, and the particles have to be prepared for each of these possibilities when they leave their source. The impossibility of instruction sets then follows exactly as it does when microsettings are not explicitly taken into account.

For more general pair distributions of microsettings, there can be instruction sets which assign more than just one color to all the microsettings that can accompany a given setting, but because of feature (i), this is only possible if the microsettings are strongly correlated in a very particular way. Microsettings associated with the same setting must be constrained so that a microsetting at one detector, instructed to flash red, cannot coexist with a microsetting at the other, instructed to flash green. This requires the microsettings for each setting and at each detector to fall into one of two distinct types, I and II, such that for each setting the underlying microsettings need not be chosen until just before the particles arrive at their detectors and since any run has a 1/3 chance of ending up with the same settings, these divisions of the microsettings into two types must hold in every run and for all three settings.

So as Hess and Philipp emphasize, the instruction sets can indeed be more elaborate than the ones I describe in Ref. 1. But their extension is highly constrained. The most general possible extended instruction set is restricted to specifying for each setting whether type I microsettings flash red and type II green, or vice versa.¹¹ This again takes us back to the instruction sets of my paper, though now we have to reinterpret them.

Now GGR in a given run means that the type-I microsettings under settings 1 and 2 result in a green flash and the type-I microsettings under setting 3 result in a red flash, while the type-II microsettings for each setting flash the color opposite to that specified by the instruction set. If particles with instruction set GGR arrive at their detectors and discover, for example, that the (common) conditions prevailing at the detectors at that moment are the kind that require type-II microsettings for settings 1 and 3 and type-I microsettings for setting 2, then the ouput produced by that instruction set for each of the nine possible pairs of settings will be that specified by the instruction set RGG of my original uncomplicated model. Since each of the nine pairs of settings are equally probable, this effective instruction set will once again result in the same colors flashing 5/9 of the time. Whatever the reinterpreted instruction set turns out to be, it will either result in

 $^{^{11}\,}$ Should all microsettings for a given setting be of the same type, call it type I.

the same colors flashing 5/9 of the time, or (if it is GGG or RRR) all the time. We are back again to the argument of my paper.

The only complication introduced by the microsettings of Hess and Philipp is that the data actually produced by each 3-entry instruction set is only determined when the particles arrive in the neighborhood of their detectors and learn the particular character of the conditions prevailing at the detectors. This has no effect on the validity of the argument that (extended) instruction sets capable of producing feature (i) of the data cannot be compatible with feature (ii).^{12,13}

I conclude that the reason Hess and Philipp have not completed the critique in Section 3.2 of quant-ph/0102038 with a counterexample tailored to that special case is that none exists. My argument against an explanation based on instruction sets is easily expanded to accomodate their "time related hidden parameters".¹⁴

Acknowledgments. I thank Richard Gill, Gregor Weihs, and Marek Zukowski for helpful comments. Supported by the National Science Foundation, Grant No. PHY0098429.

¹² Even this minor complication is not really necessary. Since the conditions prevailing at the detectors (for example the time on two synchronized clocks) must have identical predetermined consequences for whether type-I or type II microsettings underly each of the three possible settings at the two far-apart detectors, it would be a very strange model of the world that did not make the same information (synchronized clock time) directly available at the source as well thereby permitting the instruction sets to assume their final forms before the particles left the source. (The source must know how long each particle takes to reach its detector, since they must arrive at the same time.)

¹³ It is also worth remarking that according to quantum mechanics the statistical character of the data in an EPR experiment is unaffected if the two detections are separated by arbitrarily long time intervals, provided no interactions intervene to disrupt the singletstate spin correlations. To maintain this feature of quantum mechanics in a Hess and Philipp embellishment of my model it would be necessary for the choice of type I or type II microsettings for each setting to be the same for all times. The information about that choice could be available at the source, taking us back to precisely the instruction sets of Ref. 1, without even the modest embellishment described above.

¹⁴ Since their very general and elaborate local hidden-variables model based on timerelated hidden detector parameters includes my elementary example as a special case, it too must contain errors, as has been argued directly in Refs. 10.