

A modern review of the two-level approximation

Marco Frasca
Via Erasmo Gattamelata, 3,
00176 Roma (Italy)

January 16, 2022

Abstract

The paradigm of the two-level atom is revisited and its perturbative analysis is discussed in view of the principle of duality in perturbation theory. The models we consider are a two-level atom and an ensemble of two-level atoms both interacting with a single radiation mode. The aim is to see how the latter can be actually used as an amplifier of quantum fluctuations to the classical level through the thermodynamic limit of a very large ensemble of two-level atoms [M. Frasca, Phys. Lett. A 283, 271 (2001)] and how can remove Schrodinger cat states. The thermodynamic limit can be very effective for producing both classical states and decoherence on a quantum system that evolves without dissipation. Decoherence without dissipation is indeed an effect of a single two-level atom interacting with an ensemble of two-level atoms, a situation that proves to be useful to understand recent experiments on nanoscale devices showing unexpected disappearance of quantum coherence at very low temperatures.

PACS: 42.50.Lc, 42.50.Ct, 42.50.Hz, 03.65.Yz

1 Introduction

It is safe to say that the foundations of quantum optics are built on the concept of a few level atom. Indeed, the most important concept introduced so far in this field is the two-level atom [1]. A lot of physics can be derived by such an approximation and several recent experiments agree fairly well with a description given by the so-called Jaynes-Cummings model describing a two-level atom interacting with a single radiation mode [2]. Besides, by this understanding of radiation-matter interaction it has also been possible to generate Fock states of the radiation field on demand [3].

The radiation-matter interaction currently used is based on some relevant approximations that are still well verified in current experiments: Firstly, it is assumed that the dipole approximation holds, that is, the wavelength of the radiation field is much larger than the atomic dimensions; Secondly, the rotating wave approximation (RWA) is always assumed, meaning by this that just near resonant terms are effective in describing the interaction between radiation and matter, these terms being also described as energy conserving. Indeed, it is sometimes believed that, without these two approximations no two-level atom approximation can really hold [4]. Actually, in the optical regime, that statement can be supported and widely justifies the success of the Jaynes-Cummings model both theoretically and experimentally.

Actually, things are not so straightforward to describe radiation-matter interaction. In fact, Cohen-Tannoudji and coworkers were forced to introduce the concept of dressed states for the two-level atom [5] as, in the regime of microwaves, the RWA fails and a good description of the first experiments in this field were achieved through the concept of dressed states without the RWA [5].

Quantum computation exploited by ionic traps has been firstly put out by Cirac and Zoller [6]. A recent paper by Moya-Cessa et al [7] proved that the standard Jaynes-Cummings model should retain all terms for a Paul trap giving a clear example of dismissal of the rotating wave approximation in quantum optics.

The appearance of laser sources that have large intensity has made thinkable the possibility to extend the study of a two-level atom in such a field. Recent studies seem to indicate that such an approximation can give a viable model for such a physical situation

[8, 9, 10, 11, 12, 13, 14]. In view of this possibility, some methods have been recently devised to approach a solution of the two-level atom in a monochromatic field (being the laser field treated classically) [15, 16, 17]. These studies retain just the two-level and dipole approximations but give up the RWA.

In our recent analysis, it was shown that, treating the laser field classically in this situation, leaves out a relevant part of the behavior of the model [18, 19]. Particularly, if one is especially interested in a resonant behavior, it is seen that some Rabi oscillations are neglected: These oscillations have been recently observed in an experiment with Josephson junctions [20] and originate from the formation of bands for the two levels of the atom due to the radiation field [19].

The aim of this paper is to review, using the approach of duality in perturbation theory [21], the consequences of the validity of the two-level approximation relaxing the RWA approximation. We will see that a single radiation mode interacting with a large number of two-level atoms, without the RWA, provide the amplification of the quantum fluctuations of the ground state of the radiation mode producing a classical radiation field [22] and is able to remove macroscopic quantum superposition states. It is important to point out that these effects arise when the initial state of the ensemble of two-level atoms is properly prepared and the way of generating classical states by unitary evolution in the thermodynamic limit of Ref. [22] is considered. Non dissipative decoherence can also appear as interaction between an ensemble of two-level systems and a quantum system interacting with it [23]. It should be said that another approach to non dissipative decoherence has been recently proposed by Bonifacio and coworkers [24].

The paper is so structured. In section 2 we analyze the model from a general perspective deriving the two-level approximation. In section 3 we present a perturbative analysis of the two-level atom interacting with a single radiation mode, by duality in perturbation theory. In section 4 we give a brief survey of a recent proposal of appearance of classical states and decoherence by unitary evolution in the thermodynamic limit. In section 5 we show how a strong radiation field can be obtained by strong interaction of a single mode with an ensemble of two-level atoms. In section 6 we present a way to approach the measurement problem in quantum mechanics showing how, in the thermodynamic limit, Schrodinger cat states can be removed leaving

only a coherent state describing a classical field. Finally, in section 7 the conclusions are given.

2 A paradigm in quantum optics: Two-level atom

In this section we want to derive the two-level approximation on a general footing. So, let us consider a system described by a Hamiltonian H_0 such that we have a complete set of eigenstates $H_0 |n\rangle = E_n |n\rangle$. We assume, for the sake of simplicity, that the set is discrete. Then, we introduce a time-independent perturbation V . By using the identity $I = \sum_n |n\rangle \langle n|$ we can write the Hamiltonian $H = H_0 + V$ as

$$H = \sum_n (E_n + \hbar \omega) |n\rangle \langle n| + \sum_{m \neq n} \hbar \omega_{nm} |n\rangle \langle m| + \text{h.c.} \quad (1)$$

This Hamiltonian can be rewritten by introducing the operators

$$\begin{aligned} a_{nm} &= |n\rangle \langle m| \\ a_{nm}^\dagger &= |m\rangle \langle n| \\ a_{nm}^3 &= \frac{1}{2} (a_{nm} + a_{nm}^\dagger) \end{aligned} \quad (2)$$

and we can build the algebra of the Pauli matrices, currently named $su(2)$, as it is straightforward to verify that

$$\begin{aligned} [a_{nm}^3, a_{nm}^3] &= 0 \\ [a_{nm}^3, a_{nm}^\dagger] &= i a_{nm}^\dagger \\ [a_{nm}^3, a_{nm}] &= -i a_{nm} \end{aligned} \quad (3)$$

This permits us to prove that our Hamiltonian can be rewritten as the sum of two-level Hamiltonians. In fact, if we change to the interaction picture by the unitary transformation (we use units $\hbar = c = 1$)

$$U_0(t) = \exp \left(-i \sum_n E_n |n\rangle \langle n| t \right) \quad (4)$$

being $E_n = E_n + \hbar \omega$, and we rewrite the V term of the Hamiltonian as

$$V = \sum_{m \neq n} \hbar \omega_{nm} |n\rangle \langle m| + \text{h.c.} = \sum_{m > n} \hbar \omega_{nm} (a_{nm}^\dagger + a_{nm}) \quad (5)$$

we get

$$H_I = U_0^\dagger(t) V U_0(t) = \sum_{m>n} \hbar e^{i(E_m - E_n)t} \rho_{nm} \sigma_{-} + e^{i(E_n - E_m)t} \rho_{nm} \sigma_{+} \quad (6)$$

that proves our assertion: The time evolution of a quantum system can be described by a Hamiltonian being the sum of $su(2)$ Hamiltonians. The reason why this problem is not generally solvable arises from the fact that, given two $su(2)$ parts $H_{I,i}$ and $H_{I,j}$ of this Hamiltonian, it can happen that $[H_{I,i}, H_{I,j}] \neq 0$ and then, the time evolution is not straightforward to obtain analytically. Anyhow, the Hamiltonian H_I can be used to realize some approximate study of a quantum system. The simplest way to get an approximate solution is, indeed, the two-level approximation.

The two-level approximation can be easily justified by assuming that only nearest levels of the unperturbed atom really counts in the time evolution, that is, the more the separation between levels is large and the less important is the contribution to the time evolution of the system. This means that terms with the weakest time dependence in H_I are the most important. Mathematically, this means that we assume a solution by a perturbation series and recognize principally the terms where a slower time dependence is present.

We now consider the case of a single radiation mode interacting with a system having Hamiltonian H_0 . This means in turn that we can choose

$$V = -e a^\dagger a + e \frac{\hbar}{2V} (a^\dagger + a)x \quad (7)$$

being e the electron electric charge, a and a^\dagger the ladder operators of the radiation mode with frequency ω and normalization volume V , and x the coordinate where the field is oriented, having chosen a linear polarization for it. The dipole approximation is taken to hold. In this case we have

$$H_I = -e a^\dagger a + e \frac{\hbar}{2V} \sum_{m>n} e^{i(E_m - E_n)t} \rho_{nm} \sigma_{-} + e^{i(E_n - E_m)t} \rho_{nm} \sigma_{+} (a^\dagger + a) \quad (8)$$

and we are now in a position to obtain the Jaynes-Cummings model. Indeed, we can apply a new unitary transformation to the interaction

picture $U_0^0 = \exp -i\hbar a^\dagger a$ to get

$$H_I^0 = e^{-\frac{i}{2V} \sum_{m>n} \hbar e^{i(E_m - E_n)t} \langle m | \hat{x} | n \rangle \frac{1}{\sqrt{2}} (a^\dagger + a)} + e^{i(E_n - E_m)t} \langle m | \hat{x} | n \rangle \frac{1}{\sqrt{2}} (e^{-i\hbar t} a^\dagger + e^{i\hbar t} a)} : \quad (9)$$

Now, on the basis of the two-level approximation given above, we have to conclude that the only terms to retain are those having no time dependence at all, and these are the resonant terms. Here, we recover the rotating wave approximation (RWA). So, if we have two resonant levels $m = 2 > n = 1$ and we choose the phases of the eigenstates of the unperturbed system so that $\langle m | \hat{x} | n \rangle$ is real, we can finally write the Hamiltonian of the Jaynes-Cummings model as

$$H_{JC} = g(\frac{1}{2}a^\dagger + \frac{1}{2}a) \quad (10)$$

being $g = \hbar \langle 2 | \hat{x} | 1 \rangle e^{-\frac{i}{2V}}$ and the resonance condition $E_2 - E_1 = \hbar$. This gives a proper understanding of the success of the two-level atom approximation in quantum optics when weak fields are involved. It is important to note that also a small detuning can be kept, in agreement with the above discussion.

The Jaynes-Cummings model is good until the other terms in the Hamiltonian are truly negligible. The higher order corrections can be computed by a quite general approach as shown in Ref.[25]. These turn out to be corrections to the Hamiltonian at the resonance (e.g. Bloch-Siegert shift and/or a.c. Stark shift) plus the need to add higher orders of the small perturbation theory to the solution. In the optical regime it is all negligible.

So, as the small perturbation theory plays a crucial role in this analysis, one may ask what one can say if the perturbation V becomes strong. Again, by assuming that only a $su(2)$ component really contributes to the Hamiltonian (9) we need to treat the Hamiltonian

$$H_S^0 = \hbar a^\dagger a + g e^{i(E_1 - E_2)t} \frac{1}{\sqrt{2}} (a^\dagger + a) + e^{i(E_2 - E_1)t} \frac{1}{\sqrt{2}} (a^\dagger + a) : \quad (11)$$

By undoing the interaction picture transformation, this Hamiltonian can be rewritten as

$$H_S = \hbar a^\dagger a + \frac{\hbar}{2} \sigma_z + g \frac{1}{2} (a^\dagger + a); \quad (12)$$

having set $\frac{1}{2} + \frac{1}{2} = 1$, $\frac{1}{2} = \frac{1}{2}$ and $\hbar = E_2 - E_1$. Neither small perturbation theory nor rotating wave approximation apply.

Our aim in the next section is to discuss the perturbative solution of the Schrodinger equation with this Hamiltonian. But, while for the case of the Jaynes-Cummings Hamiltonian we have a fully theoretical justification for our approximation, in the strong coupling regime, the two-level approximation can be satisfactorily justified only by experiment, unless it is exact.

3 Perturbative analysis of an interacting two-level atom

In this section we will give a brief overview of the perturbative solution for a system described by the Hamiltonian (12) in the strong coupling regime. This approach has been described in Ref.[19]. To agree about what a strong coupling regime should be, one has properly to define the weak coupling regime. Indeed, if one has the Hamiltonian

$$H = H_0 + V \quad (13)$$

being an ordering parameter, the weak coupling regime is the one with very small ($\epsilon \rightarrow 0$), while the strong coupling regime is the one with very large ($\epsilon \rightarrow 1$). The duality principle in perturbation theory as devised in Ref.[21] permits to do perturbation theory in both the cases, if one is able to find the eigenstates of V , supposing known those of H_0 . Indeed, small perturbation theory by the usual Dyson series gives (we set $\epsilon = 1$ as this parameter is arbitrary)

$$|j(t)\rangle = U_0(t) T \exp \left(-i \int_0^t V_I(t') dt' \right) |j(0)\rangle \quad (14)$$

being T the time-ordering operator,

$$U_0(t) = \exp(-itH_0) \quad (15)$$

the time evolution of the unperturbed Hamiltonian, and

$$V_I(t) = U_0^\dagger(t) V U_0(t) \quad (16)$$

the transformed perturbation. The choice of a perturbation and an unperturbed part is absolutely arbitrary. So, we can exchange the role of H_0 and V , obtaining the dual Dyson series

$$|j(t)\rangle = U_F(t) T \exp \left(-i \int_0^t H_{0F}(t') dt' \right) |j(0)\rangle \quad (17)$$

being

$$U_F(t) = \exp(-itV) \quad (18)$$

the time evolution of the unperturbed Hamiltonian, and

$$H_{0F}(t) = U_F^\dagger(t) H_0 U_F(t) \quad (19)$$

the transformed perturbation. The duality principle states that, when this exchange is done, restating the series one obtains have the ordering parameters λ and $\frac{1}{\lambda}$ respectively. One is the inverse of the other. So, if we have the eigenstates of V as $|j_n\rangle$ and eigenvalues v_n , one can write

$$U_F(t) = \sum_n e^{iv_n t} |j_n\rangle \langle j_n| \quad (20)$$

If V is time dependent one has formally to rewrite the above as the adiabatic series introducing the geometric phases of the eigenvectors that now could be time dependent themselves [21].

Coming back to the Hamiltonian (12), we realize that small perturbation theory can be recovered if the unperturbed part is that of the two-level atom, otherwise one has a strong coupling perturbation series with an unperturbed Hamiltonian given by

$$V = \lambda a^\dagger a + g_1 (a^\dagger + a) \quad (21)$$

The dressed states originating by diagonalizing this Hamiltonian are well known [5] and are given by

$$|j_n, i\rangle = |j\rangle e^{\frac{g}{\lambda} (a - a^\dagger)} |j_i\rangle \quad (22)$$

being $|1j_i\rangle = |j_i\rangle$ with $j_i = 1$ and, $|j_i\rangle$ the Fock number states that are displaced by the exponential operator [26]. The eigenvalues are $E_n = n! - \frac{g^2}{\lambda}$ and are degenerate with respect to j . So, one has

$$U_F(t) = \sum_{n,j} e^{iE_n t} |j_n, i\rangle \langle j_n, j| \quad (23)$$

and the transformed Hamiltonian becomes

$$H_{0F} = U_F^\dagger(t) H_0 U_F(t) = H_0^0 + H_1 \quad (24)$$

Using the relation [26]

$$\langle j | e^{\frac{g}{\lambda} (a - a^\dagger)} | j_i \rangle = \frac{n!}{j!} \left(\frac{g}{\lambda} \right)^{1-n} e^{-\frac{2g^2}{2!^2}} L_n^{(1-n)} \left(\frac{2g^2}{!^2} \right) \quad (25)$$

with $l = n$ and $L_n^{(l=n)}(x)$ the associated Laguerre polynomial, one gets

$$H_0^0 = \frac{X}{2} e^{-\frac{2g^2}{\Gamma^2}} L_n \frac{4g^2}{\Gamma^2} [j_{n; -1} | i \hbar n; -1 \rangle \langle j | i \hbar n; -1 \rangle + j_{n; -1} | i \hbar n; -1 \rangle \langle j | i \hbar n; -1 \rangle] \quad (26)$$

being L_n the n -th Laguerre polynomial, $j_{n; -1} = e^{\frac{g}{\Gamma} (a - a^\dagger)} j_{n, i}$, and

$$H_1 = \frac{X}{2} e^{i(n-m)\Gamma t} \hbar \frac{2g}{\Gamma} (a - a^\dagger) j_n | i \hbar n; -1 \rangle \langle j | i \hbar n; -1 \rangle \langle j | i \hbar n; -1 \rangle \quad (27)$$

The Hamiltonian H_0^0 can be straightforwardly diagonalized with the eigenstates

$$j_{n; i} = \frac{1}{\sqrt{2}} [j_{n; -1} | i \rangle + j_{n; -1} | j \rangle] \quad (28)$$

and eigenvalues

$$E_{n; i} = \frac{X}{2} e^{-\frac{2g^2}{\Gamma^2}} L_n \frac{4g^2}{\Gamma^2} \quad (29)$$

being $\Gamma = 1$. We see that two bands of levels are formed and two kind of transitions are possible: interband (between levels of the two bands) and intraband (between the levels of a band). This cannot happen if we consider a classical radiation mode, the intraband transitions would be neglected. So, looking for a solution in the form

$$j_F(t) = \sum_n e^{iE_n t} a_n(t) j_{n; i} \quad (30)$$

one gets the equations for the amplitudes [19]

$$i \dot{a}_{m; 0}(t) = \frac{X}{2} \sum_{n \in m; } a_n(t) e^{i(E_n - E_{m; 0})t} e^{i(m-n)\Gamma t} \hbar \frac{2g}{\Gamma} (a - a^\dagger) j_{n; -1} \frac{0}{2} + \hbar \frac{2g}{\Gamma} (a - a^\dagger) j_{n; -1} \frac{1}{2} : \quad (31)$$

This equations can also display Rabi oscillations between the eigenstates $j_{n; i}$ that can be seen as microscopic quantum superposition states, both for interband and intraband transitions [19]. States like these could prove useful for quantum computation. These kind of Rabi oscillations in Josephson junctions have been recently observed [20].

At this stage it is very easy to do perturbation theory in the strong coupling regime for this model. One has to rewrite the initial condition $j_i(0)$ by the eigenstates $j_n; i$ obtaining in this way the amplitudes $a_n; i(0)$. Then, one has to solve eq.(31) perturbatively as done routinely in the weak coupling regime. In case of a resonance one has to apply the RWA obtaining Rabi oscillations. In this way we see that the dual Dyson series, as it should be expected, displays all the features of the standard weak coupling expansion.

4 Classical states and decoherence by unitary evolution

An ensemble of independent two-level systems can behave classically. This has been proven in Ref.[22]. Indeed, let us consider a Hamiltonian

$$H_c = \frac{1}{2} \sum_{i=1}^N \sigma_{3i} \quad (32)$$

Assuming distinguishable systems, we can take for the initial state the one given by

$$j_i(0) = \sum_{i=1}^N (\sigma_{1i} \# i + \sigma_{2i} \# i) \quad (33)$$

with $j_{1i}^2 + j_{2i}^2 = 1$, $\sigma_{1i} \# i = j_{1i}$ and $\sigma_{2i} \# i = j_{2i}$. The time evolution gives us

$$j_i(t) = \sum_{i=1}^N (\sigma_{1i} e^{i\frac{t}{2}} \# i + \sigma_{2i} e^{-i\frac{t}{2}} \# i) \quad (34)$$

For the Hamiltonian it is easy to verify that

$$H_c j_i = H_c j_i(0) = \frac{1}{2} \sum_{i=1}^N (j_{1i}^2 - j_{2i}^2) = \frac{1}{2} k_H N \quad (35)$$

being k_H a fixed number between -1 and 1. So, in a similar way, it is easy to obtain the fluctuation

$$(H_c)^2 = H_c j_i(0) H_c j_i(0) = H_c j_i(0) H_c j_i(0) = k_H^0 N \quad (36)$$

being $k_H^0 = \prod_{i=1}^N j_i^2 (1 - j_i^2) = N$ and one sees that k_H^0 is a finite number independent on N . So, as it happens in statistical thermodynamics, in the thermodynamic limit $N \rightarrow \infty$ we see that quantum fluctuations are not essential, that is

$$\frac{H_c}{H_c} / \frac{1}{N} : \quad (37)$$

The "laws of thermodynamics" are obtained by the Ehrenfest's theorem and are the classical equations of motion. That is, the variables $x = \prod_{i=1}^N x_i$, $y = \prod_{i=1}^N y_i$ and $z = \prod_{i=1}^N z_i$ follow, without any significant deviation, the classical equations of motion, when the thermodynamic limit is considered and the time evolution is computed averaging with the above $j(t)$. So, we have found a classical object out of the quantum unitary evolution. The main point here is that classical objects can be obtained by unitary evolution in the thermodynamic limit depending on their initial states. Actually, one cannot apply the above argument if e.g. the state of the system is an eigenstate of the Hamiltonian H_c . Besides, a classical state obtained by unitary evolution, per se, does not produce decoherence. Rather, it is interesting to see what happens when such a classical state interacts with some quantum system. This is a relevant problem that can prove quantum mechanics and its fluctuations to be just the bootstrap of a classical world: If by unitary evolution, in the thermodynamic limit, some classical objects are obtained and these are permitted to interact with other quantum objects, the latter can decohere or become classical by themselves.

As a relevant example, let us consider the interaction of the above system with a two-level atom. This model has been considered in Ref.[23] as a possible explanation of recent findings in some experiments with nanoscale devices that show unexpected decoherence in the low temperature limit [27, 28]. The Hamiltonian can be written as

$$H_D = \frac{\epsilon_0}{2} z + \frac{1}{2} \sum_{i=1}^N (x_i x_i + z_i z_i) - J \sum_{i=1}^N x_i \quad (38)$$

where J is the coupling. The Hamiltonian of the two-level system (second term in eq.(38)) is taken not diagonalized, but this does not change our argument as the above analysis still applies. Finally, ϵ_0 is the parameter of the Hamiltonian of the two-level atom that we want to study. We need another hypothesis to go on, that is, we

assume that the coupling J is larger than any parameter of the two-level systems ω_i, γ_i , but not with respect to ω_0 . By applying duality in perturbation theory, we have the leading order solution

$$|j(t)\rangle = \exp\left[-\frac{it}{2}\omega_0\right] \sum_{i=1}^N \exp(iJt\omega_i) |j(0)\rangle_i \quad (39)$$

Now, as already seen, we have to choose the state of the two-level system s as given by the product of the lower eigenstates of each ω_i . This can be seen as a kind of "ferromagnetic" state and is in agreement with our preceding discussion. So, we take

$$|j(0)\rangle = \prod_{i=1}^N |j\rangle_i \quad (40)$$

being $\omega_j = \omega_i$ and, similarly, $\gamma_j = \gamma_i$. The state of the ensemble of two-level systems agrees fairly well with the one of eq.(33). So, one has, by tracing away the state of the ensemble of two-level systems being not essential for our aims,

$$|j(t)\rangle = \exp\left[-\frac{it}{2}\omega_0\right] \sum_{i=1}^N \exp(iJt\omega_i) |j\rangle_i \quad (41)$$

that defines a spin coherent state [29, 30]. The point we are interested in is the thermodynamic limit. When N is taken to be large enough, the contribution ω_0 can be neglected and we have a reduced density matrix

$$\rho(t) = \exp(iJt\omega_0) \sum_{i,j} \rho_{ij} \exp(-iJt\omega_i) \quad (42)$$

being

$$\rho_{ii}(t) = \frac{1 + \cos(2N Jt)}{2} \quad (43)$$

$$\rho_{i\#}(t) = i \frac{1}{2} \sin 2N Jt \quad (44)$$

$$\rho_{\#i}(t) = i \frac{1}{2} \sin 2N Jt \quad (45)$$

$$\rho_{\#\#}(t) = \frac{1 - \cos(2N Jt)}{2}; \quad (46)$$

where we have oscillating terms with a frequency NJ that goes to infinity in the thermodynamic limit. The only meaning one can attach to

such a frequency is by an average in time (see [23] and Refs. therein) and decoherence is recovered. So, when the ensemble of two-level systems strongly interacts with a quantum system produces decoherence and quantum behavior disappears, in the thermodynamic limit. The ensemble of two-level systems should evolve unitarily, producing a classical behavior. Higher order corrections have also been studied in Ref.[23]. It is important to stress that this behavior should be expected at zero temperature as quantum coherence is lost otherwise.

Such a behavior, having a characteristic decoherence time scale depending on the number of two-level systems that interact with the quantum one, has been recently observed in quantum dots [28]. In this case, the ensemble of two-level systems can be given by the spins of the electrons that are contained in the two dimensional electron gas in the dot. Another source of decoherence in quantum dots could be given by the spins of the nuclei interacting through an hyperfine interaction with the spin of the conduction electrons [31]. The nuclei are contained in the heterostructures forming the dot. In this case we have a similar spin-spin interaction but isotropic. The mechanism that produces the decay of the off-diagonal parts of the density matrix, also in this case, appears to be the same, being the decoherence produced dynamically and dependent on the initial state.

5 Application of quantum fluctuations to the classical level

Spontaneous emission can be seen as a very simple example of decoherence in the "thermodynamic limit" of the number of radiation modes. Indeed, we can consider a two-level system interacting with N radiation modes and being resonant with one of it. In the limit of a small coupling between radiation and two-level system and very few spectator modes, one has Rabi oscillations, a clear example of quantum coherence. When the number of spectator modes is taken to go to infinity, a description with continuum is possible and this gives rise to decay, i.e. spontaneous emission. This representation of the process of decay is very well described in Ref.[5].

Here, we want to consider the opposite situation, that is, a single radiation mode strongly interacting with an ensemble of N two-level systems. We are going to show that, when the ensemble of two-level

system s behaves as a classical object if left alone, the radiation field, supposed initially in the ground state, will have the zero point fluctuations amplified to produce a classical field having intensity dependent on N , the number of two-level systems.

As done in Ref.[22], we modify the model of eq.(12) to consider N two-level systems interacting with a single radiation mode, as

$$H_S = \sum_{i=1}^N \frac{1}{2} \sigma_{3i} + g \sum_{i=1}^N \sigma_{1i} (a^\dagger + a) \quad (47)$$

Then, the strong coupling regime amounts to consider the Hamiltonian $\sum_{i=1}^N \frac{1}{2} \sigma_{3i}$ as a perturbation, as already done in sec.3 for a single two-level atom. We take as initial state of the full system $|j\rangle_{i=1}^N = \prod_{i=1}^N |j\rangle_{1i}$, so that, the ensemble of two-level systems is again in a kind of "ferromagnetic" state representing its ground state. Besides, no photon is initially present. It is a well known matter that the fluctuations of the radiation mode are not zero in this case. The unitary evolution at the leading order gives us

$$|j\rangle(t) = \exp\left[-it \sum_{i=1}^N \frac{1}{2} \sigma_{3i}\right] \exp\left[itg \sum_{i=1}^N \sigma_{1i} (a^\dagger + a)\right] \prod_{i=1}^N |j\rangle_{1i} \quad (48)$$

that, by use of a known disentangling formula [30], produces

$$|j\rangle(t) = e^{i(t)} e^{-it \sum_{i=1}^N \frac{1}{2} \sigma_{3i}} D[\alpha(t)] |j\rangle(0); \quad (49)$$

being

$$\alpha(t) = \frac{N g^2}{i^2} (1 - \sin(i t)); \quad (50)$$

$$\alpha(t) = \frac{N g}{i} (1 - e^{i t}); \quad (51)$$

and

$$D[\alpha(t)] = \exp[\alpha(t) a^\dagger - \alpha(t) a]; \quad (52)$$

We conclude that, at the leading order, the radiation mode evolves as a coherent state with a parameter given by

$$\alpha(t) = \frac{N g}{i} (e^{i t} - 1) = -i(t) e^{i t}; \quad (53)$$

In this way, we have amplified the quantum fluctuations of the field, being the fluctuation of the number of photons proportional to N , but,

as the average of the number of photons is proportional to N^2 , this ratio goes to zero as the thermodynamic limit $N \rightarrow \infty$ is taken. As it is well known [32], this produces a classical field with increasing intensity as the number of two-level systems increases, proving our initial assertion. We can see that the amplification of quantum fluctuations gives rise to a classical object, as initially no radiation field is present.

Higher order corrections have been studied in Ref.[33], showing that are not essential in the thermodynamic limit. So, this effect will prove to be a genuine example of production of a classical object by unitary evolution in the thermodynamic limit with possible technological applications.

6 Two-level Systems, Thermodynamic Limit and Schrodinger Cat States

Decoherence, as currently devised, is able to remove superposition states through interaction of the environment with a quantum system. This does not solve the measurement problem in quantum mechanics as, mixed forms of the density matrix do not give single states required by the measurement process [34]. This problem is fairly well described by the Schrodinger cat paradox as we ask that the cat has a well defined state at the observation. Schrodinger cat states have been currently produced in laboratory in a form of superposition of coherent states (see e.g. the second reference in [1])

$$|j_{cat}\rangle = N (|j e^{i\theta_1}\rangle + |j e^{i\theta_2}\rangle) \quad (54)$$

being N a normalization factor, and θ_1, θ_2 real numbers. To understand the measurement problem, we would like to get a single state out of such a superposition after unitary evolution, if possible. In this way, we can show, at least in this case, that quantum mechanics is, indeed, a self-contained theory. This possibility can be exploited by an ensemble of two-level systems interacting with a single radiation mode in the thermodynamic limit.

In order to accomplish our aim, we consider again the Hamiltonian (47) with the initial condition (54) multiplied by the "ferromagnetic state", $|j\rangle = \bigotimes_{i=1}^N |1\rangle_i$, for the ensemble of atoms to have $|j(0)\rangle = |j_{cat}\rangle |j\rangle$. The unitary evolution, assuming the Hamiltonian of two-

level atoms as a perturbation, gives in this case [35]

$$j(t) = e^{i\phi(t)} N (e^{i\phi_1(t)} j(t) e^{i\phi_2(t)} + e^{i\phi_1(t)} i + e^{i\phi_2(t)} j(t) e^{i\phi_1(t)} + e^{i\phi_1(t)} i) j(t) \quad (55)$$

where use has been made of the property of the displacement operator for coherent states so to yield

$$\phi(t) = \frac{N^2 g^2}{2} (t \sin(\phi(t))); \quad (56)$$

$$\phi(t) = \frac{N g}{2} (1 - e^{i\phi(t)}) \quad (57)$$

and

$$\phi_1(t) = \frac{i}{2} [\phi(t) e^{i\phi(t)} - \phi(t) e^{-i\phi(t)}]; \quad (58)$$

$$\phi_2(t) = \frac{i}{2} [\phi(t) e^{i\phi(t)} - \phi(t) e^{-i\phi(t)}]; \quad (59)$$

being the phases $\phi_1(t)$ and $\phi_2(t)$ generated by multiplication of two displacement operators. In the thermodynamic limit $N \rightarrow \infty$ one gets the macroscopic state $j(t) e^{i\phi(t)} i$ and the cat state seems gone away.

Actually, we have a couple of problems before one can claim that, effectively, the cat state has been removed. Firstly, all we have done is a displacement to infinity and no decoherence seems to be implied in such an operation, so all the properties of a superposition state have to be there anyway. Secondly, we have done perturbation theory and one has to prove that, in the thermodynamic limit, higher order corrections are negligible.

The first question is answered immediately by computing the interference term in the Wigner function of the state (55). In the thermodynamic limit such a term should become negligible. One has

$$\begin{aligned} W_{INT} = & \frac{2}{\pi} \exp \left[-\frac{1}{2} \left(x + \frac{p}{2N} \frac{g}{2} (1 - \cos(\phi(t))) \right)^2 - \frac{1}{2} \left(p + \frac{2N g}{2} \sin(\phi(t)) \right)^2 \right] \\ & \exp \left[-\frac{1}{2} \left(p + \frac{2N g}{2} \sin(\phi(t)) \right)^2 + \frac{1}{2} \left(p - \frac{2N g}{2} \sin(\phi(t)) \right)^2 \right] \\ & \cos \left(\frac{p}{2} \sin(\phi(t)) - x \cos(\phi(t)) + \frac{1}{2} \sin(2\phi(t)) + 8 \frac{N g}{2} \sin(\phi(t)) (1 - \cos(\phi(t))) \right) \end{aligned} \quad (60)$$

This term has a quite interesting form as displays a term that rapidly oscillates in time for N becoming increasingly large. If such oscillations become too rapid, we can invoke blurring in time to have these

terms averaged away. Otherwise, as the Wigner function can be measured, we will get a way to probe, immediately and by very simple means, Planck time physics. So, we can safely claim that we have true decoherence in the thermodynamic limit and the cat state is effectively removed generating a macroscopic classical state. It should be said that ordinary decoherence is generally invoked for blurring in space [36] and there is no reason to say that also blurring in time cannot occur. We can recognize here the same argument used in order to obtain decoherence for the model of sec.4 to explain decoherence in quantum dots. A sound mathematical basis for such an approach is given in [37].

The second question can be straightforwardly answered by computing higher order corrections through the strong coupling expansion discussed in sec.3. The proof is successfully accomplished in Ref.[35] and we do not repeat it here.

It is interesting to note that all the new points introduced so far for analyzing two-level systems can conspire to generate a new view of the way measurement is realized in quantum mechanics, possibly making the theory self-contained.

7 Discussion and Conclusions

We have presented a brief review about some new views on two-level systems. These appear to be even more important today with a lot of new effects to be described and experimentally observed. Paramount importance is acquiring the decoherence due to an ensemble of two-level systems as it is becoming ubiquitous to different fields of application as quantum computation and nanotechnology, fields that maybe could merge. To face these new ways to see the two-level approximation, we have exposed new mathematical approaches to analyze models in the strong coupling regime. This regime has been pioneered by Bender and coworkers [38] in quantum field theory in the eighties, but, with our proposal of duality in perturbation theory, a possible spreading of such ideas to other fields is now becoming possible. Indeed, a lot of useful results, as those presented here, are obtained by this new approach and, hopefully, the future should deserve some other interesting results.

The idea of a non dissipative decoherence is also relevant due to

the recent findings in the field of nanodevices, where unexpected loss of quantum coherence has appeared in experiments performed at very low temperatures. In these cases, it appears as the standard idea of decoherence, meant as interaction of a quantum system with an external environment, seems at odds with some experimental results, even if an interesting proposal through the use of quantum fluctuations has been put forward by Buttiker and coworkers [39].

The conclusion to be drawn is that, today, a lot of exciting work at the foundations of quantum mechanics is expecting us, giving insight toward new understandings and methods and, not less important, applications.

References

- [1] L. Allen, and J. H. Eberly, *Optical Resonance and Two-level Atoms*, (Wiley, New York, 1975). For a more recent exposition see W. P. Schleich, *Quantum Optics in Phase Space*, (Wiley, Berlin, 2001).
- [2] J. M. Raimond, M. Brune, and S. Haroche, *Rev. Mod. Phys.* **73**, 565 (2001) and Refs. therein.
- [3] S. Brattke, B. T. H. Varcoe, and H. Walther, *Phys. Rev. Lett.* **86**, 3534 (2001).
- [4] M. O. Scully, and M. S. Zubairy, *Quantum Optics*, (Cambridge University Press, Cambridge, 1997).
- [5] C. Cohen-Tannoudji, J. Dupont-Roc and C. Grynberg, *Atom - Photon Interactions*, (Wiley, New York, 1992).
- [6] J. I. Cirac and P. Zoller, *Phys. Rev. Lett.* **74**, 4091 (1995).
- [7] H. Moya-Cessa, A. Vidiella-Barranco, J. A. Roversi, D. S. Freitas, and S. M. Dutra, *Phys. Rev. A* **59**, 2518 (1999).
- [8] B. Sundaram, and P. W. Milonni, *Phys. Rev. A* **41**, R6571 (1990).
- [9] M. Yu. Ivanov, and P. B. Corkum, *Phys. Rev. A* **48**, 580 (1993).
- [10] Y. Dakhnovskii, and H. Metiu, *Phys. Rev. A* **48**, 2342 (1993); Y. Dakhnovskii, and R. Bavli, *Phys. Rev. B* **48**, 11020 (1993).
- [11] A. E. Kaplan, and P. L. Shkolnikov, *Phys. Rev. A* **49**, 1275 (1994).

- [12] F. I. Gauthey, C. H. Keitel, P. L. Knight, and A. Maquet, Phys. Rev. A 52, 525 (1995); 55, 615 (1997); F. I. Gauthey, B. M. Garraway, and P. L. Knight, Phys. Rev. A 56, 3093 (1997).
- [13] M. Pons, R. Taieb, and A. Maquet, Phys. Rev. A 54, 3634 (1996).
- [14] P. P. Corso, L. Lo Cascio, and F. Persico, Phys. Rev. A 58, 1549 (1998) and Refs. therein. See also A. Di Piazza and E. F. Iordilino, Phys. Rev. A 64, 013802 (2001); A. Di Piazza, E. F. Iordilino, and M. H. Mittleman, Phys. Rev. A 64, 013414 (2001)
- [15] J. C. A. Barata, and W. F. Wreszinski, Phys. Rev. Lett. 84, (2000); J. C. A. Barata, Ann. Henri Poincaré 2, 963 (2001).
- [16] V. Delgado, and J. M. Gomez Llorente, J. Phys. B 33, 5403 (2000); A. Santana, J. M. Gomez Llorente, and V. Delgado, J. Phys. B 34, 2371 (2001).
- [17] M. Frasca, Phys. Rev. A 56, 1548 (1997); 60, 573 (1999).
- [18] M. Frasca, J. Opt. B: Quant. Semicl. Opt. 3, S15 (2001). This behavior agrees with the result given in E. F. Iordilino, Phys. Rev. A 59, 876 (1999).
- [19] M. Frasca, Z. Naturforsch. 56 a, 197 (2001); Phys. Rev. A 66, 023810 (2002). For a generalization of this approach see K. Fujii, quant-ph/0203135, to appear on J. Phys. A: Math. Gen., and quant-ph/0210166.
- [20] Y. Nakamura, Yu. A. Pashkin, and J. S. Tsai, Phys. Rev. Lett. 87, 246601 (2001).
- [21] M. Frasca, Phys. Rev. A 58, 3439 (1998).
- [22] M. Frasca, Phys. Lett. A 283, 271 (2001); Erratum: ib. 306, 184 (2003). Some general theorems about this matter are given in M. Frasca, Phys. Lett. A 308, 135 (2003).
- [23] M. Frasca, Physica E 15, 252 (2002).
- [24] R. Bonifacio, N. Chin. 114B, 473 (1999); R. Bonifacio, S. Olivares, P. Tombesi and D. Vitali, Phys. Rev. A 61, 053802 (2000).
- [25] M. Frasca, Phys. Rev. A 58, 771 (1998).
- [26] F. A. M. de Oliveira, M. S. Kim, P. L. Knight, and V. Buzek, Phys. Rev. A 41, 2645 (1990). See also K. Fujii, quant-ph/0202081.

- [27] A .G .Huibers, J.A .Folk, S.R .Patel, C .M .M arcus, C .I.D unuez and J .S .Harris, Jr., Phys. Rev. Lett. 83, 5090 (1999); P .M o-hanty, E .M .Q .Jariwala and R .A .W ebb, Phys. Rev. Lett. 78, 3366 (1997); P .M ohanty and R .A .W ebb, Phys. Rev. B 55, 13452 (1997).
- [28] C .P rasad, D .K .Ferry, A .Shailos, M .E lhasan, J .P .B ird, L .H .L in, N .A oki, Y .O chiai, K .Ishibashi and Y .A oyagi, Phys. Rev. B 62, 15356 (2000); D .P .P ivin, Jr., A .A ndresen, J .P .B ird and D .K .Ferry, Phys. Rev. Lett. 82, 4687 (1999). D .K .Ferry, private com m unication.
- [29] F .T .A recchi, E .C ourtens, R .G ilm ore and H .Thom as, Phys. Rev. A 6, 2211 (1972).
- [30] W ei-M in Zhang, Da Hsuan Feng and R .G ilm ore, Rev. M od. Phys. 62, 867 (1990).
- [31] A .V .K haetskii, D .Loss, and L .G lazman, Phys. Rev. Lett. 88, 186802 (2002); J .Schliemann, A .V .K haetskii, and D .Loss, Phys. Rev. B 66, 245303 (2002). D .Loss, private com m unication.
- [32] L .M andel and E .W olf, Optical C oherence and Q uantum Optics, (C am bridge University Press, C am bridge, 1995).
- [33] M .Frasca, J .Opt. B :Q uant. Sem iclass. Opt. 4, S443 (2002). This is a contribution to Garda 2001 Conference Proceedings edited by R .B onifacio and D .V itali.
- [34] S .L .A dler, quant-ph/0112095. F .Labe, Am .J .Phys. 69, 655 (2001).
- [35] M .Frasca, quant-ph/0212119.
- [36] M .V .Berry, "Chaos and the sem iclassical lim it of Q uantum m echanics (is the m oon there when som ebody looks?)" , in Q uantum M echanics: Scienti c perspectives on divine action (eds: Robert John Russell, Philip C layton, Kirk W egter-M cNelly and John Polkinghome), Vatican Observatory CTNS publications, pp 41-54 (2001).
- [37] G .H .H ardy, D ivergent Series, (A M S C helsea Publishing, P rovi-dence, 1991), pp. 10 .
- [38] C .M .Bender, F .C ooper, G .S .G uralnik and D .H .Sharp, Phys. Rev. D 19, 1865 (1979); C .M .Bender, F .C ooper, G .S .G uralnik,

- H. Moreno, R. Roskies and D. H. Sharp, Phys. Rev. Lett. 45, 501 (1980); C. M. Bender, F. Cooper, R. Kenway and L. M. Simmons, Jr., Phys. Rev. D 24, 2693 (1981).
- [39] P. Cedraschi and M. Buttiker, Phys. Rev. B 63, 165312 (2001); Ann. Phys. 289, 1 (2001); M. Buttiker, in "Complexity from Microscopic to Macroscopic Scales: Coherence and Large Deviations", edited by Arne T. Skjeltorp and Tamás Vicsek (Kluwer, Dordrecht 2002). p. 21 – 47.