## Cold atom s in a high-Q ring-cavity

D.Knuse, M.Ruder, J.Benhelm, C.von Cube, C.Zimmermann, and Ph.W.Courteille

Physikalisches Institut, Eberhard-Karls-Universitat Tubingen,

Auf der Morgenstelle 14, D-72076 Tubingen, Germany

Th. E lsasser, B. Nagorny, and A. Hemmerich Institut fur Laser-Physik, Universitat Hamburg, Jungiusstr. 9, D-20355 Hamburg, Germany (Dated: March 29, 2024)

W e report the con nem ent of large clouds of ultra-cold  $^{85}$ R b atom s in a standing-wave dipole trap form ed by the two counter-propagating m odes of a high-Q ring-cavity. Studying the properties of this trap we dem onstrate loading of higher-order transverse cavity m odes and excite recoil-induced resonances.

PACS num bers: 32.80 P j, 32.70 Jz, 42.50 V k, 42.50 -p

The trem endous progress of ultra-cold atom ic physics in the past decades is mainly due to the invention of powerful techniques for trapping and cooling atom s. Besides m agnetic traps, optical forces are used to store atom s in tightly focused red-detuned laser beam s, in the antinodes of a standing wave formed by two counter-propagating laser beam s or even in three-dim ensional optical lattices. Spontaneous scattering processes are avoided by tuning the lasers far from resonance. High light intensities are then needed to keep the atom - eld coupling strong. Large intensity enhancements are achieved with optical cavities [1]. The coupled system made up of the cavity mode and the atom s exhibits fascinating novel features [2, 3, 4]. For example, cooling procedures have been proposed [5, 6, 7, 8], where the kinetic energy of the atom s is dissipated via the decay of the cavity eld to which the atom s are coherently coupled, rather than via spontaneous decay of atom ic excitation energy. This bears the advantage that the cooling is only limited by the technical parameter of the cavity decay rate. As the cooling procedure is quite insensitive to the details of the atom ic level structure, it should work even form olecules.

In this paper, we study atom ic trapping in a high-Q ring-cavity with both propagation directions pumped. This kind of cavity is particular in the following sense. The optical dipole force exerted by a standing wave on atom s can be understood in term s of a momentum transfer by coherent redistribution (via Rayleigh scattering) of photons between the counter-propagating laser beam s. In multimode con gurations, e.g. a ring-cavity, the photon redistribution can occur between di erent modes [9]. This im plies that the atom s have a noticeable backaction on the optical elds and that the atom ic dynamics can be monitored as an intensity [10, 11] or a phase im balance between the modes. In a ring-cavity, the scattering of a photon between the counter-propagating modes slightly shifts the phase of the standing wave. This shift, being strongly enhanced by a long cavity life time, is sensed by all atoms trapped in the standing wave. Consequently, the simultaneous interaction of the atoms with the two eld modes couples the motion of all atoms [12, 13], so that in contrast to conventional standing wave dipole traps, the atoms do not move independently of each other.

In a recent article [14] we have shown that a fardetuned optical lattice can be form ed inside a high-Q ring-cavity and that heating due to intensity uctuations can be kept at very low levels despite of the need to maintain a sharp resonance condition by a high-bandwidth servo control. Here we present a setup where we show that the atom ic motion can be probed non-destructively and in-situ by recoil-induced resonances [15]. There is an important reason to look for methods to study the atom icm otion which do not rely on sudden changes of the intra-cavity intensity. This is because the long lifetime of a high-Q cavity inhibits the non-adiabatic switch-o of deep cavity dipole traps and thus impedes a straight forward interpretation of tim e-of- ight (TOF) im ages to determ ine the axial tem perature.

We llour ring-cavity dipole trap with  $^{85}$ Rb atoms from a standard magneto-optical trap (MOT) which is baded from a vapor generated by a nubidium dispenser. Typically, we bad  $10^8$  atoms into the MOT at temperatures around 140 K with a vapor pressure of 3  $10^9$  m bar.

The geometry of the ring cavity is shown in Fig. 1. The transm issions  $T_i$  of the m innors depend on the (linear) polarization of the light modes. For s-polarization  $T_0 = 27 \quad 10^6$ ,  $T_1 = T_2 = 2 \quad 10^6$ , while for p-polarization  $T_0 = 2200 \quad 10^6$ ,  $T_1 = T_2 = 9 \quad 10^6$ . The experiments presented here are carried out with p-polarized light and a measured nesse of F = 2500, which corresponds to an intracavity intensity decay rate of 2 1.4 M H z. How ever, by sim ply rotating the polarization of the light injected into the cavity, we can switch to s-polarization, where we measure using the ring-down m ethod a much higher nesse of F = 170000 corresponding to 2 21 kH z. The round-trip length of the ring-

E lectronic address: kruse@ pit.physik.uni-tuebingen.de





FIG. 1: Geometry of the ring-cavity. The atom ic cloud is located in the free-space waist of the cavity mode. The system is characterized by the pumping parameter  $_{\rm i}$ , the atom - eld coupling g, the laser detuning with respect to the cavity  $_{\rm c}$  and to the atom ic resonance  $_{\rm a}$ , and the decay width of the atom and of the cavity eld .

cavity is L = 85 mm, the beam waists in horizontal and in vertical direction at the location of the MOT are  $w_v = 129 \, \text{m}$  and  $w_h = 124 \, \text{m}$ , respectively. This yields a cavity mode-volume of  $V_{m \ ode} = \frac{1}{4}Lw_vw_h = 1 \, \text{mm}^3$ . The intracavity power P, largely enhanced by the factor F= to values around 10 W, gives rise to an optical potential with a depth of  $k_B \,$  1.4 mK at the wavelength 799 nm. The radial and axial secular frequencies in the harm onic region close to the center of the trap are  $!_{rad}=2 \,$ = 640 H z and  $!_{ax}=2 \,$ = 450 kH z.

The ring-cavity is driven by a titanium -sapphire (T iSa) laser delivering up to 2 W output power into three opticalm odes separated by 12 G H z [16]. The centralm ode is ltered by an external confocal Fabry-Perot etalon. The T iSa laser is locked to one of the eigenfre-

FIG. 2: Experim ental setup for pumping both directions of the ring-cavity while locking the laser to a resonator eigenfrequency via the Pound-D rever-H all technique. The AOM and the piezo rule out the laser frequency uctuations. A lso shown are the dem odulated re ection signal (a) and the transm ission signal (b) of the cavity.

quencies of the ring cavity using the Pound-D rever-H all boking technique [17, 18] (see Fig. 2). A feedback servo drives a piezo-electric transducer mounted in the T iSa laser cavity, whose frequency response is limited to 10 kH z. Faster uctuations of the laser frequency are balanced by means of an external double-passed acoustooptic modulator (AOM). The servo bandwidth of 1 M H z is limited by a 100 ns time delay in the response of the AOM. The laser frequency is stable enough to yield intensity uctuations observed in the cavity transmission signal below 2 % even in the high nesse case.

The dipole trap is permanently operated, because keeping the T isa laser locked requires a certain am ount of light inside the cavity. The standing wave dipole trap is loaded from a spatially overlapping MOT for a period of 15 s. Before switching o the MOT, we apply a 40 m s tem poral dark MOT stage [19] by increasing the detuning of the MOT laser beams to 90 MHz and reducing the intensity of the repumping beams. For the conditions given above we typically capture 3  $10^7$  atom s distributed over 10000 antinodes of the standing wave. The tem perature of the atom ic ensemble is measured using the TOF method. We suddenly switch o the dipole trap and in age the shadow that the cloud in prints on a weak probe beam after a period of ballistic expansion. During the expansion, the initial momentum distribution of the cloud evolves into a density distribution, whose radial width yields the tem perature of the cloud. Depending on the potential well depth, we obtain tem peratures between 70 and 280 K, corresponding to roughly 1=5 of the well depth. The peak density is typically 3  $10^{12}$  cm<sup>3</sup>. The

lifetime of the dipole trap measured at 799 nm is 0:5 s. A thorough investigation of trap loss processes is presented in a similar system in Ref. [14].

W ith a slight m isalignment of the incoupled laser beam, the ring-cavity can be locked to higher-order transverse modes, into which the atoms can settle. Fig. 3 shows absorption pictures of atom ic clouds con ned in di erent higher-order transverse modes. Such modes exhibit an enhanced surface-to-volume ratio, which may prove advantageous for forced evaporation.



FIG.3: Absorption pictures of atoms stored in higher-order transverse cavity modes.

W e perform spectroscopy of recoil-induced resonances (RIR) [15, 20, 21], i.e. we probe 2-photon Ram an transitions between two velocity states of the same atom. Two Ram an beam s  $k_1$  and  $k_2$  enclosing a small angle 13:1 are radiated onto the atom ic cloud such that the di erence vector  $q = k_1 k_2$  is oriented nearly parallel to the dipole trap symmetry axis  $\hat{z}$ . The two beams, whose frequencies are  $!_1$  and  $!_2$ , give rise to a standing wave with periodicity 2 =q slow ly moving in  $\hat{z}$  direction with velocity  $v_z = !=q$ , where  $q = (k_1 + k_2) \sin(=2)$  and  $! = !_1 !_2$ . The light wave leads to a periodic dipole potential for the atom s which in our experiment has a well depth of 60 K in units of  $k_B$  . Only atom s satisfying the energy and m om entum conservation requirem ent can undergo Raman transitions, i.e. only atom s moving synchronously to the standing wave can scatter light from one beam into the other. This scattering is monitored as an intensity variation in one of the beams. The net rate for scattering from beam 1 into beam 2 m ay be written as W ( $v_z = !=q$ ) = ~ =2  $^{2}_{\rm B}$  N @ = Q y w here (v<sub>z</sub>) denotes the Maxwell-Boltzmann momentum distribution, N the number of atom s and  $_{\rm R} = \frac{1}{2} = 2$  with the resonant R abi-frequencies  $\ _{i}$  for each R am an beam . By varying ! the derivative of the M axwell-Boltzm ann momentum distribution is scanned from which the tem perature can be derived [21]. Trace (a) of Fig. 4 shows such RIR-scans recorded on a cloud 200 s after being released from the dipole trap. Trace (b) has been calculated assuming a 100 K cold cloud.

The scan rate must be judiciously chosen [22]. If the scan rate is too slow, the atoms are notably redistributed between the velocity classes while scanning. The above expression shows that the scattering process preferentially occurs tow ards higher velocities, so that al3

though the momentum transfer is quite small, the cloud is slightly heated under the in uence of the R am an beam s. If, on the other hand, the scan rate is too fast the signal can be strongly distorted and a ringing type oscillation is observed [23]. For an untrapped cloud of atom s rst indication of ringing can be observed for scan rates above 2:1 kH z/ s. For trapped atom s, how ever, ringing is already very pronounced at this rate (Fig. 4c). Ringing is a general feature of resonant phenom ena and can be observed for a simple mechanical oscillator as well as in an optical resonator or for standard two level quantum systems. The critical scan rate for ringing is roughly given by the square of the linew idth. For slower scan rates the system can follow adiabatically, i.e. the sweep time across the resonance is longer than the inverse linew idth.



FIG. 4: Trace (a): Spectrum of RIR resonances recorded on free atoms. The intensity of Raman beam with  $!_2$  is m easured, while the frequency of beam with  $!_1$  is scanned. Both beam swere detuned by 110 M H z from resonance and their peak intensities were 50 mW  $/cm^2$ . The scan rate was 2:1 kHz/s. Trace (b): Calculated transition rate for the same param eters assuming a 100 K cold cloud. Trace (c): Sam e conditions but recorded on trapped atoms. The trap had a well-depth of  $U_0 = h$  30 M H z =  $k_B$  1:6 m K corresponding to secular frequencies  $!_z = 2$ 700 kH z and  $!_{r} = 2$ 1 kHz. Trace (d) shows a simulation (see text) of the susceptibility of a two-level atom subject to a laser quickly swept over its resonance. The chosen decay rate was = 25 kHz and the Rabi-frequency = 0:1.

In our experiment we observe that the critical scan rate is dram atically reduced if the atom s are trapped in the standing wave dipole trap. A qualitative explanation of this e ect is possible by using the following simple picture. For an untrapped cloud of atom s the potential generated by the R am an beam s has its principle e ect on the atom s near zero detuning. Then, the speed of the potential wave is su ciently slow for the atom s to react and form a standing wave like density pattem that m irrors the shape of the potential wave. For positive detunings the potential wave starts moving with increasing speed relative to the static atom ic distribution and the atom s within the density modulation are subjected to an oscillating dipole force. This force is accompanied by a periodic photon redistribution between the Ram an beam s which is observed in the experiment. The ringing fades away as the atom ic density modulation slow ly decays due to therm al drifts of the atom s. For trapped atom s this picture does not seem to apply since the atom s are strongly con ned within the antinodes of the resonator dipole trap such that a density pattern can not form . However, if we neglect the weak con nem ent in radial direction and thus regard a therm al distribution of atoms that can still propagate freely along the valleys of the resonator dipole trap, this is only true if q is strictly parallel to the symmetry axis of the dipole trap. W ith only a slight m isalignment of q by an angle

the atom s can follow the force of the R am an potential by travelling along those valleys. In order to form a density modulation similar to the untrapped case the atom s now have to propagate a distance that is longer by a factor  $f = 1 = \sin A$ . Similarly, the lifetime of the m odulation pattern is enhanced by the same factor since the atom s have to drift over a longer distance with the sam e therm al velocity. The critical scan rate for ringing is therefore reduced by the factor f<sup>2</sup>. The sam e situation can be explained from a di erent perspective. Since the phase speed of the R am an potential along the valleys is enhanced by f the atom s that interact with the R am an potential have to move with a velocity given by f !=q. The resulting reduction of the linew idth by 1=f can easily be understood by the fact that for an axially con ned atom undergoing a 2-photon Ram an transition only the radialm om entum com ponent with  $q_r = q = f$  has to satisfy the momentum conservation requirement. From the data we can estim ate the coherence lifetim e to be 100 s. This compares well to the above described therm aldecay of the density distribution if we assume a misalignment angle of = 3.

A full quantitative description is beyond the scope of this paper, how ever it is interesting to note that the data can be well described by the dynam ics of a degenerate two level quantum system. The lower curve in Fig. 4 (d) shows the coherence Im 12 calculated by num erical integration of two-level B loch-equations ( is the density matrix). The resonance occurs at ! = 0. At the beginning of the scan, where ! is far from resonance, the population of the excited level 22 is just too sm all and the system does not react to the light eld. As soon as ! passes through the resonance, the coherence 12 is excited and can now be driven by the laser even when ! is tuned far away. The two levels can be identied with two velocity states that are coupled by the R am an-beam s. In the experim ent, the therm ald istribution causes an inhom ogeneous broadening that is accounted for in the simple

two levelm odel by introducing an e ective decay rate .

To conclude we introduced our ring-cavity standing wave dipole trap and characterized it by lifetim e and by tem perature m easurem ents. W e have show n that by driving recoil-induced resonances, we can excite and probe the motion of atoms trapped in the ring-cavity eld. In the near future we plan to look for the expected feedback of the atom ic motion on the standing light wave and for interatom ic coupling induced by the standing wave. It is also tempting to explore the predicted new types of cavity cooling in view of their aptitude of cooling below the threshold of quantum degeneracy. Bose-Einstein condensates are very appealing objects in the context of ring-cavity studies. E.g. M eystre and cow orkers [25] have discussed the use of ring-cavities for recycling superradiant light pulses produced by Rayleigh-scattering o condensates [26] and predict for such system s the possibility ofmutual coherent quantum control between optical and m atter-w ave m odes.

We acknowledge nancial support from the Landesstiftung Baden-Wurttemberg and the Deutsche Forschungsgemeinschaft.

- [L] A. Mosk, S. Jochim, H. Moritz, Th. Elsasser, M. Weidem uller, and R. Grimm, Opt. Lett. 26, 1837 (2001).
- [2] C J. Hood, T W .Lynn, A C. Doherty, A S. Parkins, and H J. K im ble, Science 287, 1447 (2000).
- [3] P.W. H. Pinske, T. Fischer, P. Maunz, and G. Rempe, Nature 404, 365 (2000).
- [4] A. Sinatra, J.F. Roch, K. Vigneron, Ph. Grelu, J.-Ph. Poizat, K. W ang, and P. Grangier, Phys. Rev. A 57, 2980 (1998).
- [5] T W . Mossberg, M. Lewenstein, and D J. Gauthier, Phys. Rev. Lett. 67, 1723 (1991).
- [6] J.I. Cirac, M. Lewenstein, and P. Zoller, Phys. Rev. A 51, 1650 (1995).
- [7] P.Horak, G.Hechenblaikner, K.M. Gheri, H. Stecher, H. Ritsch, Phys. Rev. Lett. 79, 4974 (1997).
- [8] V.Vuletic and S.Chu, Phys.Rev.Lett.84, 3787 (2000).
- [9] M.Gangland H.Ritsch, Phys. Rev. A 61, 043405 (2000).

- [10] M.Kozum a, K.Nakagawa, W.Jhe, and M.Ohtsu, Phys. Rev.Lett. 76, 2428 (1996).
- [11] G.Raithel, W D.Phillips, S.L.Rolston, Phys. Rev. Lett. 81, 3615 (1998).
- [12] A.Hemmerich, Phys.Rev.A 60, 943 (1999).
- [13] M. Gangl, P. Horak, and H. Ritsch, J. Mod. Opt. 47, 2741 (2000).
- [14] B. Nagorny, Th. Elsasser, H. Richter, A. Hemmerich, D. Kruse, C. Zimmermann, and Ph.W. Courteille, Phys. Rev. A (accepted) (2003).
- [15] J.Y. Courtois, G. Grynberg, B. Lounis, and P. Verkerk, Phys. Rev. Lett. 72, 3017 (1994).
- [16] C. Zimmermann, V. Vuletic, A. Hemmerich, L. Ricci, and T.W. Hansch, Opt.Lett. 20, 297 (1995).
- [17] R. Pound, Rev. Sci. Instr. 17, 490 (1946).
- [18] R W P.D rever, JL. Hall, F V. Kowalski, J. Hough, G M. Ford, A J. M unley, and H W. W ard, Appl. Phys. B 31,

97 (1983).

- [19] W .Ketterle, K B.Davis, M A.Jo e, A.Martin, and D E. Pritchard, Phys. Rev. Lett. 70, 2253 (1993).
- [20] J. Guo, P.R. Berman, B. Dubetsky, and G. Grynberg, Phys. Rev. A 46, 1426 (1992).
- [21] D.R.M eacher, D.Boiron, H.M etcalf, C.Salom on, and G.Grynberg, Phys.Rev.A 50, R1992 (1994).
- [22] M C. Fischer, A M. Dudarev, B. Gutierrez-Medina, and M G. Raizen, J. Opt. B; Quantum Sem iclass. Opt. 3, 279 (2001).
- [23] Sim ilar observations have been made in the case of untrapped atom s for a very rapid change of the detuning ! from zero to a non-zero value [24]. In this case, the

observed oscillations have been explained with the developm ent of a stationary atom ic density grating periodically sensed by the Raman standing wave moving with velocity !.

- [24] M.Kozum a, Y.Im ai, K.Nakagawa, and M.Ohtsu, Phys. Rev. A 52, R3421 (1995).
- [25] M G. M oore and P. M eystre, Phys. Rev. Lett. 83, 5202 (1999).
- [26] S. Inouye, A.P. Chikkatur, D.M. Stamper-Kum, J. Stenger, D.E. Pritchard, and W. Ketterle, Science 285, 571 (1999).