

## Generation of entangled states of two three-level atoms in cavity QED

XuBo Zou, K. Pahlke and W. M. Mathis  
 Electromagnetic Theory Group at THT,  
 Department of Electrical Engineering,  
 University of Hannover, Germany

We present a scheme to generate maximally entangled states of two three-level atoms with a nonresonant cavity by cavity-assisted collisions. Since the cavity field is only virtually excited no quantum information will be transferred from the atoms to the cavity.

PACS numbers: 03.67.-a, 03.65.-w, 42.50.-p

Entangled states of quantum particles give the possibility to test quantum mechanics against a local hidden variable theory [1, 2, 3]. They also have practical applications in quantum cryptography [4], quantum dense coding [5] and quantum teleportation [6]. Most of the research in quantum information processing is based on entanglement generation of quantum two-level systems (Qubits), which represent the information. Recently, there is much interest in  $N$ -level quantum systems ( $N \geq 3$ ) to represent information. It was demonstrated that key distributions based on three-level quantum systems are more secure against eavesdropping than those based on two-level systems [7]. Key distribution protocols based on entangled three-level systems were also proposed [8]. The security of these protocols is related to the violation of the Bell inequality. Recently, it was shown that the quantum prediction differs more radically from classical physics in the case of three-level systems than in the case of two-level systems. The three-level system provides in this context a much smaller level of noise [9]. The proof of Bell's theorem without the inequalities by Greenberger, Home, and Zeilinger (GHZ) was extended to multiparticle multi-level systems [10]. One way to generate multiqubit entanglement of  $N$ -level quantum systems is to use higher order parametric down conversion [11]. A more direct way is to use multilevel quantum systems. In this context entanglement generation of  $N$ -level quantum systems was reported [12, 13]. Rydberg atoms which cross superconductive cavities are an almost ideal system to generate entangled states and to perform small scale quantum information processing [14]. A number of schemes were proposed in the context of cavity QED. In particular EPR pairs [15] and GHZ states [16] were successfully generated by a successive interaction of a series of atoms in a cavity field. In these schemes cavities act as memories, which store the information of an electric system and transfer it back to the electric system. Thus, the decoherence of the cavity field becomes one of the main obstacles for the implementation of quantum information in cavity QED. Recently, significant progress was made by proposals for atoms, which interact with a nonresonant cavity [17]. In this theoretical scheme it is suggested to use a dynamic, which involves a virtual exchange of a photon with the field. To the first order of the approximation the scheme is insensitive to cavity losses or to the presence of a stray of a thermal field in the mode. Recently, this process was also suggested to generate GHZ states [9] and to implement the quantum search algorithm [20]. Following the proposal of Ref [17] an experiment was performed in which two Rydberg atoms cross a nonresonant cavity. These atoms became entangled in a controllable way by cavity-assisted collisions [18].

In this paper we show that cavity-assisted collisions of two Rydberg atoms can be used to entangle their electronic states  $|f_i\rangle$ ,  $|e_i\rangle$  and  $|g_i\rangle$  in the related three-level model (see Fig.1 (a)). The experimental setup is shown in Fig.1 (b). The Fabry-Perot resonator, which is denoted by cavity C, sustains a resonant cavity mode of frequency  $\omega_a$ . The  $|e_i\rangle \leftrightarrow |g_i\rangle$  and  $|f_i\rangle \leftrightarrow |g_i\rangle$  transitions are at 51.1 and 54.3 GHz, respectively. The cavity mode is shifted in the frequency from the transitions  $|e_i\rangle \leftrightarrow |g_i\rangle$  and  $|f_i\rangle \leftrightarrow |g_i\rangle$  by detunings  $\epsilon_g$  and  $\epsilon_f = \epsilon_g + \epsilon_{\text{det}}$ . The value  $\epsilon_{\text{det}} = 32$  GHz is the frequency difference of the transitions  $|e_i\rangle \leftrightarrow |g_i\rangle$  and  $|f_i\rangle \leftrightarrow |g_i\rangle$ . The experimental values, which are given in Ref [4] show  $\epsilon_g \approx \epsilon_f$ . Thus, we can choose the cavity frequency in a way that only the levels  $|e_i\rangle$  and  $|g_i\rangle$  are appropriately affected by the nonresonant atom-field coupling. The quantum state  $|f_i\rangle$  will in a good approximation not be affected during the atom-cavity interaction.

We write the interaction Hamiltonian inside of the cavity in the interaction picture

$$H = g[e^{-i\epsilon_g t} a^\dagger (\hat{\sigma}_1^- + \hat{\sigma}_2^-) + e^{i\epsilon_g t} a (\hat{\sigma}_1^+ + \hat{\sigma}_2^+)]: \quad (1)$$

Where  $\hat{\sigma}_j^- = |g_j\rangle\langle e_j|$  and  $\hat{\sigma}_j^+ = |e_j\rangle\langle g_j|$ ,  $a$  and  $a^\dagger$  are the annihilation and creation operator of the cavity field. The atom-cavity coupling strength is denoted by  $g$ . In the large detuning case of  $\epsilon_g \gg g$  no energy exchange between the atoms and the cavity will happen. The effective Hamiltonian is given by [17]

$$H = \sum_{i,j=1}^2 \chi_{ij}^2 (\hat{\sigma}_i^+ \hat{\sigma}_j^- a a^\dagger - \hat{\sigma}_i^- \hat{\sigma}_j^+ a^\dagger a); \quad (2)$$

where  $\Omega = g^2 = \Omega_{eg}$ . If the cavity field is at the beginning in the vacuum state the effective Hamiltonian (2) reduces to

$$H = (\sigma_{1+}\sigma_{1-} + \sigma_{2+}\sigma_{2-} + \sigma_{1+}\sigma_{2-} + \sigma_{2+}\sigma_{1-}): \quad (3)$$

The first two terms describe the Stark shift in the vacuum cavity. The dipole coupling between the two atoms, which is induced by the cavity, are considered by the other terms. In order to generate maximally entangled states of two three-level atoms, we assume that two atoms are initially prepared in the state  $|j_1 i j_2 i\rangle$ . The atom 1 crosses two classical fields, which are tuned to the transitions  $|j_1\rangle \leftrightarrow |j_i\rangle$  and  $|j_i\rangle \leftrightarrow |j_f\rangle$ , respectively. By choosing the amplitudes and phases of the classical fields appropriately this atom becomes prepared in the state  $\frac{1}{3}|j_1 i\rangle + \frac{2}{3}|j_i i\rangle$ . Then both atoms are simultaneously sent into the cavity C, which is in the vacuum state. The interaction is described by the effective Hamiltonian (3), which causes no effect on the state  $|j_1 i j_2 i\rangle$ . After the interaction time  $t_1$  the quantum state

$$|t_1\rangle = \frac{1}{3}|j_1 i j_2 i\rangle + \frac{2}{3}e^{-i t_1} [\cos(t_1)|j_i i j_2 i\rangle - i \sin(t_1)|j_1 i j_2 i\rangle] \quad (4)$$

is obtained. With the choice of  $t_1 = \pi/2$  the state (4) becomes

$$|t_1 = \pi/2\rangle = \frac{1}{3}|j_1 i j_2 i\rangle + \frac{2}{3}|j_1 i j_2 i\rangle: \quad (5)$$

The atom 2 is then addressed by a classical microwave pulse, which is tuned to the transition  $|j_i\rangle \leftrightarrow |j_f\rangle$ . This step requires separate addressing of the atoms. Since microwave field can not be focused narrow enough to address atom 2 without affecting atom 1 a method should be used, which makes the atomic transitions of the two atoms slightly different. An appropriate method was proposed by Yamaguchi et al. [20], which is based on the Stark effect. The authors suggest to use a set of electrodes in the cavity, which create an inhomogeneous electric field. Since the two atoms are located in space regions with different electric field strength their frequencies become independently controllable. The amplitudes and phases of the classical fields have to be chosen appropriately in order to let the atom 2 undergo the transition  $|j_2 i\rangle \rightarrow |j_f i\rangle$ ;  $|j_2 i\rangle \rightarrow |j_i i\rangle$ . Here we assume that the classical microwave field is strong enough so that the nonlinear interaction between atoms can be neglected during this stage. After this operation the state (5) becomes

$$|t_1 = \pi/2\rangle = \frac{1}{3}|j_1 i j_2 i\rangle + \frac{2}{3}|j_1 i j_2 i\rangle: \quad (6)$$

Then the classical microwave field is switched off and the evolution of the system is determined by the interaction (3). After another interaction time  $t_2$  the system's time evolution has transformed the state (6) to the state

$$|t_1 + t_2\rangle = \frac{1}{3}|j_1 i j_2 i\rangle + \frac{2}{3}e^{-i t_2} [\cos(t_2)|j_1 i j_2 i\rangle - i \sin(t_2)|j_i i j_2 i\rangle]: \quad (7)$$

If we choose  $t_2 = \pi/4$  the quantum state

$$|t_1 + t_2 = \pi/4\rangle = \frac{1}{3}(|j_1 i j_2 i\rangle + e^{-i \pi/4}|j_1 i j_2 i\rangle - i e^{-i \pi/4}|j_i i j_2 i\rangle) \quad (8)$$

will be obtained. After the two atoms left the cavity C the atom 2 crosses a classical field, which is tuned to the transition  $|j_i\rangle \leftrightarrow |j_f\rangle$ . If the amplitude and the phase of the classical field is chosen appropriately the atom 2 will undergo the transition  $|j_2 i\rangle \rightarrow e^{i \pi/4}|j_f i\rangle$ ;  $|j_2 i\rangle \rightarrow e^{i \pi/4}|j_i i\rangle$ . Thus, the state (8) becomes

$$|_{\text{bell}} = \frac{1}{3}(|j_1 i j_2 i\rangle + |j_1 i j_2 i\rangle + |j_i i j_2 i\rangle): \quad (9)$$

This is the maximally entangled state of a two three-level system, which is discussed in Ref[9].

One of the difficulties of this scheme is the requirement to send two atoms simultaneously through the cavity, otherwise an error will result. In the following, we discuss the case, that the second atom enters the cavity in the excited state before the first atom. This deviation from the ideal case shall be considered with the time difference  $\tau$ , which denotes a fraction of the Rabi frequency  $\Omega = \Omega_{eg} = g^2$  of the Hamiltonian (2). Then the quantum state

$$|t\rangle = \frac{1}{3}|j_1 i j_2 i\rangle + \frac{2}{3}e^{-i \tau} [\cos(\frac{\tau}{4})|j_1 i j_2 i\rangle + \sin(\frac{\tau}{4})|j_i i j_2 i\rangle] \quad (10)$$

will be generated. The difference of the state  $|0\rangle$  to the state  $|9\rangle$  can be characterized in terms of the fidelity  $F = \langle \psi_{\text{Bell}} | \psi \rangle^2$ :

$$F = \frac{5 + 4 \cos 2}{9} : \quad (11)$$

If  $\theta = 0.01$  holds, we have  $F = 0.999$ . In this case the operation is only slightly affected.

It is necessary to give a brief discussion on the experimental realization of the proposed scheme. It was reported that the cavity can have a photon storage time of  $T = 1 \text{ ms}$  (corresponding to  $Q = 3 \times 10^8$ ). The radiative time of the Rydberg atoms with the principle quantum numbers 49, 50 and 51 is about  $3 \times 10^{-2} \text{ s}$  [14]. The coupling constant of the atoms to the cavity field is  $g = 2\pi \times 25 \text{ kHz}$  [18]. In order to achieve a good entanglement in the cavity-assisted collision process, the detuning  $\Delta_{\text{eg}}$  should be much bigger than  $g$ . With the choice  $\Delta_{\text{eg}} = 10g$  the interaction time between the atom and the cavity field is in the order  $3 \times 10^{-4} \text{ s}$ . The time needed for the classical field pulse is at this scale negligible. Thus, the interaction time needed to complement the total procedure is much shorter than the radiative time and the photon lifetime  $1 \text{ ms}$  in the present cavity. For the interaction time  $1.5 \times 10^{-4} \text{ s}$  the velocity of the prepared atoms should be  $v_p \approx 0.7 \times 10^4 \text{ L/s}$ , where  $L$  is the length of the cavity. If we choose  $L = 2.75 \text{ cm}$  the velocity of the atoms should be of the order of  $192 \text{ m/s}$ , which is in the range of present experiments. Based on cavity QED techniques the present scheme seems to become realizable in a near future.

In summary, we have proposed a scheme to generate maximally entangled states of two three-level atoms. During the passage of the atoms through the cavity field they are only virtually excited. No transfer of quantum information will happen between the atoms and the cavity. The experimental implementation of the scheme demonstrates the power of cavity QED to manipulate complex entangled states for quantum information processing.

- 
- [1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. 47, 777 (1935).
  - [2] J. S. Bell, Physics (Long Island City, N.Y.) 1, 195 (1965).
  - [3] D. M. Greenberger, M. Home, A. Shimony, and A. Zeilinger, Am. J. Phys. 58, 1131 (1990).
  - [4] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991).
  - [5] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. 69, 2881 (1992).
  - [6] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. Wootters, Phys. Rev. Lett. 70, 1895 (1993).
  - [7] Mohamed Bourennane, Anders Karlsson, and Gunnar Björk, Phys. Rev. A 64, 012306 (2001); D. Bruss and C. Macchiavello, Phys. Rev. Lett. 88, 127901 (2002); Nicolas J. Cerf, Mohamed Bourennane, Anders Karlsson, and Nicolas Gisin, Phys. Rev. Lett. 88, 127902 (2002).
  - [8] Dagomir Kaszlikowski, Kelken Chang, Daniel Kuan LiO i, L. C. Kwek, C. H. Oh, quant-ph/0206170; Thomas Durt, Nicolas J. Cerf, Nicolas Gisin, Marek Żukowski, quant-ph/0207057.
  - [9] D. Kaszlikowski, P. G. Nascimbene, M. Żukowski, W. M. Itskewski, and A. Zeilinger, Phys. Rev. Lett. 85, 4418 (2000). Dagomir Kaszlikowski, Darwin Gosal, E. J. Ling, L. C. Kwek, Marek Żukowski, and C. H. Oh, Phys. Rev. A 66, 032103 (2002); A. Cincin, T. Durt, N. Gisin, and J. I. Latorre, Phys. Rev. A 65, 052325 (2002) Jing-Ling Chen, Dagomir Kaszlikowski, L. C. Kwek, C. H. Oh, and Marek Żukowski, Phys. Rev. A 64, 052109 (2001).
  - [10] Adán Cabello, Phys. Rev. A 63, 022104 (2001); Nicolas J. Cerf et al., Phys. Rev. Lett. 89, 080402 (2002).
  - [11] A. LasLinares et. al. Nature 412, 887 (2001) J. C. Howell et al. Phys. Rev. Lett. 88 030401 (2002)
  - [12] A. Mair, A. Vaziri, G. Weihs, A. Zeilinger, Nature 412, 313 (2001)
  - [13] Hugues de Riedmatten, Ivan Marcikic, Hugo Zbinden, Nicolas Gisin, quant-ph/0204165
  - [14] J. M. Raimond et al., Rev. Mod. Phys. 73, 565 (2001)
  - [15] E. Hagley, X. Maître, G. Nogues, C. Wunderlich, M. Brune, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. 79, 1 (1997).
  - [16] A. Rauschenbeutel, G. Nogues, S. Onghini, P. Bertet, M. Brune, J. M. Raimond, and S. Haroche, Science 288, 2024 (2000).
  - [17] S.-B. Zheng and G.-C. Guo, Phys. Rev. Lett. 85, 2392 (2000).
  - [18] S. Onghini et al., Phys. Rev. Lett. 87, 037902 (2001)
  - [19] Shi-Biao Zheng, Phys. Rev. Lett. 87, 230404 (2001) Guo-Ping Guo et al., Phys. Rev. A 65, 042102 (2002).
  - [20] F. Yamaguchi et al., Phys. Rev. A 66, 010302 (2002)

## Figure Captions

Figure 1.(a) This figure shows the electronic levels of the three-level atom model in the energy representation. Figure 1.(b) This figure shows the experimental apparatus. The atoms 1 and 2 cross the cavity with the same velocity but at different positions with a different electric field strength. This makes an individual manipulation of each atom by a classical field possible. Inside the cavity atom 2 is manipulated by the classical field  $S$ . Outside the cavity both atoms are manipulated by the classical fields  $R_1$  and  $R_2$ , respectively.