

BBGKY Hierarchy Underlying Many Particle Quantum Mechanics

G. Kaniadakis

Dipartimento di Fisica and Istituto Nazionale di Fisica della Materia,
Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy
(Dated: March 29, 2024)

Recently, the one particle quantum mechanics has been obtained in the framework of an entirely classical subquantum kinetics. In the present paper we argue that, within the same scheme and without any additional assumption, it is possible to obtain also the n -particle non relativistic quantum mechanics. The main goal of the present effort is to show that the classical BBGKY hierarchical equation, for the n -particle reduced distribution function, is the ancestor of the n -particle Schrodinger equation. On the other hand we show that within the scenario of the subquantum structure of quantum particle, the Fisher information measure emerges naturally in quantum mechanics.

PACS numbers: 03.65.Ta, 05.20.Dd

In ref. [1], it is shown that quantum mechanics can be obtained in a self consistent scheme of an entirely classical many body physics. Within this framework the main features of quantum mechanics (i.e. the probabilistic nature of the quantum description, the quantum potential, the Schrodinger equation, the quantum operators, the Heisenberg uncertainty principle) were obtained starting from a classical subquantum kinetics in the phase space and without invoking any additional principle. Furthermore the fundamental constant \hbar emerges naturally as an integration constant and represents a free parameter for the theory. Within this theory, the astonishing scenario of a subquantum structure for the quantum particle emerges spontaneously. The quantum particle turns out to have an internal structure and a spatial dispersion and appears to be composed by N identical point like and interacting subquantum objects, the monads, obeying to the laws of Newtonian physics. The theory can be viewed as describing a mechanism which permits to construct the quantum particle starting from its constituents and then quantum mechanics appears to describe a physical reality.

The statistical ensemble of these N monads is described in the phase space through the distribution function $f = f(t; \mathbf{x}; \mathbf{v})$ whose time evolution is governed by the kinetic equation $Lf = C(f)$, being L the Liouville operator and $C(f)$ the collision integral that takes into account the intermonad interactions. We don't make any assumption about the structure of the collision integral and the nature of the interaction between the monads except that, during the point-like collisions the monad number, momentum and energy are conserved. The projection of the above phase space kinetics into the physical space produces a hydrodynamics which naturally leads to the one particle spinless quantum mechanics when the external force field is conservative.

Recently in ref. [2], it is shown that the above kinetic

approach can be easily adopted to treat the quantum particle with spin in an external electromagnetic field. Of course in this way the spinning quantum mechanics can be obtained.

The energy conservation during the collisions considered in the present scheme plays a very important role and could be also taken into account in the stochastic models of quantum mechanics as it has been recently observed in ref. [3].

A question which naturally arises at this point is if it is possible to extend the kinetic formalism of ref. [1] in order to treat non relativistic many particle quantum systems. This point is neglected in the literature where almost the totality of works devoted to the derivation and interpretation of quantum mechanics concern the single particle.

The main goal of the present paper is to show that it is possible to obtain, within the theory developed in ref. [1] and without any additional assumption, also the n -particle quantum mechanics starting from the classical kinetics governed in the phase space by the BBGKY hierarchy.

BBGKY Hierarchy: Let us consider a statistical ensemble constituted by N identical and indistinguished point-like particles (monads) obeying the laws of classical physics. It is well known that the n -particle reduced distribution function $f_n = f_n(t; \mathbf{x}_1; \mathbf{v}_1; \mathbf{x}_2; \mathbf{v}_2; \dots; \mathbf{x}_n; \mathbf{v}_n)$ obeys the following BBGKY hierarchical evolution equation [4]

$$\frac{\partial f_n}{\partial t} + \sum_{i=1}^n \mathbf{v}_i \frac{\partial f_n}{\partial \mathbf{x}_i} + \sum_{i=1}^n \mathbf{F}_i \frac{\partial f_n}{\partial \mathbf{v}_i} = \sum_{i=1}^n C_i(f_{n+1}) ; \quad (1)$$

being $\mathbf{F}_i = -\nabla V_i = -\nabla V(\mathbf{x}_i)$ and

$$\mathbf{V}_i = V^{(ex)}(\mathbf{x}_i) + \frac{1}{2} \sum_{\substack{j=1 \\ j \neq i}}^n V^{(in)}(\mathbf{x}_i - \mathbf{x}_j) ; \quad (2)$$

$C_i(f_{n+1})$ are the collision integrals which can be expressed in term of the function f_{n+1} . The assumption that during the point-like collisions the monad number,

momentum and energy are conserved, implies that the three functions $g_1(v_i) = 1$, $g_2(v_i) = v_i$ and $g_3(v_i) = v_i^2$ are the collision invariants of $C_i(f_{n+1})$

$$\int_{\mathbb{R}^3} g_k(v_i) C_i(f_{n+1}) d^3v_i = 0 ; k = 1;2;3 : \quad (3)$$

Let us consider the 6n-dimensional phase space (x, v) being $x = (x_1; x_2; \dots; x_n)$ and $v = (v_1; v_2; \dots; v_n)$. We introduce the two gradient operators $\partial = \partial_x = (\partial = \partial_{x_1}; \partial = \partial_{x_2}; \dots; \partial = \partial_{x_n})$ and $\partial = \partial_v = (\partial = \partial_{v_1}; \partial = \partial_{v_2}; \dots; \partial = \partial_{v_n})$. Then we define the 3n-dimensional force $F = (F_1; F_2; \dots; F_n)$ which can be derived from a potential

$$F = -\frac{\partial V}{\partial x} ; V = \sum_{i=1}^N V_i ; \quad (4)$$

The BBGKY hierarchical equation (1), governing the evolution of the n-particle reduced distribution function $f_n = f_n(t; x; v)$, can be written in the following compact form

$$\frac{\partial f_n}{\partial t} + v \frac{\partial f_n}{\partial x} + \frac{F}{v} \frac{\partial f_n}{\partial v} = C(f_{n+1}) : \quad (5)$$

The collision integral

$$C(f_{n+1}) = \sum_{i=1}^N C_i(f_{n+1}) ; \quad (6)$$

admits the three collision invariants $g_1(v) = 1$, $g_2(v) = v$ and $g_3(v) = v^2$ satisfying the conditions

$$\int_{\mathbb{R}^3} g_k(v) C(f_{n+1}) d^3v = 0 ; k = 1;2;3 : \quad (7)$$

as one can verify immediately starting from Eq. (3).

Hydrodynamics: We consider now the projection of the kinetics from the 6n-dimensional phase space into the 3n-dimensional space of the coordinates x , where the system is described through the distribution function $f_n(t; x) = \int_{\mathbb{R}^3} f_n(t; x; v) d^3v$. After recalling that the total value of a given physical quantity with density $A = A(t; x; v)$ can be calculated as $\int_{\mathbb{R}^3} A f_n d^3v d^3x$ we define the mean value of A in the point x through

$$\langle A(t; x; v) \rangle_v = \frac{\int_{\mathbb{R}^3} A(t; x; v) f_n(t; x; v) d^3v}{\int_{\mathbb{R}^3} f_n(t; x; v) d^3v} : \quad (8)$$

This mean value represents the density of A in the space of the coordinates and permits to write the total value of A as $\int_{\mathbb{R}^3} \langle A \rangle_v d^3x$. We define the densities in the space of the coordinates for some quantities used in following. The densities of current, of stress tensor and of heat flux vector are defined, respectively, by

$$u = \langle v \rangle_v ; \quad (9)$$

$$ij = \langle (v_i u_i)(v_j u_j) \rangle_v ; \quad (10)$$

$$h_i = \frac{1}{2} \langle j v u_j^2 (v_i u_i) \rangle_v : \quad (11)$$

The density of energy is given by

$$E = \frac{1}{2} \langle v^2 \rangle_v + V = \frac{1}{2} u^2 + \dots + V ; \quad (12)$$

being $\dots > 0$ the density of the internal energy.

$$\dots = \frac{1}{2} \langle v^2 \rangle_v - \frac{1}{2} \langle v \rangle_v^2 : \quad (13)$$

Using the standard procedure consisting in multiplying Eq. (5) by the three collision invariants $g_k(v)$ and integrating with respect to v , the three following hydrodynamic equations can be obtained [5, 6]:

$$\frac{\partial}{\partial t} \langle f_n \rangle + \frac{\partial}{\partial x} (\dots u) = 0 ; \quad (14)$$

$$\frac{D u}{D t} = F + \dots ; \quad (15)$$

$$\frac{\partial}{\partial t} (\dots E) + \frac{\partial}{\partial x} (\dots s) = 0 : \quad (16)$$

In the above equations $D = D t = \partial = \partial_t + u \partial = \partial_x$ is the total time or substantial or Lagrangian derivative, F is the stress force

$$F_i = - \frac{1}{n} \frac{\partial}{\partial x_j} (\dots ij) ; \quad (17)$$

and $s_i = E u_i + \dots ij u_j + h_i$ is the energy flux density vector.

Eqs (14)–(16) define the most general hydrodynamics of the particle system which behaves in the physical space as a fluid. In particular Eq. (16) imposes the conservation of the total energy H of the system

$$H = \int_{\mathbb{R}^3} E_n d^3x : \quad (18)$$

After recalling that the external force F is conservative, from Eq. (15) it follows immediately that the irrotationality of u implies the conservativity of F and viceversa, namely

$$u = - \frac{1}{\rho} \frac{\partial S}{\partial x} ; \quad (19)$$

$$F = - \frac{\partial W}{\partial x} : \quad (20)$$

The condition (19) imposes that the system here considered is spinless and reduces the vector equation (15) into a scalar one

$$\frac{\partial S}{\partial t} + \frac{1}{2} \frac{\partial S}{\partial x}^2 + W + V = 0 : \quad (21)$$

Quantum potential: From the two expressions of the force F given by Eqs (17) and (20) it follows immediately

$$\frac{\partial W}{\partial x_j} = \frac{\partial}{\partial x_k} jk + jk \frac{\partial}{\partial x_k} ; \quad (22)$$

with $j = \ln n$. Note that Eq. (22) descends from the requirement that the force F is conservative and can be

viewed as a condition constraining the forms of W and σ_{jk} which result to be two functionals of the field. The solutions of the latter equation are couples of W and σ_{jk} and in principle there can exist more than one solution. Even though it can appear that Eq. (22) contains considerable degrees of freedom, in the following we will show that the particular structure of this equation restricts strongly the number of its solutions.

We suppose now that just as the left hand side, both the first and the second term in the right hand side in Eq. (22) are curl free and will take the form $\partial(\dots)/\partial x_j$. This requirement for the first term can be satisfied in two different ways. The first and most natural way is related to the choice $\sigma_{jk} = \sigma_{jk}$. Alternatively we can pose $\sigma_{jk} = \partial a_k / \partial x_j$ and after recalling the symmetry property $\sigma_{jk} = \sigma_{kj}$ we have that $a_k = \partial \phi / \partial x_k$. Then the most general form of the density of stress tensor is

$$\sigma_{jk} = \sigma_{jk} + \frac{\partial^2 \phi}{\partial x_j \partial x_k} ; \quad (23)$$

being ϕ and σ_{jk} two scalar functionals depending on ψ . After taking into account the identity

$$\frac{\partial^2 \phi}{\partial x_j \partial x_k} \frac{\partial \phi}{\partial x_k} = \frac{\partial \phi}{\partial x_j} \frac{\partial \phi}{\partial x_k} \frac{\partial \phi}{\partial x_k} - \frac{\partial \phi}{\partial x_k} \frac{\partial^2 \phi}{\partial x_k \partial x_j} ; \quad (24)$$

Eq.(22) can be written as

$$\begin{aligned} \frac{\partial}{\partial x_j} W &= \frac{\partial^2 \phi}{\partial x_k \partial x_k} \frac{\partial \phi}{\partial x_k} \frac{\partial \phi}{\partial x_k} - \frac{\partial \phi}{\partial x_j} \frac{\partial \phi}{\partial x_k} \frac{\partial^2 \phi}{\partial x_k \partial x_j} : \end{aligned} \quad (25)$$

Clearly the two terms in the right hand side of Eq. (25) must be curl free. These requirements impose the two following conditions

$$\frac{\partial}{\partial x_l} \frac{\partial \phi}{\partial x_m} = \frac{\partial \phi}{\partial x_m} \frac{\partial \phi}{\partial x_l} ; \quad (26)$$

$$\frac{\partial^2 \phi}{\partial x_l \partial x_k} \frac{\partial \phi}{\partial x_k \partial x_m} = \frac{\partial^2 \phi}{\partial x_m \partial x_k} \frac{\partial \phi}{\partial x_k \partial x_l} : \quad (27)$$

It is trivial to verify that from Eq. (26) it results that $\phi = \phi(\psi)$ is an arbitrary function of ψ while from Eq. (27) one can obtain that $\phi = c$ with c a real arbitrary constant. With these positions Eq. (25) becomes

$$\frac{\partial}{\partial x_j} W - c \frac{\partial^2 \phi}{\partial x_k \partial x_k} \frac{\partial \phi}{\partial x_k} \frac{\partial \phi}{\partial x_k} = 0 : \quad (28)$$

From this last equation we can obtain the expression of W while Eq. (23) gives the expression of σ_{ij} . Finally, we can write the most general solution of Eq. (22) under the form

$$\sigma_{jk} = c \frac{\partial^2 \ln \psi}{\partial x_j \partial x_k} + \sigma_{jk} \frac{1}{\psi} \frac{dU(\psi)}{d\psi} \psi ; \quad (29)$$

$$W = c \frac{\partial^2 \ln \psi}{\partial x^2} + \frac{1}{2} \frac{\partial}{\partial x} \ln \psi + U(\psi) : \quad (30)$$

The above expressions of σ_{jk} and W define a family of solutions of Eq. (22) and represent the hydrodynamic constitutive equations for the system. Remark that these equations have been enforced exclusively from the fact that the internal stress forces are conservative.

In the above constitutive equations the terms containing U describe the standard classical Eulerian fluid. For instance the nonlinear potential $U(\psi) = a \psi$ appearing in the so called cubic non-linear Schrodinger equation is originated from the stress tensor $\sigma_{jk} = \frac{1}{2} a \psi \delta_{jk}$. Analogously the logarithmic non-linearity $U(\psi) = b \ln \psi$ in the Schrodinger equation is originated from a constant stress tensor $\sigma_{jk} = b \delta_{jk}$.

In the following, we pose $U = 0$ and focalize our attention to the term of Eqs (29) and (30), proportional to the arbitrary constant c , which is absent in classical kinetics. From the definition of the total internal energy, it results that $H = \int d^3x \psi^2 > 0$. Starting from Eq. (29) and performing an integration by parts, we obtain $H = c \int d^3x (\partial \psi / \partial x)^2 = N$. Then we pose $c = \hbar^2/4m$ in order to put $H > 0$. The real positive constant \hbar remains a free parameter of the theory.

We introduce the normalized n -particle reduced distribution function $\psi_n = \psi^N$ being $N = \int d^3x \psi$, so that $\int \psi_n d^3x = 1$. Then we define the mass $m = N$ the internal potential $W = N W$ and set $\hbar = N$. From (30) follows the expression of W

$$W = \frac{\hbar^2}{2m} \psi_n^{1/2} \frac{\partial^2 \psi_n^{1/2}}{\partial x^2} : \quad (31)$$

We recall that the potential W given by Eq. (31) is the quantum potential (Madelung 1926) [7, 8] and it is remarkable that here it has been obtained in the framework of an entirely classical subquantum kinetics. Note that the subquantum monad structure of the system generates the stress forces and then the potential W . Clearly when this structure is suppressed the quantum potential vanishes. Furthermore are the presence of the collision integrals C_i in the BBGKY kinetic Eq. (1) which take into account the monad interactions are necessary for the consistency of the theory.

Many particle Schrodinger equation: We define the external potential $V = N V$, and set $S = N S$. Now the quantum fluid is described completely by the two scalar fields ψ_n and S whose evolution equations are (32) and (33), respectively.

$$\frac{\partial \psi_n}{\partial t} + \frac{\partial}{\partial x} \left(\psi_n \frac{1}{m} \frac{\partial S}{\partial x} \right) = 0 ; \quad (32)$$

$$\frac{\partial S}{\partial t} + \frac{1}{2m} \frac{\partial S^2}{\partial x} + W + V = 0 : \quad (33)$$

Alternatively, we can describe this fluid by means of the complex field $\psi_n = \psi_n^{1/2} \exp(iS/\hbar)$ with $\int \psi_n^2 d^3x = 1$, whose evolution equation

$$i\hbar \frac{\partial \psi_n}{\partial t} = \frac{\hbar^2}{2m} \frac{\partial^2 \psi_n}{\partial x^2} + V \psi_n : \quad (34)$$

can be obtained directly by combining Eqs (32) and (33). This later equation is the n -particle Schrodinger equation which, after recalling $V = \sum_{i=1}^n V_i$ with $V_i = N V_i$, assumes the form

$$i\hbar \frac{\partial}{\partial t} \Psi = \sum_{i=1}^n \left[\frac{\hbar^2}{2m} \nabla_i^2 + V_i \right] \Psi \quad (35)$$

The total energy H given by Eq. (18) results to be the sum of two terms: $H = H^{(cl)} + H$. The first term

$$H^{(cl)} = \sum_{i=1}^n \int d^3x \frac{1}{2} m u_i^2 + \sum_{i=1}^n \int d^3x \rho_n V_i(\mathbf{x}) \quad (36)$$

corresponds to the energy of n interacting classical particles while the second term $H = \int d^3x$ is originated by the internal structure of the n bodies and is given by

$$H = \frac{\hbar^2}{8m} \int d^3x \frac{1}{\rho_n} \left(\frac{\partial \rho_n}{\partial \mathbf{x}} \right)^2 = \frac{\hbar^2}{8m} I \quad (37)$$

Eq. (37) hints on the existence of a tight link between quantum mechanics and Fisher information theory. It is remarkable that H results to be proportional to the Fisher information measure I (Fisher 1922) which describes an important feature of quantum mechanics [9, 10, 11, 12]. Once again, within the scenario of the subquantum structure of quantum particle, another basic physical concept, namely I , emerges naturally. In the present context I has a very transparent meaning being simply a measure of the internal energy of quantum system. The fact that H is originated from the internal structure of the n -bodies becomes more transparent if we observe that H can be written also as

$$H = \sum_{i=1}^n \frac{\hbar^2}{8m} \int d^3x_i \frac{1}{\rho_n} \left(\frac{\partial \rho_n}{\partial \mathbf{x}_i} \right)^2 = \frac{\hbar^2}{8m} \sum_{i=1}^n I_i \quad (38)$$

from which it results $I = \sum_{i=1}^n I_i$ and then $H = \sum_{i=1}^n H_i$. An analogous property holds also for the quantum potential $W = \sum_{i=1}^n W_i$ being

$$W_i = \frac{\hbar^2}{2m} \rho_n^{1/2} \frac{\partial^2 \rho_n^{1/2}}{\partial \mathbf{x}_i^2} \quad (39)$$

Of course starting from $H = H^{(cl)} + H$ and taking into account Eqs (35) and (36) one obtains immediately the well known expression of the Hamiltonian of the n -particle quantum mechanics

$$H = \sum_{i=1}^n \int d^3x \frac{\hbar^2}{2m} \left(\frac{\partial \Psi}{\partial \mathbf{x}} \right)^2 + \int d^3x \rho_n V = \sum_{i=1}^n H_i \quad (40)$$

In the present picture the mass of the n -body quantum system is $M = nm$. After recalling that $m = N$ follows immediately that the system is composed by $N = nN$ monads.

Conclusions: The approach used here to obtain the Schrodinger equation for a system of n quantum particles starting from the BBGKY hierarchy describing a statistical system of $N = nN$ subquantum monads, suggests that quantum mechanics can be viewed as a hidden variables theory. This approach can be used easily also to construct the non linear quantum mechanics [it is enough to consider $U \neq 0$ in Eqs (29), (30)] and then for study the non linear effects originated for instance from the particle bosonic or fermionic nature [13, 14].

[1] G. Kaniadakis, Physica A 307, 172 (2002); quant-ph/0112049.

[2] G. Kaniadakis, Found. Phys. Lett. 16 (2), 99 (2003); quant-ph/0209033.

[3] R. C. Zopnik and P. Garbaczewski, Phys. Lett. A 299, 447 (2002).

[4] R. Balescu, Statistical Dynamics: Matter out of Equilibrium, Imperial College Press, London (2000).

[5] R. L. Libo, Kinetics Theory, Prentice-Hall International Editors, N.J. (1990).

[6] R. T. Glassey, The Cauchy problem in Kinetic Theory, SIAM, Philadelphia, PA (1996).

[7] E. Madelung, Z. Phys. 40, 322 (1926).

[8] D. Bohm, Phys. Rev. 85, 166 (1952).

[9] J. Rehacek and Z. Hradil, Phys. Rev. Lett. 88, 130401 (2002).

[10] M. Reginato, Phys. Rev. A 58, 1775 (1998).

[11] B. R. Frieden and B. H. Soer, Phys. Rev. 52, 2274 (1995).

[12] B. R. Frieden, Physics from Fisher Information: A Unification, Cambridge University Press, Cambridge (1999).

[13] G. Kaniadakis, Phys. Rev. A 55, 941 (1997).

[14] G. Kaniadakis and A. M. Scarfone, Phys. Rev. E 64, 026106 (2001).