

Special functions, raising and lowering operators

N. C. Otfas

Faculty of Physics, University of Bucharest, P.O. Box 76-54, Postal Office 76,
Bucharest, Romania, E-mail address: nootfas@yahoo.com

Abstract. The Schrodinger equations which are exactly solvable in terms of associated special functions are directly related to some self-adjoint operators defined in the theory of hypergeometric type equations. The fundamental formulae occurring in a supersymmetric approach to these Hamiltonians are consequences of some formulae concerning the general theory of associated special functions. We use this connection in order to obtain a general theory of Schrodinger equations exactly solvable in terms of associated special functions, and to extend certain results known in the case of some particular potentials.

1. Introduction

It is well-known [2, 4] that, in the case of certain potentials, the Schrodinger equation is exactly solvable and its solutions can be expressed in terms of the so-called associated special functions. Our purpose is to present a general theory of these quantum systems. Our systematic study recovers a number of earlier results in a natural unified way and also leads to new findings.

The number of articles concerning exactly solvable quantum systems and related subjects is very large (see [2, 4, 5] and references therein). Our approach is based on the raising/lowering operators presented in general form (for the first time to our knowledge) by Jafarizadeh and Fakhri [5]. We reobtain these operators in a much simpler way, and use them in a rather different way. More details can be found in [3].

2. Orthogonal polynomials and associated special functions

Many problems in quantum mechanics and mathematical physics lead to equations of hypergeometric type

$$(s)y''(s) + p(s)y'(s) + q(s)y(s) = 0 \quad (1)$$

where $p(s)$ and $q(s)$ are polynomials of at most second and first degree, respectively, and λ is a constant. This equation can be reduced to the self-adjoint form

$$[\sigma(s)y'(s)]' + \tau(s)y(s) = 0 \quad (2)$$

by choosing a function σ such that $[\sigma(s)p(s)]' = \sigma(s)q(s)$. For $\sigma = 1 = \frac{1}{2}l(l-1)$ with $l \in \mathbb{N}$ there exists a polynomial ϕ_l of degree l satisfying (1), that is,

$$(s) \phi_l''(s) + p(s) \phi_l'(s) + q(s) \phi_l(s) = 0 \quad (3)$$

If there exists a finite or infinite interval $(a; b)$ such that

$$(s) \phi_l(s) s^k \Big|_{s=a} = 0 \quad (s) \phi_l(s) s^k \Big|_{s=b} = 0 \quad \text{for all } k \in \mathbb{N} \quad (4)$$

and if $\sigma(s) > 0$, $\rho(s) > 0$ for all $s \in (a; b)$, then the polynomials p_n are orthogonal with weight function $\rho(s)$ in the interval $(a; b)$. In this case p_n are known as classical orthogonal polynomials [7].

Let $p_n(s) = p_n(s)$. By differentiating the equation (3) m times and multiplying it by $p_m(s)$, we get for each $m \in \mathbb{N}$ the associated differential equation which can be written as $H_m p_m = p_m$, where

$$H_m = \sigma(s) \frac{d^2}{ds^2} + \left(\sigma(s) \frac{d}{ds} + \frac{m(m-1)}{4} \frac{\sigma'(s)}{\sigma(s)} + \frac{m}{2} \frac{\rho'(s)}{\rho(s)} \right) p_m - \frac{1}{2} m(m-1) \rho(s) p_m$$

and $p_m(s) = p_m(s) p_m(s)$ are known as the associated special functions. The set $\{p_m; m \in \mathbb{N}\}$ is an orthogonal sequence ([7], pag. 8) in the Hilbert space $L^2_b(\sigma, \rho)$

$$H = \{f : (a; b) \rightarrow \mathbb{R} \mid \int_a^b \sigma(s) f^2(s) ds < \infty \text{ with } h' = \int_a^b \rho(s) f(s) ds = 0\}$$

For each $m \in \mathbb{N}$, let H_m be the linear span of $\{p_m; m \in \mathbb{N}\}$. In the sequel we shall restrict us to the case when H_m is dense in H for all $m \in \mathbb{N}$. For this it is sufficient the interval $(a; b)$ to be finite, but not necessary.

3. Raising and lowering operators. Factorizations for H_m

Lorente has shown recently [6] that a factorization of H_0 can be obtained by using the well-known three term recurrence relation satisfied by p_n and a consequence of Rodrigues formula. Following Lorente's idea we obtain a factorization of H_m by using the definition $p_m(s) = p_m(s) p_m(s)$ and a three term recurrence relation.

Differentiating $p_m(s) = p_m(s) p_m(s)$ we get the relation

$$p_{m+1}(s) = \left(\sigma(s) \frac{d}{ds} + \frac{m}{2} \frac{\rho'(s)}{\rho(s)} \right) p_m(s) \quad \text{for all } m \in \mathbb{N} \quad (5)$$

If we differentiate (3) $m-1$ times and multiply the obtained relation by p_{m-1} then we get for each $m \in \mathbb{N}$ the three term recurrence relation

$$p_{m+1}(s) + \frac{\sigma(s)}{\rho(s)} + 2(m-1) \rho(s) p_m(s) + (1-m) p_{m-1}(s) = 0 \quad (6)$$

and $\frac{\sigma(s)}{\rho(s)} + 2(1-m) \rho(s) p_1(s) + (1-m) p_0(s) = 0$: A direct consequence of these formulae is the relation

$$(1-m) p_m(s) = \left(\sigma(s) \frac{d}{ds} + \frac{\sigma(s)}{\rho(s)} \right) (m-1) \rho(s) p_{m+1}(s) \quad (7)$$

satisfied for all $m \in \mathbb{N}$:

The operators $A_m : H_m \rightarrow H_{m+1}$ and $A_m^+ : H_{m+1} \rightarrow H_m$ defined by

$$A_m = \left(\sigma(s) \frac{d}{ds} + \frac{\sigma(s)}{\rho(s)} \right) p_m \quad A_m^+ = \left(\sigma(s) \frac{d}{ds} + \frac{\sigma(s)}{\rho(s)} \right) (m-1) \rho(s) p_{m+1} \quad (8)$$

satisfy the relations $A_m p_m = p_{m+1}$ and $A_m^+ p_{m+1} = (1-m) p_m$ (see figure 1).




Figure 1. The functions ψ_{lm} satisfy the relation $H_m \psi_{lm} = \epsilon_{lm} \psi_{lm}$, and are related (up to some multiplicative constants) through the operators $A_m, A_m^+, a_m, a_m^+, U_m$ and $U_m^{-1} = U_m^+$.

Since $\psi_{lm}(s) = \frac{(m)!}{l!} (s)_k^{(m+1)} (s)$ is a polynomial and the function $\psi_{lm}(s)$ satisfies (4), integrating by parts one obtains $\langle \psi_{lm} | A_m^+ \psi_{lm+1} \rangle = \langle \psi_{lm} | A_m \psi_{lm+1} \rangle$, that is, the operators A_m and A_m^+ are mutually adjoint. From the relation

$\langle \psi_{lm+1} | \psi_{lm+1} \rangle = \langle \psi_{lm+1} | A_m^+ \psi_{lm+1} \rangle = \langle \psi_{lm+1} | A_m \psi_{lm+1} \rangle = \langle \psi_{lm+1} | \psi_{lm+1} \rangle$ it follows that $\epsilon_{lm} > \epsilon_{lm+1}$ for all $l > m$, and $\langle \psi_{lm+1} | \psi_{lm+1} \rangle = \langle \psi_{lm} | \psi_{lm} \rangle$. This is possible only if $\epsilon_{lm}(s) = 0$ and $\epsilon_{lm}(s) < 0$. Particularly, we have $\epsilon_{lm} \neq \epsilon_{lk}$ if and only if $l \neq k$.

The operators $H_m : H_m \rightarrow H_m$ are self-adjoint, admit the factorizations

$$H_m = A_m^+ A_m, \quad H_{m+1} = A_m A_m^+ \quad (9)$$

and satisfy the intertwining relations $H_m A_m^+ = A_m^+ H_{m+1}, A_m H_m = H_{m+1} A_m$:

4. Creation and annihilation operators. Coherent states

For each $m \in \mathbb{N}$, the sequence $\psi_{j_1 m} > \psi_{j_1+1 m} > \psi_{j_1+2 m} > \dots$, where $j_1 m > \epsilon_{j_1 m} = \langle \psi_{j_1 m} | \psi_{j_1 m} \rangle$ is an orthonormal basis of H_m , and $U_m : H_m \rightarrow H_m, U_m \psi_{j_1 m} = \psi_{j_1+1 m}$ is a unitary operator. The mutually adjoint operators (see Figure 1) $a_m, a_m^+ : H_m \rightarrow H_m, a_m = U_m^+ A_m, a_m^+ = A_m^+ U_m$ satisfy the relations

$$a_m \psi_{j_1 m} = \epsilon_{j_1 m} \psi_{j_1-1 m}, \quad a_m^+ \psi_{j_1 m} = \epsilon_{j_1+1 m} \psi_{j_1+1 m} \quad (10)$$

and allow us to factorize H_m as $H_m = a_m^+ a_m$:

The Lie algebra L_m generated by a_m^+, a_m is isomorphic to $su(1,1)$ if $\epsilon_{lm} < 0$, and it is isomorphic to the Heisenberg-Weyl algebra $h(2)$ if $\epsilon_{lm} = 0$:

Let $m \in \mathbb{N}$ be a fixed natural number, and let $j_n = j_1 + n, e_n = \epsilon_{j_n m}, e_0 = 1, e_n = e_1 e_2 \dots e_n$. Since $0 = e_0 < e_1 < e_2 < \dots < e_n < \dots$ and $a_m j_n = \epsilon_{j_n} j_n - 1, a_m^+ j_n = \epsilon_{j_n+1} j_n + 1, (H_m - e_n) j_n = e_n j_n$ (11)

we can define a system of coherent states by using the general setting presented in [1].

If $R = \limsup_{n \rightarrow \infty} \frac{1}{n!} \epsilon_n \neq 0$ then we can define

$$z_j = \frac{1}{N} \sum_{n=0}^{\infty} \frac{z^n}{n!} j_n \quad \text{where} \quad N (z_j)^2 = \sum_{n=0}^{\infty} \frac{z_j^{2n}}{n!} \quad (12)$$

for any z in the open disk $C(0; R)$ of center 0 and radius R . We get in this way a continuous family $f(z) = \sum_{j=0}^{\infty} |z_j\rangle \langle z_j| \in C(0; R)$ of normalized coherent states such that $a_m |z\rangle = z |z\rangle$:

5. Application to Schrodinger type operators

The problem of factorization of operators H_m is a very important one since it is directly related to the factorization of some Schrodinger type operators [2, 4]. If we use a change of variable $s = s(x)$ such that $ds = dx \sqrt{s(x)}$ or $ds = dx / \sqrt{s(x)}$ and define the new functions $\psi_m(x) = \sqrt{s(x)} \phi_m(s(x))$ then the relation $H_m \psi_m = E_m \psi_m$ becomes an equation of Schrodinger type

$$\frac{d^2}{dx^2} \psi_m(x) + V_m(x) \psi_m(x) = E_m \psi_m(x): \quad (13)$$

For example, by starting from the equation of Jacobi polynomials with $\alpha = 1=2$, $\beta = 1=2$, and using the change of variable $s(x) = \cos x$ we obtain the Schrodinger equation corresponding to the Poschl-Teller potential [1]

$$V_0(x) = \frac{1}{4} \frac{(1)}{\cos^2(x=2)} + \frac{(1)}{\sin^2(x=2)} - \frac{(1+1)^2}{4}: \quad (14)$$

If we choose the change of variable $s = s(x)$ such that $ds = dx \sqrt{s(x)}$, then the operators corresponding to A_m and A_m^+ are the adjoint conjugate operators

$$\begin{aligned} A_m &= [s(x)]^{1=2} A_m [s(x)]^{1=2} \frac{d}{ds} = \frac{d}{dx} + W_m(x) \\ A_m^+ &= [s(x)]^{1=2} A_m^+ [s(x)]^{1=2} \frac{d}{ds} = -\frac{d}{dx} + W_m(x) \end{aligned} \quad (15)$$

where the superpotential $W_m(x)$ is given by the formula

$$W_m(x) = \frac{(s(x))}{2(s(x))} - \frac{2m-1}{2(s(x))} \frac{d}{dx} (s(x)): \quad (16)$$

From the relations satisfied by A_m and A_m^+ we get the formulae

$$\frac{d^2}{dx^2} \psi_m(x) + V_m(x) \psi_m(x) = E_m \psi_m(x) \quad \frac{d^2}{dx^2} \psi_{m+1}(x) + V_{m+1}(x) \psi_{m+1}(x) = E_{m+1} \psi_{m+1}(x) \quad (17)$$

$$V_m(x) - E_m = W_m^2(x) - W_{m-1}(x) \quad V_{m+1}(x) - E_{m+1} = W_m^2(x) + W_m(x) \quad (18)$$

$$W_m(x) = \frac{W_{m-1}(x)}{W_m(x)} \quad V_m(x) = \frac{W_m^2(x)}{W_m(x)} + E_m \quad (19)$$

where the dot sign means derivative with respect to x .

If we choose the change of variable $s = s(x)$ such that $ds = dx / \sqrt{s(x)}$ then the formulae are very similar (only some signs are changed). In the case of Poschl-Teller potential (14) we get

$$W_0(x) = \frac{1}{2} \cot \frac{x}{2} - \tan \frac{x}{2}: \quad (20)$$

References

- [1] Antoine J-P, Gazeau J-P, Monceau P, Klauder J R and Penson K A 2001 J. Math. Phys. 42 2349-87
- [2] Cooper F, Khare A and Sukhatm U 1995 Phys. Rep. 251 267-385
- [3] Cotfas N 2002 J. Phys. A: Math. Gen. 35 9355-9365 (quant-ph/0206129)
- [4] Infeld L and Hull T E 1951 Rev. Mod. Phys. 23 21-68
- [5] Jafarizadeh M A and Fakhri H 1998 Ann. Phys., NY 262 260-76
- [6] Lorente M 2001 J. Phys. A: Math. Gen. 34 569-88
- [7] Nikiforov A F, Suslov S K and Uvarov V B 1991 Classical Orthogonal Polynomials of a Discrete Variable (Berlin: Springer) pp 2-17