

Spectrum of One-Dimensional Multiple Well Oscillators

H. A. Alhendi¹ and E. I. Lashin^{1,2}

¹ Department of Physics and Astronomy, College of Science,
King Saud University, Riyadh, Saudi Arabia

² Department of Physics, Faculty of Science,
Ain Shams University, Cairo, Egypt

March 1, 2022

We apply power series expansion to symmetric multi-well oscillators bounded by two infinite walls. The spectrum and expectation values obtained are compared with available exact and approximate values for the unbounded ones. It is shown that the method is capable of producing to a high accuracy the eigenvalues, eigenfunctions, and the expectation values $\langle x^{2k} \rangle$ of the corresponding unbounded ones as the separation between the two infinite walls becomes large.

PACS numbers: 03.65.Ge, 02.30.Hq

1 Introduction

Potential energy functions with degenerate classical minima separated by potential barriers are frequently encountered in many areas of physics, such as molecular physics, nonlinear physics, field theory, and cosmology. A widely investigated model, by variant methods ([1]–[19]), is a one-dimensional symmetric double-well potential, in which the degeneracy of the low-lying doublets is broken by quantum tunneling effects [20].

Recently, calculation of energy splitting of a one-dimensional symmetric multiwell potential energy functions has received a considerable interest ([21]–[27]). This interest stems from the fact that the topology of the multiwell potential is completely different from the double-well case. Each well, except the two outer ones, is surrounded by two barriers. In the case of odd number of wells, the positions of the minima of the potential are connected by the spatial inversion symmetry of the potential that leaves the position of the minimum at the origin invariant, for even number of wells, however, these minima are connected by the inversion symmetry. These topological properties are relevant for calculations based on path integral methods using instanton solutions [28, 29, 30].

In an earlier paper [19], we have applied the traditional power series expansion method with boundary conditions in the finite interval to one-dimensional anharmonic potential

functions having single and double-well. We have shown that the method is capable, even for a reasonable size of the interval, of producing to a high accuracy the eigenvalues and eigenfunctions of the corresponding unbounded ones. In particular for the deep double-well case, with potential function $V(x) = -\frac{1}{2}x^2 + x^4$, we have shown that the splitting of the energy levels of the ground and first excited states becomes apparent only after 26 significant digits for the mass parameter $m^2 = 25$, and for $m^2 = 35$ the splitting shows up after 42 significant digits, while for $m^2 = 50$ the splitting starts after 72 significant digits.

In the present paper we extend our previous calculation to multi-well potentials and compare with available exact and approximate values recently obtained by other methods.

The rest of the paper is organized as follows. In sec. 2 we apply the method of power series expansion to anharmonic multi-well potentials, and present our calculation of the eigenvalues, eigenfunctions, and expectation values of even powers of the x -coordinate and compare with that recently obtained by the improved Hill determinant method [21, 23], and with the exact values calculated by supersymmetric quantum mechanics [21]. It is shown that the method yields extremely very close results to the exact ones. Finally in sec. 3 we give our conclusion.

2 Calculation of energies and expectation values

In order to calculate the eigenvalues and eigenfunctions of the anharmonic multi-well potential bounded by infinitely high potential walls located at $x = \pm L$, one needs to solve the eigenvalue equation (in units $\hbar = 1$; $2m = 1$):

$$-\frac{d^2}{dx^2} \psi + E - V(x) \psi = 0; \quad (1)$$

with the boundary conditions $\psi(\pm L) = 0$. In the present work we consider potentials $V(x)$ of the form [22, 23, 27]:

$$V(x) = \sum_{k=1}^N b_{2k} x^{2k}; \quad (2)$$

Here the coefficients (b_{2m}) are real with b_{2N} being positive, and $N = 3; 4; 5$:

Making use of the power series expansion

$$\psi = \sum_{n=0}^{\infty} a_n x^n \quad (3)$$

in eq. (1) we obtain the following recurrence formula for the expansion coefficients:

$$a_n = \frac{E a_{n-2} + b_2 a_{n-4} + b_4 a_{n-6} + b_6 a_{n-8} + b_8 a_{n-10} + b_{10} a_{n-12}}{n(n-1)}; \quad n \in 0; 1$$

$$a_n = 0; \quad n < 0 \quad (4)$$

The symmetry of eq. (1) divides the solutions into two types even and odd. The even solutions can be obtained by imposing (ignoring normalization) $a_0 = 1; a_1 = 0$, while

the odd ones by imposing $a_0 = 0; a_1 = 1$. The energy eigenvalues (E) are obtained from the condition $\psi(L) = 0$ in both cases. Since we are dealing with potentials admitting power series expansion for $|x| < L$, the power series solutions of $\psi(x)$ are, according to a well known theorem in differential equations, convergent [31]. Here we present our calculation of the energy levels, wave functions and expectation values for the potentials given in eq. (2), using eqs.(3),(4) and the boundary condition at $x = L$.

Table 1 shows the first four energy eigenvalues of three-, four-, and five-well oscillators for the bounded potential-wells as compared with the obtained values reported in [22, 23] for the unbounded case. The obtained values are extremely very close to the exact ones.

Table 2 contains a comparison of the expectation values $\langle x^{2m} \rangle$; ($m = 1 \dots 5$) with the reported ones and the exact values obtained from the corresponding supersymmetric wave functions. For completeness we also present in Fig. 1 (A;C) the ground state wave functions for the potentials $A(x)$ and $C(x)$ respectively, and (B;D) are for the first excited state wave functions for the potentials $B(x)$ and $D(x)$ respectively. These coincide with the corresponding exact wave functions (ignoring normalization) obtained using supersymmetric quantum mechanics [21]:




Figure 1: These graphs show the ground state wave functions for the potentials $A(x)$ and $C(x)$, also showing the first excited state wave functions corresponding to the potentials $B(x)$ and $D(x)$ for the bounded case

| Potential | No. of wells | L | Our results | Results of supersymmetric quantum mechanics |
|-----------|--------------|---|---|---|
| A (x) | 3 | 4 | <u>2:000 000 000 000 000 000</u> <u>1:772 726 698 991 350 330</u> <u>2:078 279 891 768 595 361</u> <u>5:604 028 342 382 013 654</u> | 2 |
| B (x) | 3 | 4 | <u>9:001 720 238 527 719 715</u> <u>9:000 000 000 000 000 000</u> <u>0:639 394 262 865 333 280</u> <u>1:936 629 224 926 380 607</u> | 9 |
| C (x) | 3 | 3 | <u>0:375 000 000 000 000 000</u> <u>2:357 398 881 839 175 696</u> <u>6:988 755 014 467 723 300</u> <u>13:880 516 236 71388 190</u> | 0:375 |
| D (x) | 3 | 3 | <u>0:195 122 059 734 627 597</u> <u>1:125 000 000 000 000 000</u> <u>5:646 143 524 135 629 302</u> <u>12:384 746 745 98 872 574</u> | 1:125 |
| F (x) | 4 | 4 | <u>0:223 991 055 384 171 854</u> <u>0:083 481 557 863 966 793</u> <u>1:526 487 708 073 844 797</u> <u>3:971 174 256 474 939 459</u> | |
| G (x) | 4 | 3 | <u>0:096 291 946 230 649 098</u> <u>0:672 993 242 745 446 704</u> <u>3:111 022 328 724 771 653</u> <u>7:038 082 659 880 398 654</u> | |
| H (x) | 5 | 4 | <u>0:807 741 647 209 432 443</u> <u>3:277 946 311 571 061 982</u> <u>7:667 480 496 116 480 534</u> <u>13:578 984 131 990 285 801</u> | |

Table 1: The first four energy eigenvalues of three-, four-, and five-well oscillators: $A(x) = x^2 - 4x^4 + x^6$; $B(x) = 4x^2 - 6x^4 + x^6$; $C(x) = \frac{105}{64}x^2 - \frac{43}{8}x^4 + x^6 - x^8 + x^{10}$; $D(x) = \frac{169}{64}x^2 - \frac{59}{8}x^4 + x^6 - x^8 + x^{10}$; $F(x) = -x^2 + 2x^4 - 0.9x^6 + 0.1x^8$; $G(x) = -x^2 + 3x^4 - 2x^6 + 0.1x^{10}$; $H(x) = 2x^2 - 7.5x^4 + 5.5x^6 - 0.877x^8 + 0.04x^{10}$ as obtained in the present work for the bounded cases ($2L$ is the width of the well). The underlined numbers correspond to the unbounded ones [23].

| Potential and eigenstate | Expectation value | results | | |
|-----------------------------|--------------------------|--------------------|---------|-----------|
| | | Our results | in [22] | Exact |
| A (x); $\phi_0(x)$ | $\langle x^2 \rangle$ | 1:7042723043 | 1:704 | 1:7043 |
| | $\langle x^4 \rangle$ | 3:9085446087 | 3:908 | 3:9085 |
| | $\langle x^6 \rangle$ | 10:373497673 | 10:349 | 10:3735 |
| | $\langle x^8 \rangle$ | 30:518356869 | 30:212 | 30:5184 |
| | $\langle x^{10} \rangle$ | 97:343955598 | 93:210 | 97:3440 |
| B (x); $\phi_1(x)$ | $\langle x^2 \rangle$ | 3:1795525642 | 3:179 | 3:1796 |
| | $\langle x^4 \rangle$ | 11:0386576927 | 11:035 | 11:0387 |
| | $\langle x^6 \rangle$ | 41:0648544887 | 40:985 | 41:0649 |
| | $\langle x^8 \rangle$ | 161:8298653908 | 159:909 | 161:8299 |
| | $\langle x^{10} \rangle$ | 670:2814413720 | 625:164 | 670:82814 |
| C (x); $\phi_0(x)$ | $\langle x^2 \rangle$ | 0:45832470069 | 0:458 | 0:4583 |
| | $\langle x^4 \rangle$ | 0:43854420934 | 0:439 | 0:4385 |
| | $\langle x^6 \rangle$ | 0:54740034191 | 0:547 | 0:5474 |
| | $\langle x^8 \rangle$ | 0:79673314349 | 0:796 | 0:7967 |
| | $\langle x^{10} \rangle$ | 1:28945196690 | 1:288 | 1:2895 |
| D (x); $\phi_1(x)$ | $\langle x^2 \rangle$ | 0:956841751448 | 0:957 | 0:9568 |
| | $\langle x^4 \rangle$ | 1:194350514116 | 1:194 | 1:1944 |
| | $\langle x^6 \rangle$ | 1:738359600265 | 1:738 | 1:7384 |
| | $\langle x^8 \rangle$ | 2:8134027359576611 | 2:812 | 2:8134 |
| | $\langle x^{10} \rangle$ | 4:935043317278 | 4:930 | 4:9350 |

Table 2: Comparison of the expectation values $\langle x^{2m} \rangle$ of the supersymmetric potential A (x); B (x); C (x); and D (x) obtained in the present work for the bounded cases and that of ref. [22] for the unbounded ones along with the exact values obtained using supersymmetric quantum mechanics [21, 22]

$$\begin{aligned}
\psi_0(x) &= \exp(x^2 - x^4/4) && \text{for A (x)} \\
\psi_0(x) &= \exp(-3x^2/16 + x^2/8 - x^6/6) && \text{for C (x)} \\
\psi_1(x) &= x \exp(3x^2/2 - x^4/4) && \text{for B (x)} \\
\psi_1(x) &= x \exp(-3x^2/16 + x^4/8 - x^6/6) && \text{for D (x)}
\end{aligned}$$

These calculations show that even for a reasonable size of the interval the present method, beside being simple and direct is highly effective.

3 Conclusion

In this paper we have applied the method of power series expansion to a variant of one-dimensional multi-well potential functions bounded by two infinite walls. We have compared our calculation of the low-lying energy levels, wave functions, and expectation values with that obtained using improved Hill determinant and supersymmetric quantum mechanics. For all these potentials we have obtained results that are extremely very close to the exact ones. This is because the power series expansion in the finite range is convergent for potentials admitting power series expansion. It is easy to extend the present method to other multi-well polynomial potential functions with higher degrees and more degenerate classical minima.

Acknowledgement

This work was supported by research center at college of science, King Saud University under project number Phys=1423=02.

References

- [1] N .From an, P .O .From an, U .M yhm an, and R .Paulsson, Ann.Phys. (NY) 74, (1972) 314 .
- [2] E .M agyari, Phys.Lett. A 81, (1981) 116 .
- [3] J.D .de Deus, Phys.Rev. D 26, (1982) 2782 .
- [4] R .Balsa, M .P lo, J.G .Esteve, and A .F .Pochecco, Phys.Rev. D 28, (1983) 1945 .
- [5] C .S .H sue and J.L .Chem, Phys.Rev. D 29, (1984) 643 .
- [6] B .R .Holstein, Am .J.Phys. 56, (1988) 338 .
- [7] W .Y .Keung , E .K ovacs, and U .P .Sukhatm e, Phys.Rev.Lett. 60, (1988) 41 .
- [8] J.M .Cornwall and G .T iktopouls, Ann.Phys. (NY) 228, (1993) 365 .
- [9] F .A rias de Saavedra and E .Buendia, Phys.Rev. A 42, (1995) 5073 .
- [10] Yu.Zhou, J.D .M ancini, P .F .M eier, and S.P .Bowen, Phys.Rev. A 51, (1995) 3337 .
- [11] E .J .W eniger, Phys.Rev.Lett. 77, (1996) 2859 .
- [12] E .D elabere and F .Pham , Ann.Phys. (NY) 261, (1997) 180 .
- [13] R .Ferreira and G .Bastard, Rep.Prog.Phys. 60, (1997) 345 .
- [14] C .J .Tym czak, G .S .Japaridze, C .R .Handy, and X .-Q .W ang, Phys.Rev.Lett. 80, (1998) 3673 .
- [15] I .V .Chebotarev, Ann.Phys. (NY) 273, (1999) 114 .
- [16] P .B .Khan and Y .Zam i, J.M ath.Phys. 40, (1999) 4658 .
- [17] U .D .Jentschura and J .Zinn-Justin math-ph/0103010.
- [18] A .K .Roy, N .G upta, and B .M .Deb, Phys.Rev. A 65, (2002) 012109 .
- [19] H .A .A lhendi and E .I .Lashin quant-ph/0305128.
- [20] L .D .Landau and E .M .Lifshitz, Quantum mechanics (Pergam on Press, O xford, 1977), 3rd ed. pp.183-184.
- [21] R .A dhikari, R .D utt and Y .P .Vershni, Phys.Lett. A 141, (1989) 1 .
- [22] R .N .Chaudhuri and M .M ondal, Phys.Rev. A 43, (1991) 3241 .
- [23] R .K .A grawal and V .S .Varm a, Phys.Rev. A 49, (1994) 5089 .
- [24] Soo-Young Lee, et al Mod.Phys.Lett.A 12, (1997) 1803 .
- [25] J .Casahorran, Phys.Lett. A 283, (2001) 285 .

- [26] J. Casahorran, Fortsch. der Phys. 50, (2002) 405 .
- [27] N. Gupta, A. K. Roy, and B. M. Deb, Pramana. J. Phys. 59, (2002) 575 .
- [28] S. Coleman, " The Use of Instantons," in The W hys of Subnuclear Physics, proceeding of the International School of Subnuclear Physics, Erice, 1977, edited by A. Zichichi (Plenum , New York, 1979), pp. 270-274.
- [29] R. Rajaraman, Solutions and Instantons (North-Holland, the Netherlands, 1987).
- [30] H. Kleinert, Paths Integral in Quantum Mechanics, Statistics and Polymer Physics. Singapore: World Scientific (1990).
- [31] E. A. Coddington, An introduction to ordinary differential equations, (Prentice-Hall, Inc., Englewood Cliffs, N. J., 1966), pp. 138-142.