

# ANOTHER STATE ENTANGLEMENT MEASURE

Oscar A. Nagel<sup>1</sup>, and Guido A. Raggio<sup>2</sup>

Given a state  $\omega$  of the (minimal  $C^*$ -) tensor product  $A \otimes B$  of unital  $C^*$ -algebras  $A$  and  $B$ , its marginals are the states of  $A$  and  $B$  defined by

$$\omega^A(a) = \omega(a \otimes \mathbf{1}_B), \quad a \in A, \quad \omega^B(b) = \omega(\mathbf{1}_A \otimes b), \quad b \in B.$$

Given a state  $\rho$  of  $A$  and a state  $\phi$  of  $B$ , there is a unique state  $\omega$  of  $A \otimes B$  such that  $\omega(a \otimes b) = \rho(a)\phi(b)$  for all  $a \in A$  and all  $b \in B$ ; we denote this state by  $\rho \otimes \phi$ . A **product-state** of  $A \otimes B$  is a state  $\omega$  of  $A \otimes B$  such that  $\omega = \omega^A \otimes \omega^B$ . We write  $S_\pi(A \otimes B)$  for the product-states of  $A \otimes B$ . The convex hull of  $S_\pi$ , written  $co(S_\pi(A \otimes B))$ , is the set of finite convex combinations of product states. The states of  $A \otimes B$  in the norm-closure of  $co(S_\pi(A \otimes B))$  are usually identified with the **separable** states of the composite system whose observables are described by  $A \otimes B$ ; the states which are not separable are termed **entangled**.

For a state  $\omega$  of a unital  $C^*$ -algebra  $A$ , consider its finite convex decompositions:  $\omega = \sum_{j=1}^n \lambda_j \omega_j$ , with  $0 \leq \lambda_j \leq 1$ ,  $\sum_{j=1}^n \lambda_j = 1$ , and  $\omega_j$  a state of  $A$ . Such a decomposition will be written  $[\lambda_j, \omega_j]$  and  $\mathcal{D}_\omega$  denotes all such finite convex decompositions.

Consider the relative entropy  $(\rho, \phi) \rightarrow S(\rho, \phi)$  for pairs of states  $\rho$  and  $\phi$  of a unital  $C^*$ -algebra. We use the original convention of Araki [1]<sup>3</sup>, which is also that used in [2] which we use as a standard reference for the properties of relative entropy. We propose the following measure of entanglement

$$(1) \quad E(\omega) = \inf_{[\lambda_j, \omega_j] \in \mathcal{D}_\omega} \sum_{j=1}^n \lambda_j S(\omega_j, \omega_j^A \otimes \omega_j^B).$$

We say a map  $\alpha$  from  $A \otimes B$  into  $C \otimes D$  **commutes with marginalization** if for every state  $\omega$  of  $C \otimes D$  one has  $(\omega \circ \alpha)^A \otimes (\omega \circ \alpha)^B = (\omega^C \otimes \omega^D) \circ \alpha$ .

We have the following result, whose proof will be provided in a forthcoming paper [3], along with result about a class of entanglement measures akin to (1):

1.  $0 \leq E(\omega) \leq S(\omega, \omega^A \otimes \omega^B)$  with equality in the right-hand side inequality if  $\omega$  is a pure state.  $E(\omega) = 0$  if  $\omega$  is a product-state.
2.  $E(\cdot)$  is convex (and in general not affine).
3. If  $\alpha$  and  $\beta$  are, respectively,  $*$ -isomorphisms of  $A$  onto  $C$  and of  $B$  onto  $D$  ( $A, B, C$  and  $D$  are unital  $C^*$ -algebras) then  $E(\omega \circ (\alpha \otimes \beta)) = E(\omega)$  for every state of  $C \otimes D$ .

<sup>1</sup>Depto. de Física, Universidad Nacional del Sur, Bahía Blanca, Argentina. e-mail: nagel@uns.edu.ar

<sup>2</sup>FaMAF-Universidad Nacional de Córdoba, Córdoba, Argentina. e-mail: raggio@famaf.unc.edu.ar

<sup>3</sup>If  $A$  is the algebra of bounded linear operators on a Hilbert space, then  $S(\rho, \phi) = \text{Tr}(D_\rho(\log(D_\rho - \log(D_\phi))))$ , for normal states, where  $D_\rho$  (resp.  $D_\phi$ ) is the density operator for which  $\rho(a) = \text{Tr}(D_\rho a)$ ,  $a \in A$ .

4. If  $\gamma : A \otimes B \rightarrow C \otimes D$  is a unital, linear, continuous, Schwarz-positive map ( $\gamma(z^*z) \geq \gamma(z)^*\gamma(z)$  for every  $z \in A \otimes B$ ) which commutes with marginalization, then  $E(\omega \circ \gamma) \leq E(\omega)$  for every state  $\omega$  of  $C \otimes D$ .
5. If  $\omega$  is separable then  $E(\omega) = 0$ .
6.  $E(\omega) = 0$  iff  $\omega$  lies in the  $w^*$ -closure of  $\text{co}(S_\pi(A \otimes B))$ .
7. For  $n$  ( $n \geq 1$ ) states  $\omega_1, \omega_2, \dots, \omega_n$  of  $A \otimes B$ ,

$$(2) \quad E((\omega_1 \otimes \omega_2 \otimes \dots \otimes \omega_n) \circ \zeta_n) \leq \sum_{j=1}^n E(\omega_j) ,$$

where  $\zeta_n$  is the  $*$ -isomorphism

$$(3) \quad \left\{ \underbrace{A \otimes A \otimes \dots \otimes A}_n \right\} \otimes \left\{ \underbrace{B \otimes B \otimes \dots \otimes B}_n \right\} \xrightarrow{\zeta_n} \underbrace{(A \otimes B) \otimes (A \otimes B) \otimes \dots \otimes (A \otimes B)}_n,$$

given by  $\zeta_n((a_1 \otimes a_2 \otimes \dots \otimes a_n) \otimes (b_1 \otimes b_2 \otimes \dots \otimes b_n)) = (a_1 \otimes b_1) \otimes (a_2 \otimes b_2) \otimes \dots \otimes (a_n \otimes b_n)$ .

One has,

$$(4) \quad \lim_{n \rightarrow \infty} n^{-1} E((\omega \otimes \omega \otimes \dots \otimes \omega) \circ \zeta_n) \leq E(\omega) .$$

In both (2) and (4), the left-hand side is computed with respect to marginalization with respect to the two factors in  $\{ \}$ -brackets in (3).

8. If  $A$  or  $B$  is abelian then  $E \equiv 0$ .
9. Let  $\mathcal{M}_\omega$  be the (Radon)-measures on the state space with barycenter  $\omega$ , then

$$E(\omega) = \inf_{\{\mu \in \mathcal{M}_\omega\}} \int \mu(d\phi) S(\phi, \phi^A \otimes \phi^B) ,$$

and there exists  $\mu_o \in \mathcal{M}_\omega$  such that

$$E(\omega) = \int \mu_o(d\phi) S(\phi, \phi^A \otimes \phi^B) .$$

—

The crucial condition of “commutation with marginalization” involved in property 4. of  $E$  is met by the “LQCC” maps considered in [4]. “LQCC” means “local quantum operations” with “classical communication”, and these are the relevant maps in the games that Alice and Bob play.

Like most known entanglement measures (see e.g., [4,5]), except that devised by Vidal and Werner [6], the calculation of  $E$  involves an infimum over a rather unmanageable set. Using Kosaki’s variational expression ([7]) for the relative entropy, one obtains a lower bound on  $E$  which can be possibly used to devise a strategy to show that  $E(\omega) > 0$  for a specific state  $\omega$ .

One can replace the relative entropy in the definition of  $E$  by other, suitable functions, e.g.  $\|\phi - \phi^A \otimes \phi^B\|$ , without losing the basic properties of  $E$ ; this is studied in [3].

In a previous version of this announcement [8], we claimed that  $E$  was additive, that is, equality holds in (2). We withdraw this claim because we have found a mistake in our “proof”.

## REFERENCES

1. H. Araki: Relative entropy for states of von Neumann algebras I [& II]. Publ. RIMS Kyoto Univ. 11, 809-833 (1976); [ & 13: 173-192 (1977)].
2. M. Ohya, and D. Petz: Quantum Entropy and Its Use. Springer-Verlag Berlin Heidelberg 1993.
3. O.A. Nagel, and G.A. Raggio: A family of state entanglement measures. Work in progress.
4. M.J. Donald, M. Horodecki, and O. Rudolph: The uniqueness theorem for entanglement measures. J. Math. Phys. 43, 4252-4272 (2002).
5. V. Vedral: The role of relative entropy in quantum information theory. Rev. Mod. Phys. 74, 197-234 (2002).
6. G. Vidal, and R.F. Werner: A computable measure of entanglement. Phys. Rev. A 65, 032314 (2002).
7. H. Kosaki: Relative entropy for states: a variational expression. J. Operator Th. 16, 335-348 (1986).
8. O.A. Nagel and G.A. Raggio: Another state entanglement measure. quant-ph/0306024 v2.