arXiv:quant-ph/0307040v2 22 Jul 2003

Decoherence free algebra

Yoshiko Ogata

Abstract

We consider the decoherence free subalgebra which satisfies the minimal condition introduced by Alicki [1]. We show the manifest form of it and relate the subalgebra with the Kraus representation. The arguments also provides a new proof for generalized Lüders theorem [5].

1 Introduction

Decoherence is a non-unitary dynamics that is a consequence of system- environment coupling. As a way to protect a quantum system against decoherence, a subalgebra whose dynamics is implementable by unitary operator has attracted considerable interests; that is, the decoherence free subalgebra.

Let M_n be a C^*-algebra of all $n \times n$ matrices and ϕ a unital trace preserving completely positive map from M_n into itself. It is known that ϕ satisfies the Kadison inequality:

$$\phi(x^*x) \geq \phi(x)^*\phi(x) \quad \text{for all } x \in M_n. \tag{1}$$

The equality holds for all $x \in M_n$ if and only if ϕ is given by a unitary operator U as $\phi(x) = U^*xU$. So $\phi(x^*x) - \phi(x)^*\phi(x)$ can be interpreted as a dissipation function. Under these considerations, Alicki introduced the *decoherence free subalgebra* N_ϕ of ϕ as [1]

$$N_\phi = \{x_1 + ix_2 \in M_n; x_i = x_i^*, \phi(x_i^2) = \phi(x_i)^2,\ i = 1, 2\}.$$

It is a C^*-subalgebra of M_n. Suppose that ϕ can be joined to the identity map id, in the sense that there exists a continuous map from interval $[0, 1]$ to completely positive map $t \to \phi_t$ which satisfies $\phi_0 = \mathrm{id}$, and $\phi_1 = \phi$. Then there exists a unitary operator U_ϕ in M_n such that

$$\phi(x) = U_\phi^* x U_\phi \quad \text{for all } x \in N_\phi.$$

Hence it seems suitable to call N_ϕ *decoherence free algebra*. However, how can we find the manifest form of this subalgebra? In particular, how is it related to the Kraus representation $\phi(x) = \sum_{i=1}^k A_i^* x A_i$? In this paper, we consider this problem and show the following proposition:

Proposition 1.1 *The decoherence free subalgebra N_ϕ is equal to the commutant of the algebra generated by $A_i A_j^*$;*

$$N_\phi = \{A_i A_j^*, for\ all\ 1 \leq i, j \leq k\}',$$

which is independent of the choice of the Kraus representation.

Here and hereafter, we denote the commutant of $\mathcal{C}$ by $\mathcal{C}' \equiv \{x \in M_n; xy = yx\ for\ all\ y \in \mathcal{C}\}$. The arguments also provides a new proof for generalized Lüders theorem [5].

2 The decoherence free algebra

Let us first confirm the notations. We denote by $B(\mathcal{H})$ the set of all bounded operators on Hilbert space $\mathcal{H}$. A subset S of Hilbert space $\mathcal{H}$ is said to be total if it spans $\mathcal{H}$. We denote by $\mathfrak{h}_m$ the m-dimensional Hilbert space, and denote the orthonormal basis of it as $\{\mathrm{e}_i; 1 \leq i \leq m\}$. A linear map ϕ from $B(\mathcal{H}_2)$ to $B(\mathcal{H}_1)$ is called a unital completely positive map if it satisfies (i)$\phi(1) = 1$ and (ii)for every $n = 1, 2, \cdots$ the correspondence $(X_{ij}) \to (\phi(X_{ij}))\ 1 \leq i, j \leq n$ from $B(\mathcal{H}_2 \otimes \mathbb{C}^n)$ to $B(\mathcal{H}_1 \otimes \mathbb{C}^n)$ preserves positivity.

Before going into the proof, we introduce two famous Theorems. The first one is the Theorem of Choi [2];

Theorem 2.1 *Suppose that ϕ is a unital completely positive map from $B(\mathcal{H}_2)$ to $B(\mathcal{H}_1)$. If $\phi(a^*a) = \phi(a)^*\phi(a)$ and $\phi(aa^*) = \phi(a)\phi(a)^*$ for some $a \in B(\mathcal{H}_2)$, then for all $b \in B(\mathcal{H}_2)$, we have $\phi(ba) = \phi(b)\phi(a), \phi(ab) = \phi(a)\phi(b)$.*

The second one is the Theorem of Steinspring [3];

Theorem 2.2 *Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be finite dimensional Hilbert spaces and let ϕ be a unital completely positive map from a $B(\mathcal{H}_2)$ to $B(\mathcal{H}_1)$. Then there exists a Hilbert space $\mathcal{K}$ and isometry V from $\mathcal{H}_1$ to $\mathcal{H}_2 \otimes \mathcal{K}$ such that the following holds:*

(a) $\phi(x) = V^*(x \otimes 1)V \quad for\ all\ x \in B(\mathcal{H}_2)$

(b) $\{(x \otimes 1)Vu \mid x \in B(\mathcal{H}_2), u \in \mathcal{H}_1\}$ *is total in $\mathcal{H}_2 \otimes \mathcal{K}$.*

(c) *If $\mathcal{K}'$ is another Hilbert space and V' is an isometry from $\mathcal{H}_1$ to $\mathcal{H}_2 \otimes \mathcal{K}'$ such that (a) and (b) hold, then there exists a unitary W from $\mathcal{H}_2 \otimes \mathcal{K}$ to $\mathcal{H}_2 \otimes \mathcal{K}'$ such that $W(x \otimes 1_{\mathcal{K}})V = (x \otimes 1_{\mathcal{K}'})V'$.*

What we consider is a unital completely positive map from $B(\mathcal{H})$ to $B(\mathcal{H})$, where $\mathcal{H}$ is n-dimensional Hilbert space and $B(\mathcal{H})$ is equal to M_n. Now let us proceed to the proof of the proposition.

Proof of Proposition1.1: Let $A_i \in B(\mathcal{H}), 1 \leq i \leq m$ be Kraus operators which give the Kraus representation of ϕ as

$$\phi(a) = \sum_{1=1}^{m} A_i^* a A_i.$$

Define a linear map V from $\mathcal{H}$ to $\mathcal{H} \otimes \mathfrak{h}_m$ by

$$Vu = \sum_{i=1}^{m} A_i u \otimes \mathrm{e}_i.$$

The adjoint of V satisfies

$$V^* \left(\sum_{i=1}^{m} \psi_i \otimes \mathrm{e}_i \right) = \sum_{i=1}^{m} A_i^* \psi_i,$$

and

$$V^*(a \otimes 1)V = \sum_{1=1}^{m} A_i^* a A_i = \phi(a).$$

Therefore, the condition (a) of the Steinspring's Theorem holds for the pair $(K \equiv \mathfrak{h}_m, V)$.

First we consider the case that the condition (b) is also satisfied. Note that by the definition of N_ϕ, and the Choi's Theorem, any element a of N_ϕ satisfies $\phi(a^*a) = \phi(a)^*\phi(a)$. Then again by Choi's Theorem, we have

$$\phi(ba) = \phi(b)\phi(a), \phi(ab) = \phi(a)\phi(b) \tag{2}$$

for all $a \in N_\phi$ and all $b \in B(\mathcal{H})$. The relation (2) is represented in the Steinspring representation as

$$VV^*(a \otimes 1_K)VV^*(b \otimes 1_K)Vu = VV^*(a \otimes 1_K)(b \otimes 1_K)Vu$$

for all $u \in \mathcal{H}$, $a \in N_\phi$ and $b \in B(\mathcal{H})$. As $\{(b \otimes 1_K)Vu | b \in B(\mathcal{H}), u \in \mathcal{H}\}$ is total in $\mathcal{H} \otimes K$ by condition (b), this implies

$$VV^*(a \otimes 1_K)VV^* = VV^*(a \otimes 1_K),$$

on $\mathcal{H} \otimes K$. The decoherence free algebra N_ϕ is involutive. So we also have

$$VV^*(a^* \otimes 1_K)VV^* = VV^*(a^* \otimes 1_K).$$

for all $a \in N_\phi$. Taking the adjoint of this, we have

$$VV^*(a \otimes 1_K)VV^* = (a \otimes 1_K)VV^*.$$

Combining these, we get

$$(a \otimes 1_K)VV^* = VV^*(a \otimes 1_K),$$

for all $a \in N_\phi$. As

$$VV^* = \sum_{i,j=1}^{m} A_i A_j^* \otimes |\mathrm{e}_i\rangle\langle \mathrm{e}_j|,$$

we obtain

$$aA_iA_j^* = A_iA_j^*a \quad 1 \le i,j \le m \tag{3}$$

for all a in N_ϕ. Conversely, suppose that a self-adjoint element a in $B(\mathcal{H})$ satisfies (3). Then it follows that

$$\begin{aligned}\phi(a)^2 = \sum_{i,j=1}^{m} A_i^* a A_i A_j^* a A_j = \sum_{i,j=1}^{m} A_i^* A_i A_j^* a^2 A_j \\ = \phi(1)\phi(a^2) = \phi(a^2).\end{aligned}$$

Here we used the fact that ϕ is unital. Hence N_ϕ is equal to the commutant

$$N_\phi = \{A_iA_j^* \;\; 1 \le i,j \le k\}^{'}.$$

Second we consider the case that condition (b) of the Steinspring's Theorem fails. Let $\tilde{P}$ be a projection operator in $\mathcal{H} \otimes \mathfrak{h}_m$ onto the subspace spanned by $\{(x \otimes 1)Vu \mid x \in B(\mathcal{H}), u \in \mathcal{H}\}$. Note that $\tilde{P}$ satisfies

$$\tilde{P}(x \otimes 1)(y \otimes 1)Vu = (x \otimes 1)(y \otimes 1)Vu = (x \otimes 1)\tilde{P}(y \otimes 1)Vu.$$

This implies

$$\tilde{P}(x \otimes 1)\tilde{P} = (x \otimes 1)\tilde{P}, \quad \tilde{P}(x^* \otimes 1)\tilde{P} = (x^* \otimes 1)\tilde{P}, \quad x \in B(\mathcal{H}).$$

Taking the adjoint of the second equation, we obtain

$$(x \otimes 1)\tilde{P} = \tilde{P}(x \otimes 1), \quad x \in B(\mathcal{H}).$$

By Theorem of 5.9 of [4], all the bounded operator which commutes with $B(\mathcal{H}) \otimes 1$ belongs to $1 \otimes M_m$. So $\tilde{P}$ is of the form $\tilde{P} = 1 \otimes P$, where P is a projection in $\mathfrak{h}_m$. If (b) is not satisfied, $P \neq 1$. In this case, we define a new orthogonal basis f_j of $\mathfrak{h}_j$ so that there exists $1 \le l < m$, such that

$$\begin{aligned} Pf_j &= f_j \quad \text{for} \quad 1 \le j \le l \\ Pf_j &= 0 \quad \text{for} \quad l+1 \le j \le m. \end{aligned} \tag{4}$$

The orthonormal basis $\{e_i\}$ and $\{f_j\}$ are connected by a unitary transformation, as

$$e_i = \sum_{j=1}^{m} v_{i,j} f_j,$$

Define bounded operators B_j $1 \leq j \leq m$ as

$$B_j = \sum_{i=1}^{m} v_{i,j} A_i.$$

Using these relations, we obtain

$$Vu = \sum_{i=1}^{m} A_i u \otimes e_i = \sum_{j=1}^{m} B_j u \otimes f_j.$$

From $(1 \otimes (1-P))Vu = 0$ and (4), we have $B_j = 0$ for $l+1 \leq j \leq m$. We then see that condition (b) is satisfied if and only if A_i are linear independent. Now as $\{(x \otimes 1)Vu \mid x \in B(\mathcal{H}), u \in \mathcal{H}\}$ is total in $\mathcal{H} \otimes P\mathfrak{h}_m = \mathcal{H} \otimes \mathfrak{h}_l$, the pair $(\mathfrak{h}_l, V)$ satisfies conditions (a),(b) of the Stinespring's Theorem. So by the preceding arguments, we have

$$N_\phi = \{B_i B_j^* \;\; 1 \leq i, j \leq l\}'.$$

On the other hands, as $v_{i,j}$ satisfies

$$\sum_{j=1}^{m} \overline{v_{i,j}} v_{k,j} = \delta_{i,k},$$

we have

$$\sum_{j=1}^{m} \overline{v_{i,j}} B_j = \sum_{k,j=1}^{m} \overline{v_{i,j}} v_{k,j} A_k = A_i.$$

Hence we have

$$\{B_i B_j^* \;\; 1 \leq i, j \leq l\}' \supset \{A_i A_j^* \;\; 1 \leq i, j \leq m\}',$$

and

$$\{B_i B_j^* \;\; 1 \leq i, j \leq l\}' \subset \{A_i A_j^* \;\; 1 \leq i, j \leq m\}',$$

Therefore, we obtain

$$N_\phi = \{B_i B_j^* \;\; 1 \leq i, j \leq l\}' = \{A_i A_j^* \;\; 1 \leq i, j \leq m\}',$$

which completes the proof. □

Let us consider the relation between decoherence free algebra N_ϕ and the fixed points M_ϕ of ϕ defined by

$$M_\phi = \{x \in M_n | \phi(x) = x\}.$$

We have the following Lemma:

Lemma 2.1 *Let ϕ be a unital trace preserving completely positive map from M_n into itself.Then N_ϕ includes M_ϕ.*

Proof
Any $x \in M_\phi$ is decomposed into $x = x_1 + ix_2$, where x_1, x_2 are self adjoint elements of M_ϕ. So it suffices to show that any self-adjoint element x of M_ϕ satisfies $\phi(x^2) = \phi(x)^2$. By the inequality (1), we have

$$\phi(x^2) - x^2 \geq 0.$$

On the other hands, we have

$$\mathrm{Tr}\left(\phi(x^2) - x^2\right) = 0,$$

because ϕ is trace preserving. As trace is faithful, we have

$$\phi(x^2) = x^2 = \phi(x)^2,$$

which completes the proof. □

Let $\mathcal{A}$ be a C^*-algebra generated by A_i, and $\mathcal{B}$, a C^*-algebra generated by $A_i A_j^*$. Clearly, $\mathcal{B} \subset \mathcal{A}$ and the following inclusion relation holds:

$$\mathcal{A}' \subset M_\phi \subset N_\phi = \mathcal{B}'$$

If A_i are positive, then $A_i = \sqrt{A_i A_i^*} \in \mathcal{B}$, i.e., all the generators of $\mathcal{A}$ is included in $\mathcal{B}$. So we have $\mathcal{A} = \mathcal{B}$, which implies $\mathcal{A}' = \mathcal{B}'$. In this case, we have

$$\mathcal{A}' = M_\phi = N_\phi = \mathcal{B}'.$$

That is, all the decoherence free elements are fixed points if A_i are positive. This is a new proof of a generalized Lüders theorem [5], [6].

References

[1] Robert Alicki *Controlled Quantum Open Systems quant-ph/0302132*

[2] Choi, Man Duen *A Schwarz inequality for positive linear maps on C^*-algebras.* Illinois J. Math. **18** (1974), 565–574.

[3] Stinespring, W. Forrest *Positive functions on C^*-algebras.* Proc. Amer. Math. Soc. 6, (1955). 211–216.

[4] Takesaki, Masamichi *Theory of operator algebras. I.* Springer-Verlag, New York-Heidelberg, 1979.

[5] Paul Busch and Javed Singh *Lüders theorem for unsharp quantum measurements* Phys.Lett. A **249** (1998) 10-12

[6] A.Arias, A.Gheondea and S.Gudder *Fixed points of quantum operations* J.Math.Phys **43** No.12 (2002)