

Renormalisation and fixed points in Hilbert Space

Tarek Khalil and Jean Richert

Laboratoire de Physique Theorique, UMR 7085 CNRS/ULP,
Universite Louis Pasteur, 67084 Strasbourg Cedex,

France

(Dated: April 17, 2024)

The energies of low-lying bound states of a microscopic quantum many-body system of particles can be worked out in a reduced Hilbert space. We present here and test a specific non-perturbative truncation procedure. We also show that real exceptional points which may be present in the spectrum can be identified as fixed points of coupling constants in the truncation procedure.

PACS numbers: 03.65.-w 05.70.Fh 24.10.Cn

The *ab initio* construction of a rigorous quantum many-body theory able to describe bound particle systems like molecules, atoms, aggregates, atomic nuclei and condensed systems has developed over a long period of time starting in the sixties [1].

In practice the explicit resolution of the problem necessitates the diagonalisation of the many-body Hamiltonian in Hilbert space which is spanned by a complete set of basis states, in principle of infinite dimension, at least generally very large. In many cases the information of interest is however restricted to the knowledge of a few energetically low-lying states which often possess collective properties. Hence it would be convenient to work in a finite truncated subspace of the original Hilbert space. Many different procedures have been proposed and successfully applied in the framework of the nuclear shell model and related microscopic descriptions.

Rigorous projection methods lead to effective Hamiltonians which can in principle be explicitly generated by means of perturbation techniques [1, 2]. Unfortunately there exists no straightforward control on the convergence properties of the perturbation expansions which are involved, especially not when the interaction between the particles is strong as it is the case in atomic nuclei or in quantum spin systems for instance. Many attempts have been made in order to overcome this problem [3, 4, 5, 6]. Pragmatic phenomenological procedures have also been introduced [7, 8]. More recently effective 2-body interactions have been constructed by means of a non-perturbative renormalisation technique which cuts off the large momentum components of the interaction [9].

The investigations which follow are developed in the spirit of former work based on renormalization concepts [11, 12, 13, 14]. We introduce here a possible non-perturbative scheme for the bound state many-body problem which relies on a systematic reduction of Hilbert space.

Formal framework. Consider a system with a fixed but arbitrary number of bound quantum objects in a Hilbert space $H^{(N)}$ of dimension N governed by a Hamiltonian $H^{(N)}(g_1^{(N)}; g_2^{(N)}; \dots; g_p^{(N)})$ where

n
 $g_1^{(N)}; g_2^{(N)}; \dots; g_p^{(N)}$ are a set of parameters (coupling constants) which characterise $H^{(N)}$. The eigenvectors $j_i^{(N)}(g^{(N)})$ $i = 1; \dots; N$ of $H^{(N)}$ span the Hilbert space and are the solutions of the Schrodinger equation

$$H^{(N)}(g^{(N)})j_i^{(N)}(g^{(N)}) = \epsilon_i(g^{(N)})j_i^{(N)}(g^{(N)}) : (1)$$

The diagonalisation of $H^{(N)}$ delivers both the eigenvalues $\epsilon_i(g^{(N)})$; $i = 1; \dots; N$ and eigenvectors $j_i^{(N)}(g^{(N)})$; $i = 1; \dots; N$ in terms of a linear combination of orthogonal basis states f_j ; $j = 1; \dots; N$. Since $\dim H^{(N)} = N$ is generally very large if not infinite and the information needed reduces to a finite part of the spectrum it makes sense to try to restrict the space dimensions. If the relevant quantities of interest are for instance M eigenvalues out of the set $\epsilon_i(g^{(N)})$ then one looks for a new Hamiltonian $H^{(M)}(g^{(M)})$ such that

$$H^{(M)}(g^{(M)})j_i^{(M)}(g^{(M)}) = \epsilon_i(g^{(M)})j_i^{(M)}(g^{(M)}) : (2)$$

with the constraints

$$\epsilon_i(g^{(M)}) = \epsilon_i(g^{(N)}) : (3)$$

for $i = 1; \dots; M$. Eq. (3) implies relations between the sets of coupling constants $g^{(M)}$ and $g^{(N)}$

$$g_k^{(M)} = f_k(g_1^{(N)}; g_2^{(N)}; \dots; g_p^{(N)}) : (4)$$

with $k = 1; \dots; p$. The solution of this set of equations generates the effective Hamiltonian $H^{(M)}$ whose spectrum in the restricted space $H^{(M)}$ is the same as the corresponding one in the original space.

Renormalisation algorithm for systems at temperature $T = 0$. We develop an explicit but general approach which allows to implement the former procedure by following the evolution of the effective Hamiltonian of the system when the dimensions of the Hilbert space are systematically reduced. Using the Feshbach formalism [15, 16] we divide the Hilbert space $H^{(N)}$ into two subspaces, $P H^{(N)}$ and $Q H^{(N)}$ with

$$\dim P H^{(N)} = N - 1; \dim Q H^{(N)} = 1 : (5)$$

In the projected subspace $P H^{(N)}$ the system with energy E is described by the effective Hamiltonian

$$H_{\text{eff}}(E) = P H P + P H Q (E - Q H Q)^{-1} Q H P : \quad (6)$$

H is written in the general form

$$H = H_0 + g H_1 \quad (7)$$

where H_0 and H_1 are Hamiltonian operators and g a strength parameter (coupling constant) which takes the value $g^{(N)}$ in $H^{(N)}$.

The complete set of basis states $|j_i\rangle; i = 1; \dots; N$ may f.i. be chosen as the eigenvectors of H_0 with the corresponding eigenvalues $\epsilon_i; i = 1; \dots; N$. The expression $H_{\text{eff}}(E)$ is generally the starting point of theories which use perturbation expansions [1]. Here we proceed differently. We consider

$$P |j_1^{(N)}\rangle = \sum_{i=1}^N a_{1i}^{(N)} (g^{(N)}) |j_i\rangle \quad (8)$$

which is the projection on $P H^{(N)}$ of an eigenvector

$$|j_1^{(N)}\rangle = \sum_{i=1}^N a_{1i}^{(N)} (g^{(N)}) |j_i\rangle \quad (9)$$

of $H^{(N)}$. If $\epsilon_1^{(N)}$ is the eigenvalue corresponding to $|j_1^{(N)}\rangle$ we look for the solution of

$$H_{\text{eff}}(\epsilon_1^{(N)}) P |j_1^{(N)}\rangle = \epsilon_1^{(N)} P |j_1^{(N)}\rangle : \quad (10)$$

We consider $P |j_1^{(N)}\rangle$ to be the lowest energy eigenstate and $Q H^{(N)}$ to contain f. i. the highest one in energy. Any other state may in principle be chosen. We impose the lowest eigenvalue in the $P H^{(N)}$ subspace to be the same as the one in the complete space

$$\epsilon_1^{(N-1)} = \epsilon_1^{(N)} : \quad (11)$$

Projecting the expression of Eq. (10) on $|h_{1j}\rangle$ which is the eigenvector of H_0 with lowest energy

$$h_{1j} H_{\text{eff}}(\epsilon_1^{(N)}) P |j_1^{(N)}\rangle = \epsilon_1^{(N)} (g^{(N)}) a_{1j}^{(N)} (g^{(N)}) : \quad (12)$$

and defining the Hamiltonian which acts in the projected space $P H^{(N)}$ as $H^{(N-1)} = H_0 + g^{(N-1)} H_1$, Eq. (11) leads to

$$h_{1j} H_{\text{eff}}(\epsilon_1^{(N)}) P |j_1^{(N)}\rangle = F(g^{(N-1)}) : \quad (13)$$

where

$$F(g^{(N-1)}) = H_{1N}^{(N-1)} + H_{1N}(\epsilon_1^{(N-1)}) H_{NN}^{(N-1)} H_{N1}^{(N-1)} : \quad (14)$$

with

$$H_{ij}^{(N-1)} = h_{ij} H^{(N-1)} |j\rangle \quad (15)$$

and

$$H_{1N}^{(N-1)} = h_{1j} H^{(N-1)} P |j_1^{(N)}\rangle : \quad (16)$$

$H_{N1}^{(N-1)}$ is the same expression as $H_{1N}^{(N-1)}$ with h_{1j} replaced by h_{Nj} .

The r.h.s. of Eq. (13) can be worked out explicitly. The denominator in the second term of $H_{\text{eff}}(\epsilon_1^{(N)})$ of Eq. (14) is a scalar quantity since $\dim Q H^{(N)} = 1$ where N designates the highest-lying energy eigenstate of H_0 , $|j_N\rangle$. Imposing the constraint introduced through Eq. (11) leads to a relation which fixes $g^{(N-1)}$. One gets explicitly

$$a^{(N-1)} g^{(N-1)^2} + b^{(N-1)} g^{(N-1)} + c^{(N-1)} = 0 \quad (17)$$

where

$$a^{(N-1)} = G_{1N} H_{NN} F_{1N} \quad (18)$$

with

$$H_{ij} = h_{ij} H |j\rangle \quad (19)$$

$$b^{(N-1)} = a_{11}^{(N)} H_{NN}(\epsilon_1^{(N-1)}) + F_{1N}(\epsilon_1^{(N-1)}) \quad (20)$$

$$c^{(N-1)} = a_{11}^{(N)}(\epsilon_1^{(N-1)})(\epsilon_1^{(N-1)} - \epsilon_N^{(N)}) \quad (21)$$

with

$$F_{1N} = \sum_{i=1}^N a_{1i}^{(N)} h_{1j} H |j_i\rangle : \quad (22)$$

and

$$G_{1N} = H_{1N} \sum_{i=1}^N a_{1i}^{(N)} h_{Nj} H |j_i\rangle : \quad (23)$$

Since Eq. (17) is non-linear in $g^{(N-1)}$ and has two solutions, $g^{(N-1)}$ is chosen as the one closest to $g^{(N)}$ by continuity. The process can be iterated step by step by projection from the space of dimension $N-1$ to $N-2$ and further, generating subsequently a succession of values of the strength parameter (coupling constant) $g^{(k)}$ at each iteration. At each step the projected wavefunction $P |j_1^{(k)}\rangle$ is obtained from $|j_1^{(k)}\rangle$ by elimination of a state $|j_k\rangle$. Taking the continuum limit for large N leads to a differential equation for g .

Example.

We consider a real symmetric tight-binding Hamiltonian with $H_0 = 0$, diagonal elements $h_{ij} H |j_i\rangle = \epsilon_i$ and non-diagonal ones $h_{ij} H |j_{i+1}\rangle = h_{ij} H |j_i\rangle = \epsilon_{ij}$ which is typical of strongly coupled systems. By essence, a reduction of Hilbert space by means of perturbation expansions is meaningless in this context. Starting

with an initial Hilbert space dimension N we apply the renormalisation procedure described above to g starting from an initial value $g^{(N)}$. The evolution of the lowest eigenvalues and the flow of g are shown in Table 1 for different values of g and N . Several conclusions can be drawn. First, as expected, the stability of the spectrum of low-lying states is the better the smaller the non-diagonal coupling between neighbouring states. Second, the stability increases when the initial dimension of the space increases. Third, the coupling constant may considerably change, decreasing in the present case. As a general prescription for the algorithm to work one should always keep the states which are strongly coupled those basis states which contribute in an essential way to the low-lying states of the physical spectrum.

N	n	g	1	2	3	4	5
10	10	20	0.81	3.17	6.90	11.69	17.15
	7	13.4	1.02	3.93	8.29	13.43	18.57
	5	8.18	1.10	4.09	8.18	12.27	15.26
20	20	20	0.22	0.89	1.98	3.47	5.34
	10	3.28	0.26	1.04	2.26	3.83	5.62
	5	1.13	0.30	1.13	2.25	3.38	4.21
30	30	20	0.10	0.41	0.92	1.62	2.51
	15	6.02	0.12	0.46	1.02	1.77	2.68
	5	1.03	0.14	0.52	1.03	1.55	1.93
50	50	20	0.04	0.15	0.34	0.60	0.94
	20	3.73	0.04	0.17	0.37	0.65	0.99
	5	0.38	0.05	0.19	0.38	0.57	0.71
20	20	1.0	0.01	0.04	0.10	0.17	0.27
	10	0.33	0.01	0.05	0.11	0.19	0.28
	5	0.11	0.015	0.06	0.11	0.17	0.21

Table 1.

Evolution of the coupling constant and the 5 lowest eigenvalues of the tight-binding matrix described in the text. Here $\beta = 1$, $\alpha = 0.5$. N is the initial space dimension, n the restricted dimension and g the running coupling constant.

Exceptional points and fixed points.

In the present renormalisation scheme exceptional points which generate divergences in perturbation expansions are related to the existence of fixed points of the coupling constant g .

It has been rigorously established that the eigenvalues $\epsilon_k(g)$ of $H(g) = H_0 + gH_1$ are analytic functions of g with only algebraic singularities [10, 17, 18]. They get singular at so called exceptional points $g = g_e$ which are first order branch points in the complex g -plane. Branch points appear if two (or more) eigenvalues get degenerate.

This can happen if g takes values such that $H_{kk} = H_{ll}$ where $H_{kk} = \langle k | H | k \rangle$ which corresponds to a so-called level crossing. As a consequence, if a level belonging to the PH subspace defined above crosses a level lying in the complementary QH subspace the perturbation development constructed from $H_{eff}(E)$ diverges [10]. Exceptional points are defined as the solutions of [18]

$$f(g_e) = \det[H(g_e) - (g_e)I] = 0 \quad (24)$$

and

$$\frac{df(g_e)}{dg} \Big|_{g=g_e} = 0 \quad (25)$$

where $f(g)$ is the secular determinant. It is now possible to show that exceptional points are connected to fixed points corresponding to $dg/dx = 0$ in specific cases. If $\epsilon_i(g)$ are the set of eigenvalues the secular equation can be written as

$$\prod_{i=1}^N (\epsilon_i - g) = 0 \quad (26)$$

Consider $g = g_p$ which satisfies Eq. (24). Then Eq. (25) can only be satisfied if there exists another eigenvalue $\epsilon_q = g_p$, hence if a degeneracy appears in the spectrum. This is the case at an exceptional point.

Going back to the algorithm described above consider the case where the fixed eigenvalue ϵ_1 gets degenerate with some other eigenvalue $\epsilon_i^{(k)}(g = g_e)$ at some step k in the space reduction process. Since ϵ_1 is constrained to be constant,

$$\epsilon_1^{(k)}(g_e) = \epsilon_1^{(l)}(g_e^0) \quad (27)$$

which is realised in any projected subspace of size k and l containing states $j_1 i$ and $j_l i$. Going over to the continuum limit for large values of N and considering the subspaces of dimension x and $x + dx$ in this limit one can write

$$\frac{d\epsilon_1}{dx} = 0 = \frac{d\epsilon_i(x)}{dx} \quad (28)$$

Consequently

$$\frac{d\epsilon_i}{dg_e} \frac{dg_e}{dx} = 0 \quad (29)$$

Due to the Wigner-Neumann avoided crossing rule the degeneracy of eigenvalues is generally not fulfilled for real values of the coupling constant and the derivative of ϵ_i with respect to g vanishes. There exist however specific situations, like systems with special symmetry properties [10, 21] or in finite systems [19] for which degeneracy for real g can occur. In these cases Eq. (29) is realised if

$$\frac{dg_e}{dx} = 0 \quad \text{and} \quad \frac{d\epsilon_i}{dg_e} \neq 0 \quad (30)$$

The second relation works if crossing takes place and g_c is a fixed point in the sense of renormalisation theory.

Eq. (30) shows the connection between exceptional and fixed points in the framework of the present approach. Ground state degeneracy due to level crossing is indeed a signature for the existence of phase transitions [20], perturbation expansions break down at these points. The ground state wavefunction changes its properties when the (real) coupling constant g crosses the exceptional point g_c . There the eigenstates exchange the main components of their projection on the set of basis states $|j_i\rangle; i=1; \dots; N$. It is worthwhile to notice that the same result is not restricted to the ground state, it is valid at any physical level crossing in which one of the eigenvalues stays constant for any value of the coupling constant. We have tested this property on several systems.

Example.

Consider an $N \times N$ matrix which corresponds to a highly degenerate and strongly coupled system located at a fixed point whose representative matrix possesses equal diagonal matrix elements ($H_{00} = 0; H_{ii} = 0.5$ in the present example) and equal non-diagonal elements ($H_{ij} = +0.5$). H_{ij} is defined in Eq. (19). The diagonalisation and the application of the Hilbert space reduction procedure leads to rigorously stable lowest eigenstates equal to 20 and the coupling constant which characterises the system stays at its initial value, here $g = 20$, for any value of N . The system is located at a fixed point corresponding to a state level crossing (exceptional point) as discussed above.

Conclusions.

In summary, we developed a non-perturbative effective theory of the bound state many-body quantum problem based on a reduction process of the dimensions of the initial Hilbert space. The central point concerns the renormalisation of the coupling constant which characterises the initial Hamiltonian under the constraint that the lowest eigenenergy which corresponds to the ground state of the system should not change. We showed the relationship which exists between exceptional points corresponding to level crossings in the spectrum where perturbation expansions break down and fixed points of the coupling constants which characterise phase transitions. The approach was tested and discussed on explicit examples of strongly coupled and highly degenerate systems encountered in condensed matter physics. The formalism can be applied to other physical quantum systems like

atoms, molecules, aggregates as well. It can be extended to systems characterised by several coupling constants and at finite temperature. Effective operators acting in reduced space can be worked out. We shall present these developments in forthcoming work.

One of us (J.R.) would like to thank J. Polonyi, H.A. Weidenmüller, J.M. Carmona, M. Henkel, D.W. Heiss and I. Rotter for their encouragements, comments, criticisms and advices.

-
- [1] B.H. Brandow, Rev. Mod. Phys. 39 (1967) 711
 - [2] T.H. Schucan and H.A. Weidenmüller, Ann. Phys. (N.Y.) 73 (1972) 108
 - [3] K. Suzuki and S.Y. Lee, Prog. Theor. Phys. vol. 64 (1980) 2091
 - [4] H. Kummel, K.H. Lühmann, J.G. Zabolitzky, Phys. Rep. 38C (1978) 1
 - [5] B.M. Hoffman, S.Y. Lee, J. Richert, H.A. Weidenmüller, and T.H. Schucan, Ann. Phys. (N.Y.) 85 (1974) 410
 - [6] W.C. Haxton and T. Luu, Phys. Rev. Lett. 89 (2002) 182503
 - [7] G.E. Brown and T.T.S. Kuo, Nucl. Phys. A 92 (1967) 481
 - [8] B.A. Brown and B.H. Wildenthal, Ann. Rev. Nucl. Sci. 32 (1982) 65
 - [9] S. Bogner, T.T.S. Kuo, L. Coraggio, A. Covello and N. Itaco, Phys. Rev. C 65 (2002) 051301 (R)
 - [10] T.H. Schucan and H.A. Weidenmüller, Ann. Phys. (N.Y.) 76 (1973) 483
 - [11] S.D. Glazek and Kenneth G. Wilson, Phys. Rev. D 57 (1998) 3558
 - [12] J. Rau, cond-mat/0110061
 - [13] H. Mueller, J. Piekarewicz and J.R. Shepard, nucl-th/0110061
 - [14] K.W. Becker, A. Huebsch and T. Sommer, cond-mat/0208351 (1999) 1514
 - [15] H. Feshbach, Nuclear Spectroscopy part B (1960), Academic Press
 - [16] C. Bloch and J. Horowitz, Nucl. Phys. 8 (1958) 91
 - [17] T. Kato, "Perturbation Theory for Linear Operators", Springer Verlag, Berlin, 1966
 - [18] W.D. Heiss, Phys. Rev. E 61 (2000) 929
 - [19] T.D. Schultz, D.C. Mattis and E.H. Lieb, Rev. Mod. Phys. 36 (1964) 856
 - [20] S. Sachdev, "Quantum Phase Transitions", Cambridge University Press, 1999
 - [21] E.A. Yuzbashyan, B.L. Altshuler and B.S. Shastry, J. Phys. A: Math. Gen. 35 (2002) 7525